



Turkana County, Kenya

Multiple Indicator Cluster Survey, 2013/14

Final Report

February, 2016



The Turkana County Multiple Indicator Cluster Survey (MICS) was carried out in 2013/14 by the Population Studies and Research Institute, University of Nairobi, in collaboration with Kenya National Bureau of Statistics, as part of the global MICS programme. Technical support was provided by the United Nations Children’s Fund (UNICEF). UNICEF provided financial support.

The global MICS programme was developed by UNICEF in the 1990s as an international household survey programme to support countries in the collection of internationally comparable data on a wide range of indicators on the situation of children and women. MICS surveys measure key indicators that allow countries to generate data for use in policies and programmes, and to monitor progress towards the Millennium Development Goals (MDGs) and other internationally agreed upon commitments.

Suggested citation:

Kenya National Bureau of Statistics, Population Studies and Research Institute and United Nations Children’s Fund. 2016. *Turkana County Multiple Indicator Cluster Survey 2013/14, Final Report*. Nairobi, Kenya: Kenya National Bureau of Statistics, Population Studies and Research Institute and United Nations Children’s Fund.

Summary Table of Survey Implementation and the Survey Population, Turkana County, 2013/14

Survey implementation			
Sample frame	National Sample Survey and Evaluation Programme V (NASSEP V)	Questionnaires	Household Women (age 15-49) Children under-five
Updated	November 2013		
Interviewer training	October 2013	Fieldwork	November 2013 to January 2014
Survey sample			
Households		Children under-five	
Sampled	1,680	Eligible	1,117
Occupied	1,379	Mothers/caretakers interviewed	1,067
Interviewed	1,277	Response rate (Percent)	95.5
Response rate (Percent)	92.6		
Women			
Eligible for interviews	1,300		
Interviewed	1,104		
Response rate (Percent)	84.9		

Survey population			
Average household size	5.2	Percentage of population living in	
Percentage of population under:		Urban areas	54.6
Age 5	16.6	Rural areas	45.4
Age 18	56.1		
Percentage of women age 15-49 years with at least one live birth in the last 2 years	35.1		

Housing characteristics	
Percentage of households with	
Electricity	10.8
Finished floor	16.3
Finished roofing	43.2
Finished walls	20.1
Mean number of persons per room used for sleeping	4.04

Household or personal assets	
Percentage of households that own	
A television	10.4
A refrigerator	2.7
Agricultural land	13.3
Farm animals/livestock	36.2
Percentage of households where at least a member has or owns a	
Mobile phone	45.4
Car or truck	1.9

Summary Table of Findings¹

Multiple Indicator Cluster Surveys (MICS) and Millennium Development Goals (MDG) Indicators, Turkana County MICS, 2013/14

NUTRITION			
Breastfeeding and infant feeding			
MICS Indicator	Indicator	Description	Value
2.5	Children ever breastfed	Percentage of women with a live birth in the last 2 years who breastfed their last live-born child at any time	97.2
2.6	Early initiation of breastfeeding	Percentage of women with a live birth in the last 2 years who put their last newborn to the breast within one hour of birth	54.0
2.7	Exclusive breastfeeding under 6 months	Percentage of infants under 6 months of age who are exclusively breastfed	68.5
2.8	Predominant breastfeeding under 6 months	Percentage of infants under 6 months of age who received breast milk as the predominant source of nourishment during the previous day	86.1
2.9	Continued breastfeeding at 1 year	Percentage of children age 12-15 months who received breast milk during the previous day	91.8
2.10	Continued breastfeeding at 2 years	Percentage of children age 20-23 months who received breast milk during the previous day	60.5
2.11	Median duration of breastfeeding	The age in months when 50 percent of children age 0-35 months did not receive breast milk during the previous day	21.8
2.12	Age-appropriate breastfeeding	Percentage of children age 0-23 months appropriately fed during the previous day	52.3
2.13	Introduction of solid, semi-solid or soft foods	Percentage of infants age 6-8 months who received solid, semi-solid or soft foods during the previous day	31.2
2.14	Milk feeding frequency for non-breastfed children	Percentage of non-breastfed children age 6-23 months who received at least 2 milk feedings during the previous day	(30.4)
2.15	Minimum meal frequency	Percentage of children age 6-23 months who received solid, semi-solid and soft foods (plus milk feeds for non-breastfed children) the minimum number of times or more during the previous day	18.1
2.16	Minimum dietary diversity	Percentage of children age 6-23 months who received foods from 4 or more food groups during the previous day	9.5
2.17a	Minimum acceptable diet	(a) Percentage of breastfed children age 6-23 months who had at least the minimum dietary diversity and the minimum meal frequency during the previous day	3.7
2.17b		(b) Percentage of non-breastfed children age 6-23 months who received at least 2 milk feedings and had at least the minimum dietary diversity not including milk feeds and the minimum meal frequency during the previous day	(1.7)
2.18	Bottle feeding	Percentage of children age 0-23 months who were fed with a bottle during the previous day	9.7
Salt iodization			
2.19	Iodized salt consumption	Percentage of households with salt testing 15 parts per million or more of iodate	63.4
Low-birthweight			
2.20	Low-birthweight infants	Percentage of most recent live births in the last 2 years weighing below 2,500 grams at birth	8.2

¹ See Appendix G for a detailed description of MICS indicators

2.21	Infants weighed at birth	Percentage of most recent live births in the last 2 years who were weighed at birth	38.8
------	--------------------------	---	------

CHILD HEALTH

Vaccinations

MICS Indicator	Indicator	Description	Value
3.1	Tuberculosis immunization coverage	Percentage of children age 12-23 months who received BCG vaccine by their first birthday	96.4
3.2	Polio immunization coverage	Percentage of children age 12-23 months who received the third dose of OPV vaccine (OPV3) by their first birthday	88.0
3.3	Diphtheria, pertussis and tetanus (DPT) immunization coverage	Percentage of children age 12-23 months who received the third dose of DPT vaccine (DPT3) by their first birthday	90.5
3.4	MDG 4.3 Measles immunization coverage	Percentage of children age 12-23 months who received measles vaccine by their first birthday	81.3
3.5	Hepatitis B immunization coverage	Percentage of children age 12-23 months who received the third dose of Hepatitis B vaccine (HepB3) by their first birthday	89.7
3.6	Haemophilus influenzae type B (Hib) immunization coverage	Percentage of children age 12-23 months who received the third dose of Hib vaccine (Hib3) by their first birthday	86.2
3.8	Full immunization coverage	Percentage of children age 12-23 months who received all vaccinations recommended in the national immunization schedule by their first birthday	70.3

Tetanus toxoid

3.9	Neonatal tetanus protection	Percentage of women age 15-49 years with a live birth in the last 2 years who were given at least two doses of tetanus toxoid vaccine within the appropriate interval prior to the most recent birth	44.7
-----	-----------------------------	--	------

Diarrhoea

-	Children with diarrhoea	Percentage of children under age 5 with diarrhoea in the last 2 weeks	17.4
3.10	Care-seeking for diarrhoea	Percentage of children under age 5 with diarrhoea in the last 2 weeks for whom advice or treatment was sought from a health facility or provider	70.1
3.11	Diarrhoea treatment with oral rehydration salts (ORS) and zinc	Percentage of children under age 5 with diarrhoea in the last 2 weeks who received ORS and zinc	27.3
3.12	Diarrhoea treatment with oral rehydration therapy (ORT) and continued feeding	Percentage of children under age 5 with diarrhoea in the last 2 weeks who received ORT (ORS packet, pre-packaged ORS fluid, recommended homemade fluid or increased fluids) and continued feeding during the episode of diarrhoea	43.2

Acute Respiratory Infection (ARI) symptoms

-	Children with ARI symptoms	Percentage of children under age 5 with ARI symptoms in the last 2 weeks	5.1
3.13	Care-seeking for children with ARI symptoms	Percentage of children under age 5 with ARI symptoms in the last 2 weeks for whom advice or treatment was sought from a health facility or provider	77.8
3.14	Antibiotic treatment for children with ARI symptoms	Percentage of children under age 5 with ARI symptoms in the last 2 weeks who received antibiotics	39.3

Solid fuel use

3.15	Use of solid fuels for cooking	Percentage of household members in households that use solid fuels as the primary source of domestic energy to cook	99.3
------	--------------------------------	---	------

Malaria / Fever			
MICS Indicator	Indicator	Description	Value
-	Children with fever	Percentage of children under age 5 with fever in the last 2 weeks	18.6
3.16a	Household availability of insecticide-treated nets (ITNs)	Percentage of households with (a) at least one ITN	36.6
3.16b		(b) at least one ITN for every two people	13.4
3.17a	Household vector control	Percentage of households (a) with at least one ITN or that have been sprayed by IRS in the last 12 months	37.3
3.17b		(b) with at least one ITN for every two people or that have been sprayed by IRS in the last 12 months	14.8
3.18	MDG 6.7	Children under age 5 who slept under an ITN	24.7
3.19		Population that slept under an ITN	53.7
3.20		Care-seeking for fever	66.9
3.21		Malaria diagnostics usage	45.4
3.22	MDG 6.8	Anti-malarial treatment of children under age 5	17.5
3.23		Treatment with Artemisinin-based Combination Therapy (ACT) among children who received anti-malarial treatment	(50.3)
3.24		Pregnant women who slept under an ITN	33.5
3.25		Intermittent preventive treatment for malaria during pregnancy	30.1

WATER AND SANITATION			
MICS Indicator	Indicator	Description	Value
4.1	MDG 7.8	Use of improved drinking water sources	71.9
4.2		Water treatment	3.3
4.3	MDG 7.9	Use of improved sanitation	12.3
4.4		Safe disposal of child's faeces	26.8
4.5		Place for handwashing	3.3
4.6		Availability of soap or other cleansing agent	33.8

REPRODUCTIVE HEALTH				
Contraception and unmet need				
MICS Indicator	Indicator	Description	Value	
-	Total fertility rate	Total fertility rate for women age 15-49 years	6.0	
5.1	MDG 5.4	Adolescent birth rate	Age-specific fertility rate for women age 15-19 years	101
5.2		Early childbearing	Percentage of women age 20-24 years who had at least one live birth before age 18	29.2
5.3	MDG 5.3	Contraceptive prevalence rate	Percentage of women age 15-49 years currently married or in union who are using (or whose partner is using) a (modern or traditional) contraceptive method	14.1
5.4	MDG 5.6	Unmet need	Percentage of women age 15-49 years who are currently married or in union who are fecund and want to space their births or limit the number of children they have and who are not currently using contraception	34.0
Maternal and newborn health				
5.5a	MDG 5.5	Antenatal care coverage	Percentage of women age 15-49 years with a live birth in the last 2 years who were attended during their last pregnancy that led to a live birth (a) at least once by skilled health personnel (b) at least four times by any provider	93.6
5.5b	MDG 5.5			44.0
5.6		Content of antenatal care	Percentage of women age 15-49 years with a live birth in the last 2 years who had their blood pressure measured and gave urine and blood samples during the last pregnancy that led to a live birth	70.2
5.7	MDG 5.2	Skilled attendant at delivery	Percentage of women age 15-49 years with a live birth in the last 2 years who were attended by skilled health personnel during their most recent live birth	35.1
5.8		Institutional deliveries	Percentage of women age 15-49 years with a live birth in the last 2 years whose most recent live birth was delivered in a health facility	34.8
5.9		Caesarean section	Percentage of women age 15-49 years whose most recent live birth in the last 2 years was delivered by caesarean section	2.3
Post-natal health checks				
5.10		Post-partum stay in health facility	Percentage of women age 15-49 years who stayed in the health facility for 12 hours or more after the delivery of their most recent live birth in the last 2 years	70.6
5.11		Post-natal health check for the newborn	Percentage of last live births in the last 2 years who received a health check while in facility or at home following delivery, or a post-natal care visit within 2 days after delivery	37.1
5.12		Post-natal health check for the mother	Percentage of women age 15-49 years who received a health check while in facility or at home following delivery, or a post-natal care visit within 2 days after delivery of their most recent live birth in the last 2 years	37.1

CHILD DEVELOPMENT				
MICS Indicator	Indicator	Description	Value	
6.1		Attendance to early childhood education	Percentage of children age 36-59 months who are attending an early childhood education programme	30.8
6.2		Support for learning	Percentage of children age 36-59 months with whom an adult has engaged in four or more activities to promote learning and school readiness in the last 3 days	52.9

CHILD DEVELOPMENT			
MICS Indicator	Indicator	Description	Value
6.3	Father's support for learning	Percentage of children age 36-59 months whose biological father has engaged in four or more activities to promote learning and school readiness in the last 3 days	0.8
6.4	Mother's support for learning	Percentage of children age 36-59 months whose biological mother has engaged in four or more activities to promote learning and school readiness in the last 3 days	3.1
6.5	Availability of children's books	Percentage of children under age 5 who have three or more children's books	0.3
6.6	Availability of playthings	Percentage of children under age 5 who play with two or more types of playthings	19.4
6.7	Inadequate care	Percentage of children under age 5 left alone or in the care of another child younger than 10 years of age for more than one hour at least once in the last week	54.1
6.8	Early child development index	Percentage of children age 36-59 months who are developmentally on track in at least three of the following four domains: literacy-numeracy, physical, social-emotional, and learning	50.1

LITERACY AND EDUCATION^A			
MICS Indicator	Indicator	Description	Value
7.1 MDG 2.3	Literacy rate among young people	Percentage of young women age 15-24 years who are able to read a short simple statement about everyday life or who attended secondary or higher education	54.7
7.2	School readiness	Percentage of children in first grade of primary school who attended pre-school during the previous school year	43.8
7.3	Net intake rate in primary education	Percentage of children of school-entry age who enter the first grade of primary school	33.6
7.4 MDG 2.1	Primary school net attendance ratio (adjusted)	Percentage of children of primary school age currently attending primary (primary 1-6; ISCED 1) or secondary school	62.4
7.S1	Primary school net attendance ratio (adjusted)	Percentage of children of primary school age currently attending primary (primary 1-8; national) or secondary school	65.4
7.5	Secondary school net attendance ratio (adjusted)	Percentage of children of secondary school age currently attending secondary (primary 7-8 included; ISCED) school or higher	36.8
7.S2	Secondary school net attendance ratio (adjusted)	Percentage of children of secondary school age currently attending secondary school (national) or higher	19.4
7.6 MDG 2.2	Children reaching last grade of primary	Percentage of children entering the first grade of primary school who eventually reach last grade (primary 6; ISCED)	98.4
7.S3	Children reaching last grade of primary	Percentage of children entering the first grade of primary school who eventually reach last grade (primary 8; national)	96.0
7.7	Primary completion rate	Number of children attending the last grade of primary school (excluding repeaters) divided by number of children of primary school completion age (age appropriate to final grade of primary school) (ISCED)	84.2
7.S4	Primary completion rate	Number of children attending the last grade of primary school (excluding repeaters) divided by number of children of primary school completion age (age appropriate to final grade of primary school) (national)	74.0

7.8		Transition rate to secondary school	Number of children attending the last grade of primary school during the previous school year who are in the first grade of secondary school during the current school year divided by number of children attending the last grade of primary school during the previous school year (ISCED)	96.4
7.9	MDG 3.1	Gender parity index (primary school)	Primary school net attendance ratio (adjusted) for girls divided by primary school net attendance ratio (adjusted) for boys (ISCED)	0.97
7.S5		Gender parity index (primary school)	Primary school net attendance ratio (adjusted) for girls divided by primary school net attendance ratio (adjusted) for boys (national)	0.97
7.10	MDG 3.1	Gender parity index (secondary school)	Secondary school net attendance ratio (adjusted) for girls divided by secondary school net attendance ratio (adjusted) for boys (ISCED)	1.00
7.S6		Gender parity index (secondary school)	Secondary school net attendance ratio (adjusted) for girls divided by secondary school net attendance ratio (adjusted) for boys (national)	0.97
<p>^A For Kenya, the International Standard Classification of Education (ISCED) 1997 classifies Primary 7 and 8 as Lower Secondary education. The indicators labelled ISCED calculates Primary School indicators based on Primary 1-6 only, whereas Primary 7 and 8 are included in Secondary School indicators. Those indicators labelled national and marked with S are based on the national education system, which includes Primary 7-8 in Primary School indicators.</p>				
(*) Figures that are based on less than 25 unweighted cases				

CHILD PROTECTION

Birth registration

MICS Indicator	Indicator	Description	Value
8.1	Birth registration	Percentage of children under age 5 whose births are reported registered	40.1

Child labour

8.2	Child labour	Percentage of children age 5-17 years who are involved in child labour	32.8
-----	--------------	--	------

Child discipline

8.3	Violent discipline	Percentage of children age 1-14 years who experienced psychological aggression or physical punishment during the last one month	64.3
-----	--------------------	---	------

Early marriage and polygyny

8.4	Marriage before age 15	Percentage of women age 15-49 years who were first married or in union before age 15	8.9
8.5	Marriage before age 18	Percentage of women age 20-49 years who were first married or in union before age 18	32.2
8.6	Young women age 15-19 years currently married or in union	Percentage of young women age 15-19 years who are married or in union	13.9
8.7	Polygyny	Percentage of women people age 15-49 years who are in a polygynous union	30.0
8.8a	Spousal age difference	Percentage of young women who are married or in union and whose spouse is 10 or more years older, (a) among women age 15-19 years, (b) among women age 20-24 years	(14.2)
8.8b			21.3

Female genital mutilation/cutting			
8.9	Approval for female genital mutilation/cutting (FGM/C)	Percentage of women age 15-49 years who state that FGM/C should be continued	5.9
8.10	Prevalence of FGM/C among women	Percentage of women age 15-49 years who report to have undergone any form of FGM/C	3.2
8.11	Prevalence of FGM/C among girls	Percentage of daughters age 0-14 years who have undergone any form of FGM/C, as reported by mothers age 15-49 years	1.9
Attitudes towards domestic violence			
8.12	Attitudes towards domestic violence	Percentage of women age 15-49 years who state that a husband is justified in hitting or beating his wife in at least one of the following circumstances: (1) she goes out without telling him, (2) she neglects the children, (3) she argues with him, (4) she refuses sex with him, (5) she burns the food	70.1
Children's living arrangements			
8.13	Children's living arrangements	Percentage of children age 0-17 years living with neither biological parent	17.7
8.14	Prevalence of children with one or both parents dead	Percentage of children age 0-17 years with one or both biological parents dead	18.4
8.15	Children with at least one parent living abroad	Percentage of children 0-17 years with at least one biological parent living abroad	1.0

HIV/AIDS AND SEXUAL BEHAVIOUR

HIV/AIDS knowledge and attitudes

MICS Indicator	Indicator	Description	Value
-	Have heard of AIDS	Percentage of women age 15-49 years who have heard of AIDS	98.2
9.1 MDG 6.3	Knowledge about HIV prevention among young people	Percentage of young women age 15-24 years who correctly identify ways of preventing the sexual transmission of HIV, and who reject major misconceptions about HIV transmission	34.7
9.2	Knowledge of mother-to-child transmission of HIV	Percentage of women age 15-49 years who correctly identify all three means of mother-to-child transmission of HIV	22.3
9.3	Accepting attitudes towards people living with HIV	Percentage of women age 15-49 years expressing accepting attitudes on all four questions toward women living with HIV	16.7

HIV testing

9.4	Women who know where to be tested for HIV	Percentage of women age 15-49 years who state knowledge of a place to be tested for HIV	88.1
9.5	Women who have been tested for HIV and know the results	Percentage of women age 15-49 years who have been tested for HIV in the last 12 months and who know their results	53.0
9.6	Sexually active young women who have been tested for HIV and know the results	Percentage of young women age 15-24 years who have had sex in the last 12 months, who have been tested for HIV in the last 12 months and who know their results	67.7

9.7	HIV counselling during antenatal care	Percentage of women age 15-49 years who had a live birth in the last 2 years and received antenatal care during the pregnancy of their most recent birth, reporting that they received counselling on HIV during antenatal care	83.6
9.8	HIV testing during antenatal care	Percentage of women age 15-49 years who had a live birth in the last 2 years and received antenatal care during the pregnancy of their most recent birth, reporting that they were offered and accepted an HIV test during antenatal care and received their results	83.7
Sexual behaviour			
9.9	Young women who have never had sex	Percentage of never married young women age 15-24 years who have never had sex	66.9
9.10	Sex before age 15 among young women	Percentage of young women age 15-24 years who had sexual intercourse before age 15	8.2
9.11	Age-mixing among sexual partners	Percentage of women age 15-24 years who had sex in the last 12 months with a partner who was 10 or more years older	17.6
9.12	Multiple sexual partnerships	Percentage of women age 15-49 years who had sexual intercourse with more than one partner in the last 12 months	1.8
9.13	Condom use at last sex among people with multiple sexual partnerships	Percentage of women age 15-49 years who report having had more than one sexual partner in the last 12 months who also reported that a condom was used the last time they had sex	(*)
9.14	Sex with non-regular partners	Percentage of sexually active young women age 15-24 years who had sex with a non-marital, non-cohabiting partner in the last 12 months	17.0
9.15	MDG 6.2 Condom use with non-regular partners	Percentage of young women age 15-24 years reporting the use of a condom during the last sexual intercourse with a non-marital, non-cohabiting sex partner in the last 12 months	31.2
Orphans			
9.16	MDG 6.4 Ratio of school attendance of orphans to school attendance of non-orphans	Proportion attending school among children age 10-14 years who have lost both parents divided by proportion attending school among children age 10-14 years whose parents are alive and who are living with one or both parents	1.19

ACCESS TO MASS MEDIA AND ICT

Access to mass media

MICS Indicator	Indicator	Description	Value
10.1	Exposure to mass media	Percentage of women age 15-49 years who, at least once a week, read a newspaper or magazine, listen to the radio, and watch television	3.2
Use of information/communication technology			
10.2	Use of computers	Percentage of young women age 15-24 years who used a computer during the last 12 months	11.2

10.3	Use of internet	Percentage of young women age 15-24 years who used the internet during the last 12 months	11.8
------	-----------------	---	------

SUBJECTIVE WELL-BEING

MICS Indicator	Indicator	Description	Value
11.1	Life satisfaction	Percentage of young women age 15-24 years who are very or somewhat satisfied with their life, overall	89.7
11.2	Happiness	Percentage of young women age 15-24 years who are very or somewhat happy	91.5
11.3	Perception of a better life	Percentage of young women age 15-24 years whose life improved during the last one year, and who expect that their life will be better after one year	62.9

TOBACCO AND ALCOHOL USE

Tobacco use

MICS Indicator	Indicator	Description	Value
12.1	Tobacco use	Percentage of women age 15-49 years who smoked cigarettes, or used smoked or smokeless tobacco products at any time during the last one month	20.7
12.2	Smoking before age 15	Percentage of women age 15-49 years who smoked a whole cigarette before age 15	0.0

Alcohol use

12.3	Use of alcohol	Percentage of women age 15-49 years who had at least one alcoholic drink at any time during the last one month	9.8
12.4	Use of alcohol before age 15	Percentage of women age 15-49 years who had at least one alcoholic drink before age 15	4.2

Table of Contents

Summary Table of Survey Implementation and the Survey Population, Turkana County, 2013/14.....	iii
Summary Table of Findings.....	iv
Table of Contents.....	xiii
List of Tables.....	xvi
List of Figures.....	xx
List of Abbreviations.....	xxi
Foreword.....	xxiii
Acknowledgements.....	xxv
Executive Summary.....	xxvi
1. Introduction.....	1
1.1 Background.....	1
1.2 Survey Objectives.....	3
2. Sample and Survey Methodology.....	4
2.1 Sample Design.....	4
2.2 Questionnaires.....	4
2.3 Training and Fieldwork.....	6
2.4 Data Processing.....	6
3. Sample Coverage and the Characteristics of Households and Respondents.....	7
3.1 Sample Coverage.....	7
3.2 Characteristics of Households.....	8
3.3 Characteristics of Female Respondents 15-49 Years of Age and Children Under-5 Years.....	11
3.4 Housing characteristics, asset ownership, and wealth quintiles.....	15
4. Nutrition.....	19
4.1 Birth Weight.....	19
4.2 Breastfeeding and Infant and Young Child Feeding.....	21
4.3 Salt Iodization.....	31
5. Child Health.....	33
5.1 Vaccinations.....	33
5.2 Neonatal Tetanus Protection.....	39
5.3 Care of Illness.....	40
5.3.1 Diarrhoea.....	42

5.3.2 Acute Respiratory Infections	50
5.3.3 Solid Fuel Use	51
5.3.4 Malaria/Fever	53
6. Water and Sanitation	69
6.1 Use of Improved Water Sources	69
6.2 Use of Improved Sanitation	75
6.3 Handwashing	85
7. Reproductive Health	90
7.1 Fertility	90
7.2 Contraception	95
7.3 Unmet Need	98
7.4 Antenatal Care	101
7.5 Assistance at Delivery	105
7.6 Place of Delivery	108
7.7 Post-natal Health Checks	109
8. Early Childhood Development	116
8.1 Early Childhood Care and Education	116
8.2 Quality of Care	117
8.3 Developmental Status of Children	123
9. Literacy and Education	126
9.1 Literacy among Young Women	126
9.2 School Readiness	127
9.3 Primary and Secondary School Participation	128
10. Child Protection	141
10.1 Birth Registration	141
10.2 Child Labour	144
10.3 Child Discipline	149
10.4 Early Marriage and Polygyny	153
10.5 Female Genital Mutilation/Cutting	159
10.6 Attitudes toward Domestic Violence	162
10.7 Children's Living Arrangements	165
11. HIV/AIDS and Sexual Behaviour	168
11.1 Knowledge about HIV Transmission and Misconceptions about HIV	168
11.2 Knowledge of mother-to-child HIV transmission (MTCT)	172
11.3 Accepting Attitudes toward People Living with HIV	174
11.4 Knowledge of a Place for HIV Counselling and Testing during Antenatal Care	175

11.5 Sexual Behaviour Related to HIV Transmission	178
11.6 HIV Indicators for Young Women	179
11.7 Orphans	185
12. Access to Mass Media and Use of Information/Communication Technology.....	187
12.1 Access to Mass Media	187
12.2 Use of Information/Communication Technology	188
13. Subjective well-being	190
14. Tobacco and Alcohol Use.....	195
14.1 Tobacco Use	195
14.2 Alcohol Use	197
Appendix A. Documents Reviewed	199
Appendix B. Education ISCED Tables.....	206
Appendix C. Sample Design	212
Appendix D. Estimates of Sampling Errors	216
Appendix E. List of Personnel Involved in the Survey	221
Appendix F. Data Quality Tables	223
Appendix G. Turkana County MICS5 Indicators: Numerators and Denominators	238
Appendix H. Turkana County MICS Questionnaires	251

List of Tables

Table HH.1: Results of household, women's, men's and under-5 interviews	7
Table HH.2: Household age distribution by sex.....	8
Table HH.3: Household composition	10
Table HH.4: Women's background characteristics.....	12
Table HH.5: Under-5's background characteristics.....	13
Table HH.6: Housing characteristics	14
Table HH.7: Household and personal assets	15
Table HH.8: Wealth quintiles	16
Table NU.1: Low birth weight infants	19
Table NU.2: Guiding Principles for Feeding children age 6-23 months	20
Table NU.3: Initial breastfeeding	21
Table NU.4: Breastfeeding	23
Table NU.5: Duration of breastfeeding.....	23
Table NU.6: Age-appropriate breastfeeding.....	24
Table NU.7: Introduction of solid, semi-solid, or soft foods.....	25
Table NU.8: Infant and young child feeding (IYCF) practices.....	25
Table NU.9: Bottle feeding.....	26
Table NU.10: Iodized salt consumption	28
Table CH.1: Vaccinations in the first years of life.....	31
Table CH.2: Vaccinations by background characteristics	34
Table CH.3: Neonatal tetanus protection.....	35
Table CH.4: Reported disease episodes	37
Table CH.5: Care-seeking during diarrhoea.....	38
Table CH.6: Feeding practices during diarrhoea	39
Table CH.7: Oral rehydration solutions, recommended homemade fluids, and zinc	40
Table CH.8: Oral rehydration therapy with continued feeding and other treatments.....	42
Table CH.9: Source of ORS and zinc.....	44
Table CH.10: Care-seeking for and antibiotic treatment of symptoms of acute respiratory infection (ARI)	45
Table CH.11: Knowledge of the two danger signs of pneumonia	46
Table CH.12: Solid fuel use	47
Table CH.13: Solid fuel use by place of cooking	48
Table CH.14: Household availability of insecticide treated nets and protection by a vector control method.....	50
Table CH.15: Access to an insecticide treated net (ITN) - number of household members	52
Table CH.16: Access to an insecticide treated net (ITN) - background characteristics	52
Table CH.17: Use of ITNs	53
Table CH.18: Children sleeping under mosquito nets.....	55
Table CH.19: Use of mosquito nets by the household population.....	57
Table CH.20: Care-seeking during fever	58
Table CH.21: Treatment of children with fever	59
Table CH.22: Diagnostics and anti-malarial treatment of children	60
Table CH.23: Source of anti-malarial	60
Table CH.24: Pregnant women sleeping under mosquito nets.....	61
Table CH.25: Intermittent preventive treatment for malaria	62

Table WS.1: Use of improved water sources	65
Table WS.2: Household water treatment	67
Table WS.3: Time to source of drinking water	68
Table WS.4: Person collecting water	68
Table WS.5: Types of sanitation facilities	70
Table WS.6: Use and sharing of sanitation facilities	72
Table WS.7: Drinking water and sanitation ladders	75
Table WS.8: Disposal of child's faeces	77
Table WS.9: Water and soap at place for handwashing	79
Table WS.10: Availability of soap or other cleansing agent	81
Table RH.1: Fertility rates	83
Table RH.2: Adolescent birth rate and total fertility rate	84
Table RH.3: Early childbearing	84
Table RH.4: Trends in early childbearing	86
Table RH.5: Use of contraception	88
Table RH.6: Unmet need for contraception	91
Table RH.7: Antenatal care coverage	93
Table RH.8: Number of antenatal care visits and timing of first visit	95
Table RH.9: Content of antenatal care	96
Table RH.10: Assistance during delivery and caesarean section	98
Table RH.11: Place of delivery	101
Table RH.12: Post-partum stay in health facility	102
Table RH.13: Post-natal health checks for newborns	103
Table RH.14: Post-natal health checks for mothers	104
Table RH.15: Post-natal health checks for mothers and newborns	105
Table CD.1: Early childhood education	108
Table CD.2: Support for learning	110
Table CD.3: Learning materials	113
Table CD.4: Inadequate care	114
Table CD.5: Early child development index	115
Table ED.1: Literacy (young women)	117
Table ED.2: School readiness	118
Table ED.3: Primary school entry	119
Table ED.4: Primary school attendance and out of school children	121
Table ED.5: Secondary school attendance and out of school children	124
Table ED.6: Children reaching last grade of primary school	126
Table ED.7: Primary school completion and transition to secondary school	127
Table ED.8: Education gender parity	128
Table ED.9: Out of school gender parity	128
Table ED.10: Summary of education indicators (ISCED)	130
Table CP.1: Birth registration	134
Table CP.2: Children's involvement in economic activities	136
Table CP.3: Children's involvement in household chores	137
Table CP.4: Child labour	138
Table CP.5: Child discipline	140
Table CP.6: Attitudes toward physical punishment	142
Table CP.7: Early marriage and polygyny (women)	145

Table CP.8:	Trends in early marriage (women).....	148
Table CP.9:	Spousal age difference	149
Table CP.10:	Female genital mutilation/cutting (FGM/C) among women	150
Table CP.11:	Female genital mutilation/cutting (FGM/C) among girls	151
Table CP.12:	Approval of female genital mutilation/cutting (FGM/C)	152
Table CP.13:	Attitudes toward domestic violence (women)	153
Table CP.14:	Children's living arrangements and orphanhood.....	155
Table CP.15:	Children with parents living abroad.....	156
Table HA.1:	Knowledge about HIV transmission, misconceptions about HIV, and comprehensive knowledge about HIV transmission (women).....	159
Table HA.2:	Knowledge of mother-to-child HIV transmission (women)	162
Table HA.3:	Accepting attitudes toward people living with HIV (women).....	163
Table HA.4:	Knowledge of a place for HIV testing (women)	165
Table HA.5:	HIV counselling and testing during antenatal care	166
Table HA.6:	Sex with multiple partners (women)	167
Table HA.7:	Key HIV and AIDS indicators (young women)	169
Table HA.8:	Key sexual behaviour indicators (young women)	172
Table HA.9:	School attendance of orphans and non-orphans.....	175
Table MT.1:	Exposure to mass media (women).....	176
Table MT.2:	Use of computers and internet (women)	178
Table SW.1:	Domains of life satisfaction (women)	180
Table SW.2:	Overall life satisfaction and happiness (women)	182
Table SW.3:	Perception of a better life (women)	183
Table TA.1:	Current and ever use of tobacco (women)	185
Table TA.2:	Use of alcohol (women)	187
Appendices:		
Table ED.4:	Primary school attendance and out of school children	195
Table ED.5:	Secondary school attendance and out of school children	196
Table ED.7:	Primary school completion and transition to secondary school.....	198
Table ED.8:	Education gender parity.....	198
Table ED.9:	Out of school gender parity	200
Table SD.1:	Allocation of Sample Clusters (Primary Sampling Units) to Sampling Strata.....	202
Table SE.1:	Indicators selected for sampling error calculations.....	206
Table SE.2:	Sampling errors: Total sample	207
Table SE.3:	Sampling errors: Urban	208
Table SE.4:	Sampling errors: Rural.....	209
Table DQ.1:	Age distribution of household population	212
Table DQ.2:	Age distribution of eligible and interviewed women.....	214
Table DQ.4:	Age distribution of children in household and under-5 questionnaires.....	215
Table DQ.5:	Birth date reporting: Household population	215
Table DQ.6:	Birth date and age reporting: Women.....	216
Table DQ.8:	Birth date and age reporting: Under-5s.....	216

Table DQ.9: Birth date reporting: Children, adolescents and young people.....	216
Table DQ.10: Birth date reporting: First and last births.....	217
Table DQ.11: Completeness of reporting	217
Table DQ.12: Completeness of information for anthropometric indicators: Underweight	218
Table DQ.13: Completeness of information for anthropometric indicators: Stunting.....	218
Table DQ.14: Completeness of information for anthropometric indicators: Wasting	219
Table DQ.15: Heaping in anthropometric measurements.....	219
Table DQ.16: Observation of birth certificates	220
Table DQ.17: Observation of vaccination cards.....	221
Table DQ.18: Observation of women's health cards	221
Table DQ.19: Observation of bednets and places for handwashing	222
Table DQ.20: Presence of mother in the household and the person interviewed for the under-5 questionnaire	222
Table DQ.21: Selection of children age 1-17 years for the child labour and child discipline modules.....	223
Table DQ.22: School attendance by single age	224
Table DQ.23: Sex ratio at birth among children ever born and living	225
Table DQ.24: Births in years preceding the survey.....	225
Table DQ.25: Reporting of age at death in days	226
Table DQ.26: Reporting of age at death in months	226

List of Figures

Figure HH.1: Age and sex distribution of household population.....	9
Figure NU.1: Infant feeding patterns by age.....	22
Figure NU.2: Consumption of iodized salt	28
Figure CH.1: Vaccinations by age 12 months	32
Figure CH.2: Children under-5 with diarrhoea who received ORS or recommended homemade liquids	41
Figure CH.3: Children under-5 with diarrhoea receiving oral rehydration therapy (ORT) and continued feeding	43
Figure CH.4: Percentage of household population with access to an ITN in the household	53
Figure WS.1: Percent distribution of household members by source of drinking water.....	66
Figure WS.2: Percent distribution of household members by use and sharing of sanitation facilities	73
Figure WS.3: Use of improved drinking water sources and improved sanitation facilities by household members	76
Figure WS.4: Use of Improved water and sanitation in urban and rural areas.....	77
Figure RH.1: Differentials in contraceptive use	90
Figure RH.2: Person assisting at delivery	100
Figure RH.3: Place of delivery and post-natal health checks.....	106
Figure ED.1: Education indicators by sex.....	129
Figure CP.1: Children under-5 whose births are registered.....	135
Figure CP.2: Child disciplining methods, children age 1-14 years.....	142
Figure CP.3: Early marriage among women	149
Figure HA.1: Women with comprehensive knowledge of HIV transmission.....	161
Figure HA.2: Accepting attitudes toward people living with HIV/AIDS	164
Figure HA.3: Sexual behaviour that increases the risk of HIV infection, young people age 15-24.....	174
Figure TA.1: Ever and current smokers	186
Appendix:	
Figure DQ.1: Number of household population by single ages	213
Figure DQ.2: Weight and height/length measurements by digits reported for the decimal points.....	220

List of Abbreviations

ACRWC	African Charter on the Rights and Welfare of the Child
ACT	Artemisinin-based Combination therapy
AIDS	Acquired Immune Deficiency Syndrome
ANC	Antenatal Care
ARI	Acute Respiratory Infection
ART	Anti-retroviral Therapy
ASFRs	Age-specific Fertility Rates
BCC	Behaviour Change Communication
BCG	Bacillus Calmette-Guérin (Tuberculosis)
CARMMA	Campaign on Accelerated Reduction of Maternal Mortality in Africa
CBR	Crude Birth Rate
CEDAW	Convention on the Elimination of all forms of Discrimination Against Women
CRC	Convention on the rights of the Child
CSP	Country Strategy Paper
CSPro	Census and Survey Processing System
DOMC	Division of Malaria Control
DPT	Diphtheria Pertussis Tetanus
DVI	Division of Vaccine and Immunisation
EA	Enumeration area
ECD	Early Childhood Development
ECDE	Early Childhood Development and Development
ECDI	Early Child Development Index
EFA	Education for All
EHP	Essential Health Package
EMTCT	Elimination of Mother-to-Child Transmission of HIV
EPI	Expanded Programme on Immunization
FCTC	Framework Convention on Tobacco Control
FGM/C	Female genital mutilation/cutting
FNSP	Food and Nutrition Security Policy
GAPPD	Global Action Plan for the Prevention and Control of Pneumonia and Diarrhoea
GARPR	Global AIDS Response Progress Reporting
GFR	General Fertility Rate
GIPA	Greater Involvement of People Living with HIV and AIDS
GMAP	Global Malaria Action Plan
GPI	Gender Parity Index
GVAP	Global Vaccine Action Plan
HIV	Human Immunodeficiency Virus
ICPD	International Conference on Population and Development
ICT	Information and Communications Technology
IDD	Iodine Deficiency Disorders
ILO	International Labour Organization
IPT	Intermittent Preventive Treatment
IPTp	Intermittent Preventive Treatment of Pregnant women
IRS	Indoor Residual Spraying
ITN	Insecticide Treated Net
IUD	Intrauterine Device
JMP	Joint Monitoring Programme
KASF	Kenya AIDS Strategic Framework

KCPE	Kenya Certificate of Primary Education
KCSE	Kenya Certificate of Secondary Education
KDHS	Kenya Demographic and Health Survey
KEBS	Kenya Bureau of Standards
KEPI	Kenya Expanded Programme on Immunization
KHPF	Kenya Health Policy Framework
KNASP	Kenya National AIDS Strategic Plan
KNBS	Kenya National Bureau of Statistics
LAM	Lactational Amenorrhea Method
MDG	Millennium Development Goals
MICS	Multiple Indicator Cluster Survey
MICS5	Fifth global round of Multiple Indicator Clusters Surveys programme
MoH	Ministry of Health
MTP	Medium Term Plans
NAR	Net Attendance Rate
NASSEP V	National Sample Survey and Evaluation Programme V
NHSSP II	National Health Sector Strategic Plan II
NNAP	National Nutrition Action Plan
NTFIC	National Tobacco Free Initiative Committee
ORS	Oral Rehydration Salts
ORT	Oral rehydration treatment
PMI	Presidents Malaria Initiative
PMTC	Prevention of Mother to Child Transmission
PNC	Post-natal Care
PNHC	Post-natal Health Checks
PPM	Parts Per Million
PSRI	Population Studies and Research Institute, University of Nairobi
RHF	Recommended Home Fluid
SP	Sulfadoxine-Pyrimethamine
SPSS	Statistical Package for Social Sciences
STIs	Sexually Transmitted Infections
SUN	Scaling Up Nutrition
TFR	Total Fertility Rate
UNAIDS	United Nations Programme on HIV/AIDS
UNDP	United Nations Development Programme
UNFPA	United Nations Population Fund
UNGASS	United Nations General Assembly Special Session on HIV/AIDS
UNICEF	United Nations Children's Fund
WFFC	World Fit for Children
WHO	World Health Organization

Foreword

The 2013/14 Multiple Indicator Cluster Survey (MICS5) covering Bungoma, Kakamega and Turkana Counties are part of the fifth global round of Multiple Indicator Cluster Survey series conducted worldwide to provide up-to-date information on the situation of children and women. This survey was conducted in collaboration with the Population Studies and Research Institute (PSRI) of the University of Nairobi, the Kenya National Bureau of Statistics (KNBS) and United Nations Children's Fund (UNICEF).

The results of this survey provide requisite baseline information that can be used to facilitate evidence-based planning, budgeting and programming by policymakers and stakeholders at the county levels. The reports will go a long way in encouraging increased demand for use of statistics by policy makers at devolved levels; ensure that resources at both county and national levels are used most effectively through well-planned projects/programmes that will benefit especially the women and children of the three counties.

MICS5 was conducted at county level to provide comprehensive and disaggregated data to partly fill the existing data gaps at this level. This survey is the second of its kind to be conducted at the devolved level after MICS4 conducted in the six counties of the Nyanza region in 2011. MICS3 was conducted in all the 13 districts of the then Eastern Province in 2008.

The MICS5 results are critical in gauging milestones achieved in the field of education, nutrition, child development, and health for women and children in the three counties and in evaluating the various health based policies that the Government has formulated over the years towards achieving the national welfare objectives.

More specifically, the 2013/14 MICS5 data is critical in informing the future planning for the three counties, especially in view of the new constitutional dispensation and Vision 2030. It is anticipated that MICS5 will supplement the data collected during 2014 Kenya Demographic and Health Survey (KDHS). In addition, the information collected will inform strategic communication for social and behaviour change interventions by Government and partners including UNICEF. Furthermore, the data will contribute to the improvement of data and monitoring systems in the three counties.

The survey laid emphasis on quality in every step of the process, right from the design of the tools, training of interviewers, monitoring of data collection, and the whole process of data processing. The MICS5 has much to offer to the health and family planning professionals, government planners, NGOs, researchers, and gender specialists. The potential users are numerous. It is, therefore, our appeal that the findings of MICS5 be put into good use so as to improve the well-being of people in the counties; to prepare reasonable and realistic objectives for county projects; to draw attention to critical problems and inequities; and to determine budgetary priorities.

This report is a culmination of concerted efforts of various organizations and individuals. I have the greatest pleasure to give credit to the technical and financial assistance from UNICEF. I wish to appreciate the organizations, especially Population Studies and Research Institute of the University of Nairobi, that have contributed so much time, energy, and expertise to providing these findings and results. In addition I commend the hard work and dedication of Kenya National Bureau of Statistics (KNBS) staff in assisting to plan and implement this Survey. I thank the interviewers, editors, supervisors, who traversed the three counties, knocking on doors and spending hours talking to household respondents to generate the data. They faced a variety of challenges from occasional vehicle breakdowns, bad terrains, changing weather to basic accommodation. I wish to thank the

respondents who generously and voluntarily provided the information. Without them, there would have been no report to talk about. Much gratitude goes to the data processing specialists and data editors for dedicating their time and expertise to put together quality data. All of them did a tremendous job.

Zachary Mwangi
Director General, Kenya National Bureau of Statistics

Acknowledgements

Kenya implemented the Multiple Indicator Cluster Survey (MICS5) in 2013/2014 in the three counties of Bungoma, Kakamega and Turkana as part of Global MICS round five. MICS is an international household survey programme developed by UNICEF. MICS provides up-to-date information on the situation of children and women and measures key indicators that allow countries to monitor progress towards the Millennium Development Goals (MDGs) and other internationally agreed upon commitments. In Kenya, this information is important to guide the planning and implementation of new development plans targeting the new administrative County -levels of governance.

The successful implementation of the MICS5 was due to the great support and dedication of the partners. Kenya would like to thank the following collaborating organizations:

- United Nations Children’s Fund
- Kenya National Bureau of Statistics

We do appreciate the financial support provided by the United Nations Children’s Fund. Special thanks go to the technical experts from the Kenya National Bureau of Statistics and Population Studies and Research Institute (PSRI) who ensured that the survey was implemented efficiently and effectively to produce quality results. These experts included officers from the collaborating institutions. They exhibited high degree of professionalism during the preparatory work prior and during the implementation stage as well as during the data analysis and report writing. We also thank the UNICEF Regional Office for East and Southern Africa and UNICEF Kenya Country Office for the technical support provided to Kenya during MICS5. We especially recognize and appreciate the support of Dr. Paul Mpuga, Dr. Monica Chizororo, Mr. Nicholas Oloo, Dr. Robert Ndugwa, Dr John Ndegwa Wagai and Dr. Nyasha Madzingira.

Our deepest gratitude goes to the Kenyan Core Technical team responsible for implementing the MICS. The team consisted of technical staff from the PSRI lead by Prof. Lawrence Ikamari supported by Mr. Ben Obonyo, Dr. Wanjiru, Dr. Samuel Wakibi, Dr Andrew Mutuku and Dr. Odipo. The survey could not have been such a success without the guidance and expertise of the Kenya National Bureau of Statistics. In particular, the immeasurable support, advice and guidance of Mr. Zachary Mwangi – Director General, KNBS, Mr Macdonald Obutho – Director Population and Social Statistics, Mr. Robert Buluma, Mr. James Ng’ang’a and Bernard Obasi. This core team effectively implemented the entire MICS5 household survey.

Finally, the most heartfelt gratitude goes to the County Statistical Officers in Bungoma, Kakamega and Turkana; Supervisors, KNBS enumerators, Research Assistants, the Village Elders and all the respondents who participated in the generation of data that made this survey successful.

Prof. Murungaru Kimani
Director
Population Studies and Research Institute
University of Nairobi

Executive Summary

The Turkana County Multiple Indicator Survey (MICS) is a representative sample survey designed to provide estimates for a large number of indicators on the situation of children and women at the county level, for urban and rural areas. The survey used two-stage stratified cluster sampling where the first stage selected 58 clusters from the KNBS fifth National Sample Survey and Evaluation Program (NASSEP V) household-based master sampling frame using equal probability selection method (EPSEM). The second stage randomly selected a uniform sample of 30 households in each cluster from a list of households in the cluster using systematic random sampling method. The survey was implemented by the University of Nairobi through Population studies and Research Institute in collaboration with Kenya National Bureau of Statistics (KNBS) with support from UNICEF Kenya.

Information was collected from a total of 1,277 households representing 93 percent response rate. The composition of these households was 6,594 household members comprising 3,274 males and 3,321 females. The mean household size was 5.2 persons. About 49 percent of the sampled households' population is below 15 years, 48 percent are age 15-64 years and three percent are age 65 years and above.

Due to data quality issues, data relating to mortality and anthropometric measures were not analyzed and reported. Anthropometric data suffered from digit preference for both weight and height, while for mortality, deaths especially among under-5 years old were under reported. KDHS 2014 had similar shortcomings.

Nutrition

Weight at birth is a good indicator not only of a mother's health and nutritional status but also the new-born's chances for survival, growth, long-term health and psychosocial development. The survey findings show that 39 percent of the live-born births in the two years preceding the survey were weighed at birth, and approximately eight percent of infants weighed less than 2,500 grams at birth.

Ninety-seven percent of the children were ever breastfed and 54 percent of babies were breastfed for the first time within one hour of birth. By age 12-15 months, 92 percent of children continued breastfeeding and by age 20-23 months, only 61 percent were still being breastfed. Among children under age 3 years, the median duration of any breastfeeding was 22 months, 5 months for exclusive breastfeeding, and 7 months for predominant breastfeeding. Age appropriate breastfeeding was 54 percent for girls and 51 percent for boys. Only 18 percent of the children age 6-23 months were receiving solid, semi-solid and soft foods the minimum number of times. The overall assessment using the indicator of minimum acceptable diet revealed that only three percent were benefitting from a diet sufficient in both diversity and frequency. About 10 percent of children under-six months were fed using a bottle with a nipple. The findings indicate that bottle feeding becomes prevalent mainly from 6 months of age and older. In 63 percent of households, salt was found to contain at least 15 parts per million (ppm) or more of iodine.

Child Health

Immunization plays a key part in reducing preventable child diseases and mortality. The percentage of children who were fully vaccinated by their first birthday is 64 percent. Overall, 79 percent of children age 12-23 months were fully immunized against vaccine preventable childhood diseases.

Seventeen percent of children under five years of age were reported to have had diarrhoea in the two weeks preceding the survey, five percent had symptoms of ARI, and 19 percent had an episode of fever. Overall, a health facility or provider was seen in 70 percent of cases among children with diarrhoea. Approximately 72 percent of children with diarrhoea received one or more of the recommended home treatments (i.e. were treated with ORS or any recommended homemade fluid), while about half of them received zinc. In addition, 27 percent received ORS and zinc. About 37 percent of households had at least one insecticide treated net and 25 percent of children under the age of five years slept under an ITN the night preceding the survey. Thirty-seven percent of pregnant women slept under any mosquito net the night prior to the survey and 34 percent slept under an insecticide treated net. Overall, 45 percent of children with a fever in the previous two weeks preceding the survey had blood taken from a finger or heel for testing. Eighteen percent of children who had fever in the two weeks preceding the survey were treated with any antimalarial drugs.

Water and Sanitation

Safe drinking water is a basic necessity for good health. Unsafe drinking water can be a significant determinant of diseases such as cholera, typhoid, and schistosomiasis. Drinking water can also be contaminated with chemical and physical contaminants with harmful effects on human health. In addition to preventing disease, improved access to drinking water may be particularly important for women and children, especially in rural areas, who bear the primary responsibility for carrying water, often for long distances.

In Turkana, 72 percent of the population use an improved source of drinking water. Only three percent of household members in households using unimproved drinking water sources are using an appropriate water treatment method. For 22 percent of the household population in the survey, it takes 30 minutes or more to get to the water source and bring water from an improved water source. In the majority of households (80 percent), an adult female usually collects drinking water when the source is not on the premises. Twenty five percent of the household population is using an improved sanitation facility (12 percent not shared and 13 percent public or shared with other households). In 27 percent of the cases, children's stool is disposed of safely. The percentage of households where a place for hand washing was observed is only four percent. Eighty-seven percent of the households had no specific place for hand washing in the dwelling, yard, or plot. The percentage of households with soap or other cleansing agent anywhere in the dwelling was 34 percent.

Reproductive Health

Empowering women and adolescent girls to exercise their sexual and reproductive health rights is a necessary condition for sustainable development. The findings show that age specific fertility rate and birth rate for the three years preceding the survey fertility is 101 births per 1,000 women among adolescents age 15-19 years. Sixteen percent of women age 15-19 years had begun childbearing, three percent were pregnant with their first child, and three percent have had a live birth before age 15. The proportion of women with a live birth before age 15 is seven percent in urban areas and five percent in rural areas.

Contraception by women currently married or in union is 14 percent and one in ten women use injectables. Unmet need for family planning was 22 percent. The percentage of women age 15-49 years who gave birth in the two years preceding the survey and received antenatal care were 94 percent. Ninety-four percent received ANC from a skilled provider and 44 percent of the women had four or more ANC visits. Among those women who had a live birth during the last two years preceding the survey, 70 percent had blood pressure checked, and urine and blood samples taken. Thirty-five percent of births occurring in the two years preceding the MICS were delivered by skilled personnel. Thirty-five percent of births were delivered in a health facility. Overall, 71 percent of women who gave

birth in a health facility stayed 12 hours or more in the facility after delivery. Overall, 36 percent of newborns received a health check following birth while in a health facility or at home. A total of 37 percent of all mothers received a post-natal health check.

Early Childhood Development

In Turkana County, about 31 percent of children age 36-59 months are attending an organised early childhood education programme. Fifty-three percent of children age 36-59 months have an adult household member engaged in four or more activities that promote learning and school readiness. The father's involvement in such activities was low, with only one percent of children age 36-59 months with fathers involved in four or more activities. Mother's engagement in four or more activities that promote learning during the three days preceding the survey is also very low, at three percent. Availability of children's books for those age 0-59 months was virtually non-existent. Nineteen percent of children age 0-59 months had two or more types of playthings to play with in their homes. A total of 54 percent of children were left with inadequate care, either by being left alone or in the care of another child. Child development index is calculated as the percentage of children who are developmentally on target in at least three of the four component domains such as language-cognitive, physical, social-emotional, and approaches to learning. In Turkana County, half of children age 36-59 months are developmentally on track.

Literacy and Education

Youth Literacy Rate as a measure of the effectiveness of the primary education system is often seen as a proxy measure of social progress and economic achievement. Forty-four percent of children who were attending the first grade of primary school at the time of the survey were attending pre-primary school the previous year. About 55 percent of young women age 15-24 years were literate. Among those with primary school as their highest level of education, 71 percent were able to read the statement shown to them. Thirty-three percent of children age 6-13 years were out of school, with a low attendance rate of 47 percent for children age 6, who appeared to be starting late in school. Thirteen percent of the children of secondary school age were out of school. The majority of all children starting grade one were expected to reach grade 8 (96 percent). The gender parity index (GPI) for primary and secondary school was 0.97, suggesting boys and girls of primary school age attended primary education at the same rate.

Child Protection

A name and nationality is every child's right, enshrined in the Convention on the Rights of the Child (CRC) and other international treaties. The findings show that the births of 40 percent of children under five years are registered. Births registration is higher among urban children (54 percent) compared with rural children (26 percent). The combined percentage of children age 5-17 years by involvement in economic activities or household chores, plus percentage working under hazardous conditions during, and percentage engaged in child labour during the week preceding the survey was 23 percent. About 64 percent of children age 1-14 years were subjected to at least one form of psychological aggression or physical punishment by household members during the past month. Among women age 15-49 years, nine percent were married before age 15 and, among women age 20-49 years, 11 percent were married before age 15 while 32 percent were married before age 18. Among currently married/in union women age 20-24 years, about one in five are married/in union with a man who is older by 10 years or more (21 percent).

Three percent of women have some form of female genital mutilation. The practice is five percent prevalent in urban areas and less than 1 percent in rural areas. Six percent of women think FGM should be continued while 87 percent believe it should be discontinued. Overall, 70 percent of women feel that a husband/partner is justified in hitting or beating his wife in at least one of the five situations (if she goes out without telling her husband, neglects children, argues with husband, if the wife refuses to have sex with the husband if she burns the food). Nearly 18 percent of children live with neither of their biological parents and the proportion is higher in urban areas (21 percent) than rural areas (14 percent). One percent of children age 0-17 has one or both parents living abroad.

HIV/AIDS and Sexual Behaviour

Almost all women age 15-49 years (98 percent) have knowledge of AIDS. Thirty-five percent know of the two main ways of preventing HIV transmission with 76 percent knowing having only one faithful uninfected partner and 38 percent know using a condom every time as main ways of preventing HIV transmission.

Overall, 29 percent of women have comprehensive knowledge of HIV prevention methods and transmission which is higher in urban (34 percent) than rural areas (20 percent) and also varies with education and wealth status. In total, 59 percent of women rejected the two most common misconceptions that HIV can be transmitted through mosquito bites and by sharing food with someone with HIV and know that a healthy-looking person (81 percent) can be HIV-positive. About 88 percent and 72 percent of women know that supernatural means and mosquito bites cannot transmit HIV, respectively. Ninety-one percent of women age 15-49 years know that HIV can be transmitted from mother to child by at least one of the three means; during pregnancy, delivery and breastfeeding while 22 percent of women know all three ways of mother-to-child transmission. Ninety-seven percent of women age 15-49 years who have heard of AIDS agree with at least one accepting attitude. The most common accepting attitude is willingness to care for a family member with AIDS in own home (84 percent). More educated women tend to have a more accepting attitude than those with no education. Eighty-eight percent of women age 15-49 years know of a place where to be tested, while 78 percent have been tested. Fifty-seven percent of women know the result of their most recent test. The proportion of women age 15-49 years that had been tested within the last 12 months preceding the survey is 58 percent, while those who had been tested within the last 12 months and know the result is 53 percent. Eighty-four percent of women age 15-49 years with a live birth in the last two years preceding the survey received HIV counselling during ANC, 85 percent were offered an HIV test and were tested for HIV; and 78 percent received HIV counselling, were offered an HIV test, accepted and received the results.

Two percent of women 15-49 years of age reported that they had sex with more than one partner in the last 12 months with a mean number of lifetime sexual partners as 1.3. Thirty-five percent of young women have comprehensive knowledge. Young women who know of three means of HIV transmission from mother-to-child are 20 percent and 86 percent have knowledge of a place to get tested. About 57 percent of young women age 15-24 years, who were sexually active, had been tested for HIV in the last 12 months and know the result. The proportion is low among young women with secondary/higher education (72 percent) compared with those with primary education (75 percent). Overall, eight percent of young women age 15-24 years reported ever having sex before age 15. Further, three percent of young women had sex with more than one partner in the last 12 months.

Only 31 percent of women used a condom the last time they had sex. About 18 percent of women age 15-24 years who had sex in the last 12 months, had sex with a man 10 or more years older.

Access to Mass Media and Use of Information/Communication Technology

About nine percent of women age 15-49 years in Turkana County read a newspaper or magazine, 19 percent listen to the radio, and 16 percent watch television at least once a week. Overall, 72 percent do not have regular exposure to any of the three media, while 27 percent are exposed to at least one, and three percent to all the three types of media on a weekly basis. Women below the age of 25 years are more likely compared to older women to report exposure to all three types of mass media. Women with higher education are more likely to have been exposed to all three types of media (16 percent) than women with primary education (2 percent). Similarly, women from the richest households are more likely to have been exposed to all three types of media (12 percent) compared to women from the poorest households (0 percent).

Overall, 12 percent of young women age 15-24 years ever used the internet, while a similar proportion had used the internet during the 12 months preceding the survey. The proportion of young women who used the internet more frequently, at least once a week during the last month, was lower, at 10 percent. Both computer and internet use during the last 12 months were more widespread among the 20-24 year old women. Use of a computer and the internet is associated with area, education, and household wealth. Only about one percent of women with primary education reported using a computer once a week during the last month, while about a quarter of the women with higher education used a computer. Similarly, higher utilisation of the internet is observed among young women in urban areas (14 percent) compared with 6 percent in rural areas.

Subjective Well-being

Young women are the most satisfied with the way they look (98 percent), their health (96 percent), and their family life (90 percent). The percentage of women age 15-24 years who are very or somewhat satisfied with school is 91 percent. In Turkana County, 90 percent of women age 15-24 years are satisfied with their life. The proportions do not vary significantly by age and marital status. The results indicate that 92 percent of women age 15-24 years are very or somewhat happy. About 69 percent of young women age 15-24 years believe that their lives had improved during the last one year and expect that it would get better after one year, while the corresponding proportion for young women age 20-24 years is 56 percent; and by urban/rural areas, 68 percent and 48 percent, respectively.

Tobacco and Alcohol Use

In Turkana County MICS, ever use of any tobacco product among women is 22 percent, while 21 percent smoke cigarettes, or used smoked or smokeless tobacco products on one or more days during the last one month prior to the survey. Smoking was more prevalent in rural than urban areas, and among women with no education. The results further indicate that majority of the smokers are from the poorest and the second poorest wealth quintiles. Only one woman age 15-49 years in our sample stated to be a current cigarette smoker.

About 10 percent of women age 15-49 years, had at least one drink of alcohol on one or more days during the last one month preceding the survey while four percent have had at least one alcoholic drink before the age of 15 years. The proportion who had an alcoholic drink in the month preceding

the survey ranged between three percent and 18 percent by age. Women age 15-49 years in urban areas are twice (5 percent) as likely to have had at least one alcoholic drink before age 16 than their rural counterparts (2 percent). The results further indicate that drinking of at least one alcoholic drink before age 15 increases with increase in wealth quintiles. About two percent of the women in the poorest wealth quintile have had at least one alcoholic drink before age 15 compared to five percent of those from the richest wealth quintile.

1. Introduction

Turkana County is one of the 47 counties in Kenya. Turkana County is situated in the North-western part of Kenya and is bordered by Uganda to the West, South Sudan and Ethiopia to the North and North-east and Lake Turkana to the East. The county has an estimated population of 855 399.²

1.1 Background

This report is based on the Turkana County Multiple Indicator Cluster Survey (MICS), conducted in 2013/14 by the Population Studies and Research Institute, University of Nairobi, in collaboration with Kenya National Bureau of Statistics, as part of the global MICS programme. The survey provides statistically sound and internationally comparable data essential for developing evidence-based policies and programmes, and for monitoring progress toward national goals and global commitments. Among these global commitments are those emanating from the World Fit for Children Declaration and Plan of Action (2002)³, the goals of the United Nations General Assembly Special Session on HIV/AIDS (2001)⁴, the Education for All Declaration (2000)⁵ and the Millennium Development Goals (MDGs) 2000.⁶

A Commitment to Action: National and International Reporting Responsibilities

The governments that signed the Millennium Declaration and the World Fit for Children Declaration and Plan of Action also committed themselves to monitoring progress towards the goals and objectives they contained:

“We will monitor regularly at the national level and, where appropriate, at the regional level and assess progress towards the goals and targets of the present Plan of Action at the national, regional and global levels. Accordingly, we will strengthen our national statistical capacity to collect, analyse and disaggregate data, including by sex, age and other relevant factors that may lead to disparities, and support a wide range of child-focused research. We will enhance international cooperation to support statistical capacity-building efforts and build community capacity for monitoring, assessment, and planning.” (**A World Fit for Children**, paragraph 60)

“...We will conduct periodic reviews at the national and sub-national levels of progress in order to address obstacles more effectively and accelerate actions....” (**A World Fit for Children**, paragraph 61)

The Plan of Action of the World Fit for Children (paragraph 61) also calls for the specific involvement of UNICEF in the preparation of periodic progress reports:

“... As the world’s lead agency for children, the United Nations Children’s Fund is requested to continue to prepare and disseminate, in close collaboration with Governments, relevant funds, programmes and the specialized agencies of the United Nations system, and all other relevant actors, as appropriate, information on the progress made in the implementation of the Declaration and the Plan of Action.”

²Kenya National Bureau of Statistics, 2013. Statistical Abstract 2013.

³A World Fit for Children. Resolution adopted by the United Nations General Assembly 10 May 2002.

⁴United Nations General Assembly Special Session on HIV/AIDS 2001. Summary of the Declaration of Commitment on HIV/AIDS 25-27 June 2001, New York

⁵<http://www.unesco.org/new/en/education/themes/leading-the-international-agenda/education-for-all/>

⁶http://www.who.int/topics/millennium_development_goals/en/

Similarly, the **Millennium Declaration** (paragraph 31) calls for periodic reporting on progress:

“...We request the General Assembly to review on a regular basis the progress made in implementing the provisions of this Declaration, and ask the Secretary-General to issue periodic reports for consideration by the General Assembly and as a basis for further action.”

Kenya’s GDP has grown by an annual average of four percent in the past five years. In 2013, Kenya adopted its second five-year Medium Term Plan (MTP II 2013-17) to implement its ‘Vision 2030’, which represents a solid strategic framework to transform Kenya into a newly industrializing, middle-income country by 2030.⁷ The overarching objective of the Bank’s Country Strategy Paper (CSP) 2014-18 is job creation to realize country’s developmental goals.⁶ The African Development Bank’s Country Strategy Paper (CSP) 2014-18 for Kenya supports the country’s ambitions and addresses its main developmental challenges by promoting job creation as the overarching objective.

The Turkana County MICS results are expected to form part of the baseline data for the post-2015 era. The survey findings are also expected to contribute to the evidence base of several other important initiatives, including Committing to Child Survival: **A Promise Renewed**⁸, a global movement to end child deaths from preventable causes, and the accountability framework proposed by the **Commission on Information and Accountability for the Global Strategy for Women’s and Children’s Health**.⁹

This final report presents the results of the indicators and topics covered in the survey. There are 14 chapters presented as follows:

- Chapter 1: An introductory note to the Turkana County MICS Report;
- Chapter 2: Sample and survey methodology
- Chapter 3: Sample coverage and characteristics of households and respondents
- Chapter 4: Child nutrition
- Chapter 5: Child health
- Chapter 6: Water and sanitation
- Chapter 7: Reproductive health
- Chapter 8: Early childhood development
- Chapter 9: Literacy and education
- Chapter 10: Child protection
- Chapter 11: HIV, AIDS and sexual behaviour
- Chapter 12: Mass Media, Information, and Communication Technology (ICT)
- Chapter 13: Subjective well-being
- Chapter 14: Tobacco and alcohol use

⁷African Development Bank Group, 2014, Kenya Country Strategy Paper 2014-2018.

⁸United Nations Children’s Fund (UNICEF), September 2014. Committing to Child Survival: A Promise Renewed - Progress Report 2014.

⁹WHO. 2014. Implementing the Commission on Information and Accountability Recommendations 2014: Progress Report Accountability for Women’s and Children’s Health.

1.2 Survey Objectives

The 2013/14 Turkana County MICS has as its primary objectives to:

- Provide up-to-date information for assessing the situation of children and women in Turkana County;
- Generate data for the critical assessment of the progress made in various areas, and to put additional efforts in those areas that require more attention;
- Furnish data needed for monitoring progress toward goals established in the Millennium Declaration, and other internationally agreed upon goals, as a basis for future action;
- Collect disaggregated data for the identification of disparities, to allow for evidence based policy-making aimed at social inclusion of the most vulnerable;
- Contribute to the generation of baseline data for the post-2015 agenda;
- Validate data from other sources and the results of focused interventions; and
- Contribute to the improvement of data and monitoring systems in Kenya and to strengthen technical expertise in the design, implementation, and analysis of such systems.

2. Sample and Survey Methodology

Chapter Two presents the survey sample design and methodology, content for the three questionnaires used in the survey, the interviewer training process, fieldwork, and data management and processing.

2.1 Sample Design

The sample for the Turkana County MICS, 2013/14 was designed to provide estimates for a large number of indicators on the situation of children and women at the county level. The urban and rural areas within the county were the main sampling strata. The sample was selected in two stages: cluster and household. The survey utilized the fifth National Sample Survey and Evaluation Program (NASSEP V) household-based master sampling frame which is created and maintained by the Kenya National Bureau of Statistics (KNBS). The primary sampling unit for the frame is a cluster, which constitutes one or more EAs, with an average of 100 households.

For the NASSEP V master sample the EAs were selected within each stratum using systematic sampling with probabilities proportion to size (PPS). For the MICS, within each stratum a specified number of census enumeration areas was selected from the master sample using an equal probability selection method (EPSEM). After a household listing was carried out in the selected clusters, a systematic sample of 30 households was drawn in each sampled cluster. In total, there were 58 clusters which were selected for the survey in Turkana County. The sample was stratified by urban and rural areas, and was not self-weighting. All selected clusters were visited during fieldwork. For reporting county level results, sample weights are used. A more detailed description of the sample design is provided in Appendix C.

2.2 Questionnaires

A set of three questionnaires was used in the survey: 1) a household questionnaire which was administered to the household head or any other responsible member of the household; 2) a questionnaire for individual women administered in each household to all women age 15-49 years; 3) an under-5 questionnaire, administered to mothers (or caretakers) for all children under 5 years living in the household.

The questionnaires included the following modules:

The Household Questionnaire included the following modules:

- List of Household Members
- Education
- Child Labour
- Child Discipline
- Household Characteristics
- Insecticide Treated Nets
- Indoor Residual Spraying
- Water and Sanitation
- Handwashing

- Salt Iodization

The Questionnaire for Individual Women age 15-49 years included the following modules:

- Woman's Background
- Access to Mass Media and Use of Information/Communication Technology
- Fertility/Birth History
- Desire for Last Birth
- Maternal and Newborn Health
- Post-natal Health Checks
- Illness Symptoms
- Contraception
- Unmet Need
- Female Genital Mutilation/Cutting
- Attitudes Toward Domestic Violence
- Marriage/Union
- Sexual Behaviour
- HIV/AIDS
- Tobacco and Alcohol Use
- Life Satisfaction

The Questionnaire for Children Under5 was administered to mothers (or caretakers) of children under 5 years of age¹⁰ living in the households. Normally, the questionnaire was administered to mothers of under-5 children; in cases when the mother was not listed in the household roster, a primary caretaker for the child was identified and interviewed. The questionnaire included the following modules:

- Age
- Birth Registration
- Early Childhood Development
- Immunization
- Breastfeeding and Dietary Intake
- Care of Illness
- Anthropometry

Due to data quality issues, data relating to mortality and anthropometric measures were not analyzed and reported. Anthropometric data suffered digit preference for both weight and height, while for mortality, deaths especially among children under-five years were under reported. The recommendation to remove the Mortality Chapter and the anthropometric measures section from the Nutrition Chapter was adopted at the final reports validation workshop organized by KNBS, PSRI and UNICEF. KDHS 2014 had similar shortcomings. The DQ tables are included in the report for reference. The MICS data set can be accessed and evaluated by researchers for further analysis. The survey team, KNBS and the Population Studies and Research Institute will review the data in detail to identify challenges encountered and to address them before the next round of surveys.

The questionnaires are based on the MICS5 model questionnaire.¹¹ From the MICS5 model English version, the questionnaires were customised and translated into Kiswahili and Turkana languages and

¹⁰ The terms "children under 5", "children age 0-4 years", and "children age 0-59 months" are used interchangeably in this report.

¹¹ The model MICS5 questionnaires can be found at http://www.childinfo.org/mics5_questionnaire.html

were pre-tested in four clusters (rural and urban) in Trans Nzoia County. Based on the results of the pre-test, modifications were made to the wording and translation of the questionnaires. A copy of the Turkana County MICS questionnaires is provided in Appendix H.

In addition to administering of questionnaires, fieldwork teams tested the salt used for cooking in the households for iodine, observed the place for handwashing, and measured the weights and heights of children under-5 years of age. Details and findings of these observations and measurements are provided in the respective sections of the report.

2.3 Training and Fieldwork

Training for the fieldwork was conducted in Kitale town for 14 days from 24th October to 6th November, 2013. Training included lectures on interviewing techniques and the content of the questionnaires, and mock interviews between trainees to gain practice in asking questions. Facilitators used a variety of methods which included PowerPoint presentations, illustrations on flip charts, question and answer, case studies, group work and group discussions. Towards the end of the training period, trainees spent two days practising the research tools by interviewing respondents in selected urban and rural clusters in Trans Nzoia County.

Fieldwork began in November 2013 and concluded in February 2014. The survey team was divided into two groups. Each group comprised of five interviewers, one driver, one editor, one measurer and a supervisor.

2.4 Data Processing

CSPRO software, Version 5.0 running on desktop computers was used for data entry. Data entry was done by a trained team of 14 data entry operators, one Archivist/System administrator and one data entry supervisor. For quality assurance purposes, all questionnaires were double-entered and internal consistency checks performed. Procedures and standard programs developed under the global MICS programme and adapted to the Turkana County MICS questionnaire were used throughout. Data processing began simultaneously with data collection in November 2013 and was completed in February 2014. Data were analysed using the Statistical Package for Social Sciences (SPSS) software, Version 21. Model syntax and tabulation plans developed by UNICEF were customized and used for this purpose.

3. Sample Coverage and the Characteristics of Households and Respondents

This chapter discusses sample coverage, characteristics of households and female respondents age 15-49 years and children under-five years of age. The chapter also presents housing characteristics, asset ownership and household wealth quintiles.

3.1 Sample Coverage

Table HH1 shows the results of household, women's and under-5 interviews for Turkana County. A total of 1,680 households were selected for the sample in Turkana County out of which 1,379 were occupied. Of these, 1,277 were successfully interviewed giving a household response rate of 93 percent. In the interviewed households, 1,300 women age 15-49 years were identified as eligible. Of these, 1,104 were successfully interviewed, yielding a response rate of 85 percent.

There were 1,117 eligible children under age five who were listed in the interviewed households. The questionnaires were completed for 1,067 of these children by their mothers or caretakers giving a response rate of 96 percent for eligible children.

The overall response rate was 80 percent for women in urban areas and 78 percent for those in rural areas. For children under-5 years, the overall response rate was 89 percent in rural areas and 88 percent in urban areas. The response rate for women in urban and rural areas was generally low and this may be due to failure by interviewers to reach the respondents (lack of access). Table DQ.2 in Appendix F shows that women age 15-19 years were those who had the lowest response rate (76 percent).

Table HH.1: Results of household, women's, men's and under-5 interviews

Number of households, women, men, and children under 5 by interview results, and household, women's and under-5's response rates, Turkana County MICS, 2013/14

	Total	Area	
		Urban	Rural
Households			
Sampled	1,680	660	1,020
Occupied	1,379	575	804
Interviewed	1,277	533	744
Household response rate	92.6	92.7	92.5
Women			
Eligible	1,300	645	655
Interviewed	1,104	554	550
Women's response rate	84.9	85.9	84.0
Women's overall response rate	78.6	79.6	77.7
Children under 5			
Eligible	1,117	449	668
Mothers/caretakers interviewed	1,067	424	643
Under-5's response rate	95.5	94.4	96.3
Under-5's overall response rate	88.5	87.5	89.1

3.2 Characteristics of Households

Table HH2 provides information on the age distribution of household population by sex. The population distribution is also used to produce the population pyramid in Figure HH.1. A total of 1,277 households successfully interviewed in Turkana with 6,594 household members listed. Of these, 3,274 (50 percent) are males, and 3,321 are females. The county population distribution is similar to the national population distribution by sex as per the 2009 Housing and Population Census.¹²

Table HH.2: Age distribution of household population by sex

Percent and frequency distribution of the household population by five-year age groups, dependency age groups, and by child (age 0-17 years) and adult populations (age 18 or more), by sex, Turkana County MICS, 2013/14

	Total		Males		Females	
	Number	Percent	Number	Percent	Number	Percent
Total	6,594	100.0	3,274	100.0	3,321	100.0
Age						
0-4	1,095	16.6	551	16.8	544	16.4
5-9	1,096	16.6	561	17.1	534	16.1
10-14	1,018	15.4	508	15.5	511	15.4
15-19	765	11.6	413	12.6	351	10.6
20-24	527	8.0	266	8.1	262	7.9
25-29	477	7.2	228	7.0	249	7.5
30-34	302	4.6	140	4.3	161	4.9
35-39	310	4.7	153	4.7	157	4.7
40-44	209	3.2	109	3.3	100	3.0
45-49	161	2.4	76	2.3	85	2.6
50-54	194	2.9	85	2.6	109	3.3
55-59	138	2.1	41	1.2	97	2.9
60-64	108	1.6	55	1.7	53	1.6
65-69	96	1.5	41	1.3	55	1.6
70-74	57	0.9	29	0.9	28	0.9
75-79	21	0.3	4	0.1	17	0.5
80-84	14	0.2	8	0.3	6	0.2
85+	4	0.1	3	0.1	2	0.0
Missing/DK	2	0.0	2	0.1	0	0.0
Dependency age groups						
0-14	3,209	48.7	1,620	49.5	1,589	47.9
15-64	3,190	48.4	1,567	47.9	1,624	48.9
65+	193	2.9	85	2.6	108	3.2
Missing/DK	2	0.0	2	0.1	0	0.0
Child and adult populations						
Children age 0-17 years	3,699	56.1	1,876	57.3	1,823	54.9
Adults age 18+ years	2,894	43.9	1,396	42.6	1,498	45.1
Missing/DK	2	0.0	2	0.1	0	0.0


The population pyramid (Figure HH.1) is broad based. However, the pattern exhibited is slightly different from the national population pyramid from the 2009 Housing and Population Census. The national population pyramid from the 2009 census was smooth and showed a higher percentage of

¹² Kenya National Bureau of Statistics, 2010. The 2009 Kenya Population and Housing Census

the population in the 0-4 year age group than in the 5-9 year age group, with the proportion for females age 0-4 slightly higher. On the contrary, the population pyramid from the MICS5 shows more or less a similar percentage of the population in the 0-4 year age group and in the 5-9 year age group. This may be due to interviewers’ bias (out transference) in order to reduce the number of under-five questionnaires to administer. There is also a noticeable drop in the age group 20-24 years especially for males, which may be an indication of out-migration of the population from the county to other areas either for further education or for employment opportunities or other reasons.

In Turkana County, a high proportion of the population (49 percent) is below 15 years of age (Table HH.2). Forty-eight percent of the population comprises of people in the age group 15 to 64 years while three percent are age 65 years and above. This implies that the dependent population in Turkana County (i.e. persons below age 15 years and those 65 and above) account for 52 percent. Fifty-six percent of the population is under the age of 18 years. The percentage of males under the age of 18 years is 57 percent, with females at 55 percent.

Figure HH.1: Age and sex distribution of household population, Turkana County MICS, 2013/14



Note: 2 household members with missing age and/or sex are excluded

Tables HH.3, HH.4 and HH.5 provide basic information on the households, female respondents age 15-49 years, and children under-5 years. Both unweighted and weighted numbers are presented. Such information is essential for the interpretation of findings presented later in this report and provides background information on the representativeness of the survey sample. The remaining tables in this report are presented only with weighted numbers.¹³

¹³ See Appendix C: Sample Design, for more details on sample weights.

Table HH.3 provides basic background information on the household composition which include the sex of the household head, area, number of household members, education of household head, and ethnicity of the household head. These background characteristics are used in subsequent tables in this report. The figures in the table are also intended to show the numbers of observations by major categories of analysis in the report.

Table HH.3: Household composition			
Percent and frequency distribution of households by selected characteristics, Turkana County MICS, 2013/14			
	Weighted percent	Number of households	
		Weighted	Unweighted
Total	100.0	1,277	1,277
Sex of household head			
Male	51.3	656	642
Female	48.7	621	635
Area			
Urban	53.5	684	533
Rural	46.5	593	744
Number of household members			
1	7.1	91	85
2	9.8	125	122
3	10.6	135	132
4	16.1	205	207
5	15.1	193	200
6	13.4	172	181
7	10.9	139	134
8	6.1	78	83
9	4.3	54	56
10+	6.7	85	77
Education of household head			
None	64.3	822	912
Primary	16.5	211	177
Secondary+	18.3	234	180
Missing/DK	0.9	11	8
Ethnicity of household head			
Turkana	81.9	1046	1092
Other ethnic group	17.7	226	180
Missing/DK	0.4	5	5
Mean household size	5.2	1,277	1,277

The weighted and unweighted total number of households are equal, since sample weights were normalized.¹³ The table shows the weighted mean household size of 5.2 persons estimated by the survey. The proportion of households headed by males and females in Turkana County is almost the same. Fifty-one percent of the households are headed by males while 49 percent are headed by females. Fifty-four percent of the households are in urban areas while 46 percent are in rural areas. About a third (31 percent) of the households have household sizes of 4-5 persons, 20 percent have 2-3 persons, 24 percent had 6-7 persons, seven percent have one person, 10 percent had 8-9 persons and seven percent have 10 or more persons. Most heads of households have no education (64 percent), while 17 percent had primary education and 18 percent have secondary/higher education.

Most of the heads of households (82 percent) are headed by persons of the Turkana ethnic group while 18 percent are headed by persons of other ethnic groups.

3.3 Characteristics of Female Respondents 15-49 Years of Age and Children Under-5 Years

Tables HH.4 and HH.5 provide information on the background characteristics of female respondents age 15-49 years and of children under-5 years. In all three tables, the total number of weighted and unweighted observations are equal, since sample weights have been normalized (standardized).¹³ In addition to providing useful information on the background characteristics of women and children under age five, the tables are also intended to show the numbers of observations in each background category. These categories are used in the subsequent tabulations of this report.

Table HH.4 provides background characteristics of female respondents, age 15-49 years. The table includes information on the distribution of women according to area, age, marital/union status, motherhood status, births in last two years, education¹⁴, wealth index quintiles^{15, 16}, and ethnicity of the household head. Sixty-two percent of the women interviewed reside in urban areas while 38 percent are in rural areas. Disaggregation of the data by age of the woman shows that 23 percent of the women are age 15-19 years, 19 percent are 20-24 years and 25-29 years age category.

The data shows that 56 percent of the women interviewed are currently married/in union, while 29 percent have never married. About nine percent of the women are either divorced or separated while

¹⁴ Throughout this report, unless otherwise stated, “education” refers to highest educational level ever attended by the respondent when it is used as a background variable.

¹⁵ The wealth index is a composite indicator of wealth. To construct the wealth index, principal components analysis is performed by using information on the ownership of consumer goods, dwelling characteristics, water and sanitation, and other characteristics that are related to the household’s wealth, to generate weights (factor scores) for each of the items used. First, initial factor scores are calculated for the total sample. Then, separate factor scores are calculated for households in urban and rural areas. Finally, the urban and rural factor scores are regressed on the initial factor scores to obtain the combined, final factor scores for the total sample. This is carried out to minimize the urban bias in the wealth index values.

Each household in the total sample is then assigned a wealth score based on the assets owned by that household and on the final factor scores obtained as described above. The survey household population is then ranked according to the wealth score of the household they are living in, and is finally divided into 5 equal parts (quintiles) from lowest (poorest) to highest (richest).

In Turkana County MICS, the following assets were used in these calculations: radio, television, non-mobile telephone, refrigerator, agricultural land, farm animals/livestock, watch, mobile telephone, bicycle, motorcycle or scooter, animal-drawn cart, car or truck, boat with a motor, and ownership of dwelling.

The wealth index is assumed to capture the underlying long-term wealth through information on the household assets, and is intended to produce a ranking of households by wealth, from poorest to richest. The wealth index does not provide information on absolute poverty, current income or expenditure levels. The wealth scores calculated are applicable for only the particular data set they are based on.

Further information on the construction of the wealth index can be found in Filmer, D and Pritchett, L. 2001. *Estimating wealth effects without expenditure data – or tears: An application to educational enrolments in states of India*. Demography 38(1): 115-132; Rutstein, SO and Johnson, K. 2004. *The DHS Wealth Index*. DHS Comparative Reports No. 6; and Rutstein, SO. 2008. *The DHS Wealth Index: Approaches for Rural and Urban Areas*. DHS Working Papers No. 60.

¹⁶ When describing survey results by wealth quintiles, appropriate terminology is used when referring to individual household members, such as for instance “women in the richest population quintile”, which is used interchangeably with “women in the wealthiest survey population”, “women living in households in the richest population wealth quintile”, and similar.

seven percent are widowed.

A high proportion (72 percent) of women age 15-49 years had ever given birth and 35 percent had given birth in the last two years preceding the survey while 37 had never given birth in the last two years. The majority of women (56 percent) have no formal education while 27 percent have primary level of education and 17 percent have secondary or higher level of education.

Table HH.4: Women's background characteristics			
Percent and frequency distribution of women age 15-49 years by selected background characteristics, Turkana County MICS, 2013/14			
	Weighted percent	Number of women	
		Weighted	Unweighted
Total	100.0	1,104	1,104
Area			
Urban	61.9	683	554
Rural	38.1	421	550
Age			
15-19	22.8	252	228
20-24	18.9	209	197
25-29	19.0	210	216
30-34	13.0	144	151
35-39	12.1	134	141
40-44	7.4	82	92
45-49	6.8	75	79
Marital/Union status			
Currently married/in union	55.7	615	649
Widowed	6.5	72	76
Divorced	2.6	29	25
Separated	6.4	71	64
Never married/in union	28.5	314	288
Missing	0.3	3	2
Motherhood and recent births			
Never gave birth	27.6	304	279
Ever gave birth	72.4	800	825
Gave birth in last two years	35.1	387	412
No birth in last two years	37.4	412	413
Education			
None	56.3	622	702
Primary	26.8	296	254
Secondary+	16.9	186	148
Wealth index quintile			
Poorest	15.3	169	232
Second	17.3	191	232
Middle	20.4	226	214
Fourth	22.5	249	215
Richest	24.4	270	211
Ethnicity of household head			
Turkana	80.0	883	916
Other ethnic group	19.8	218	184
Missing/DK	0.2	3	4

In households where there were children under the age of five years, the mothers/caretakers were interviewed. Background characteristics of children under-5 years are presented in Table HH.5. These include the distribution of children by several attributes: sex, area, age in months, respondent type, mother's (or caretaker's) education, wealth, and ethnicity.

The results show that there is no difference between the number of male and female children under-5 years covered in the survey. Fifty-one percent of children under-5 years are in urban areas, while 49 percent are in rural areas. About 21 percent of the children surveyed are age 0-11 months while 18 percent are age 12-23 years. Ninety-three percent of the women who responded to the questions about the child under-5 years are mothers of the children and seven percent are caretakers. Seventy-one percent of caregivers of children under-5 years of age have no education while 19 percent have. About 43 percent of the children under-5 years are in the poorest and second poorest wealth quintiles.

Table HH.5: Under-5's background characteristics

Percent and frequency distribution of children under five years of age by selected characteristics, Turkana County MICS, 2013/14

	Weighted percent	Number of under-5 children	
		Weighted	Unweighted
Total	100.0	1,067	1,067
Sex			
Male	50.3	537	536
Female	49.7	530	531
Area			
Urban	51.2	546	424
Rural	48.8	521	643
Age			
0-5 months	10.0	107	111
6-11 months	11.2	120	120
12-23 months	18.3	196	191
24-35 months	19.2	205	204
36-47 months	20.8	222	221
48-59 months	20.4	217	220
Respondent to the under-5 questionnaire			
Mother	93.3	995	987
Other primary caretaker	6.7	72	80
Mother's education^a			
None	71.1	758	818
Primary	19.4	207	173
Secondary+	9.3	99	74
Missing/DK	0.3	3	2
Wealth index quintile			
Poorest	20.3	216	271
Second	22.9	244	283
Middle	20.3	217	207
Fourth	21.3	227	184
Richest	15.2	163	122
Ethnicity of household head			
Turkana	84.2	898	932
Other ethnic group	15.7	167	132
Missing/DK	0.1	1	3

^a In this table and throughout the report, mother's education refers to educational attainment of mothers as well as caretakers of children under 5, who are the respondents to the under-5 questionnaire if the mother is deceased or is living elsewhere.

3.4 Housing characteristics, asset ownership, and wealth quintiles

Tables HH.6, HH.7 and HH.8 provide results on household characteristics and assets in connection to household wealth. Table HH.6 presents characteristics of housing, disaggregated by area and region, distributed by whether the dwelling is connected to electricity, the main materials of the floor, roof, and exterior walls, as well as the number of rooms used for sleeping.

The data indicates that only a few households in Turkana County have electricity connections. Only 11 percent of households have electricity (19 percent in urban areas and one percent in rural areas). About 83 percent of the households have natural floors¹⁷ (71 percent in urban and 96 percent in rural areas), while 16 percent have a finished floor¹⁸ (29 percent in urban areas and two percent in rural areas). Fifty-one percent of households have natural roofing¹⁹ (32 percent in urban areas and 74 percent in rural areas) and 43 percent have finished roofing²⁰ (63 percent in urban areas and 20 percent in rural areas). The results further show that 53 percent of the households have natural walls²¹, 24 percent have rudimentary walls²², and 20 percent have finished walls²³. Data was also collected on the number of sleeping rooms and number of persons sleeping in one room. The majority of the households (68 percent) have one room for sleeping followed by 22 percent which have two rooms. Only eight percent of the households have 3 or more rooms for sleeping. The mean number of persons per room used for sleeping is four.

¹⁷ Natural flooring – earth/sand or dung

¹⁸ Finished floor - Parquet or polished wood, vinyl or asphalt strips, ceramic tiles, cement or carpet

¹⁹ Natural roofing - No Roof, thatch/palm leaf ., or sod

²⁰ Metal/Tin, wood, calamine/cement fibre, ceramic tiles, cement, or roofing shingles

²¹ Natural walls - No walls, cane /palm / trunks or dirt.

²² Rudimentary walls - Bamboo with mud, stone with mud, uncovered adobe, plywood, cardboard, or reused wood

²³ Finished walls – Cement, stone with lime / cement, bricks, cement blocks, covered adobe or wood planks / shingles.

Additional definitions for housing characteristics (Table HH.6) are in Appendix G

Table HH.6: Housing characteristics

Percent distribution of households by selected housing characteristics, according to area of residence and regions, Turkana County MICS, 2013/14

	Area		
	Total	Urban	Rural
Electricity			
Yes	10.8	19.1	1.3
No	89.2	80.9	98.7
Flooring			
Natural floor	83.0	71.4	96.3
Rudimentary floor	0.6	0.0	1.4
Finished floor	16.3	28.6	2.1
Other	0.1	0.0	0.2
Roof			
Natural roofing	51.4	31.5	74.3
Rudimentary roofing	2.2	2.3	2.2
Finished roofing	43.2	63.4	19.9
Other	2.8	2.3	3.5
Missing/DK	0.3	0.5	0.1
Exterior walls			
Natural walls	53.1	32.7	76.7
Rudimentary walls	23.9	34.4	11.8
Finished walls	20.1	29.9	8.8
Other	2.5	2.4	2.7
Missing/DK	0.4	0.7	0.1
Rooms used for sleeping			
1	67.6	64.2	71.5
2	22.0	22.9	21.0
3 or more	7.7	9.5	5.5
Missing/DK	2.7	3.4	1.9
Total	100.0	100.0	100.0
Number of households	1,277	684	593
Mean number of persons per room used for sleeping	4.04	3.97	4.13

In Table HH.7, households are distributed according to ownership of assets by households and by individual household members. This also includes ownership of dwelling unit. The results indicate that 17 percent of the households own a radio (29 in urban areas and four in rural areas) while 10 percent own a television set (18 percent in urban areas and one percent in rural areas). Thirty-six of households own farm animals/livestock (19 percent in urban areas and 56 percent in rural areas) while 13 percent own agricultural land (6 percent in urban areas and 22 percent in rural areas).

The table further shows that 45 percent of households have at least one member who owns a mobile phone (66 percent in urban areas and 22 percent in rural areas), 19 percent have at least one member who has a bank account, seven percent have at least one member who owns a bicycle, and six percent have at least one member who owns a watch. Eighty-one percent of the dwelling units are owned by a household member. Ownership of dwelling unit is higher in rural areas (98 percent) than urban areas

(67 percent). About 16 percent of the dwelling unit are rented (30 percent in urban areas compared to only one percent in rural areas).

Table HH.7: Household and personal assets			
Percentage of households by ownership of selected household and personal assets, and percent distribution by ownership of dwelling, according to area of residence and regions, Turkana County MICS, 2013/14			
	Total	Area	
		Urban	Rural
Percentage of households that own a			
Radio	17.4	28.7	4.4
Television	10.4	18.3	1.3
Non-mobile telephone	2.3	4.1	0.2
Refrigerator	2.7	4.9	0.2
Solar Panel	2.0	2.0	2.0
Chair	1.7	1.5	1.9
Sofa Set	1.9	1.7	2.0
Table	1.7	1.5	1.9
Cupboard	1.9	1.8	2.0
Bed	1.7	1.4	1.9
Clock	2.0	1.9	2.0
Camera	2.0	2.0	2.0
Computer	2.0	2.0	2.0
Percentage of households that own			
Agricultural land	13.3	5.5	22.3
Farm animals/Livestock	36.2	19.2	55.8
Percentage of households where at least one member owns or has a			
Watch	5.9	9.2	2.1
Mobile telephone	45.4	66.0	21.7
Bicycle	7.4	10.1	4.1
Motorcycle or scooter	2.8	4.4	1.0
Animal-drawn cart	0.4	0.8	0.0
Car or truck	1.9	3.6	0.0
Boat with a motor	0.0	0.0	0.1
Bank account	18.9	33.1	2.5
Ownership of dwelling			
Owned by a household member	80.9	66.6	97.5
Not owned	18.9	33.3	2.4
Rented	16.2	29.5	0.9
Other	2.7	3.7	1.4
Missing/DK	0.2	0.2	0.2
Total	100.0	100.0	100.0
Number of households	1,277	684	593

Table HH.8 shows how the household populations in urban and rural areas are distributed according to household wealth quintiles. Most of the households (60 percent) in Turkana County are in the poorest to middle wealth quintiles. The data further shows that whereas only 12 percent of the population residing in urban areas are in the poorest and second poorest wealth quintiles, a large

proportion (74 percent) of the rural populations are in the poorest and second poorest wealth quintiles.

Table HH.8: Wealth quintiles

Percent distribution of the household population by wealth index quintile, according to area of residence and regions, Turkana County MICS, 2013/14

	Wealth index quintile					Total	Number of household members
	Poorest	Second	Middle	Fourth	Richest		
Total	20.0	20.0	20.0	19.9	20.0	100.0	6,594
Area							
Urban	4.4	7.5	23.1	29.6	35.3	100.0	3,598
Rural	38.7	35.0	16.4	8.3	1.6	100.0	2,996

4. Nutrition

About half of Kenya's estimated 38.5 million people are poor, and some 7.5 million people live in extreme poverty, while over 10 million people suffer from chronic food insecurity and poor nutrition. Children are undernourished and micronutrient deficiencies are widespread.^{24, 25}

The Government of Kenya is strongly committed to reducing hunger and malnutrition. Policies and strategies were developed to guide the nutrition interventions and activities in the country. These include the Food and Nutrition Security Policy (FNSP) 2011, National Nutrition Action Plan (NNAP) 2012-2017 and Kenya Health Strategic Plan 2008-2012. Most of these interventions were part of Scaling Up Nutrition (SUN) actions that were implemented globally to accelerate efforts towards achieving MDG 4 and 5. The NNAP is aligned to the government's Medium Term Plans (MTPs) to enable mainstreaming of the nutrition budgeting process into national development plans, and facilitate allocation of resources to nutrition programmes.

This chapter presents the results on birth weight; breastfeeding, and infant and young child feeding practices; and use of iodized salt at household.²⁶

4.1 Birth Weight

Weight at birth is a good indicator not only of a mother's health and nutritional status but also the newborn's chances for survival, growth, long-term health and psychosocial development. Low birth weight (defined as less than 2,500 grams) carries a range of grave health risks for children. Babies who were undernourished in the womb face a greatly increased risk of dying during their early days, months and years. Those who survive may have impaired immune function and increased risk of disease; they are likely to remain undernourished, with reduced muscle strength, throughout their lives, and suffer a higher incidence of diabetes and heart disease in later life. Children born with low birth weight also risk a lower IQ and cognitive disabilities, affecting their performance in school and their job opportunities as adults.

In the developing world, low birth weight stems primarily from the mother's poor health and nutrition. Three factors have most impact: the mother's poor nutritional status before conception, short stature (due mostly to under nutrition and infections during her childhood), and poor nutrition during pregnancy. Inadequate weight gain during pregnancy is particularly important since it accounts for a large proportion of foetal growth retardation. Moreover, diseases such as diarrhoea and malaria, which are common in many developing countries, can significantly impair foetal growth if the mother becomes infected while pregnant.

In the industrialized world, cigarette smoking during pregnancy is the leading cause of low birth weight. In developed and developing countries alike, teenagers who give birth when their own bodies have yet to finish growing run a higher risk of bearing low birth weight babies.

²⁴ Government of Kenya, 2011. National Food and Nutrition Security Policy.

²⁵ The Partnership for Maternal, Newborn and Child Health, 2012. Maternal and Child Health: Kenya

²⁶ A section on anthropometric indicators was excluded from the report due to data quality issues.

One of the major challenges in measuring the incidence of low birth weight is that more than half of infants in the developing world are not weighed at birth. In the past, most estimates of low birth weight for developing countries were based on data compiled from health facilities. However, these estimates are biased for most developing countries because the majority of newborns are not delivered in health facilities, and those who are, represent only a sample of all births.

Since many infants are not weighed at birth and those who are weighed may be a biased sample of all births, the reported birth weights usually cannot be used to estimate the prevalence of low birth weight among all children. Therefore, the percentage of births weighing below 2,500 grams is estimated from two items in the questionnaire: the mother's assessment of the child's size at birth (i.e., very small, smaller than average, average, larger than average, very large) and the mother's recall of the child's weight or the weight as recorded on a health card if the child was weighed at birth.²⁷

In Turkana County, 39 percent of the last live-born births in the last two years preceding the survey were weighed at birth and approximately eight percent of infants weighed less than 2,500 grams at birth (Table NU.1). The prevalence of low birth weight according to background characteristics shows no meaningful variations at this sample size.

²⁷ For a detailed description of the methodology, see Boerma, JT et al. 1996. *Data on Birth Weight in Developing Countries: Can Surveys Help?* Bulletin of the World Health Organization 74(2): 209-16.

Table NU.1: Low birth weight infants

Percentage of last live-born children in the last two years that are estimated to have weighed below 2,500 grams at birth and percentage of live births weighed at birth, Turkana County MICS, 2013/14

	Percent distribution of births by mother's assessment of size at birth					Total	Percentage of live births:		Number of last live-born children in the last two years
	Very small	Smaller than average	Average	Larger than average or very large	DK		Below 2,500 grams ¹	Weighed at birth ²	
Total	3.7	6.3	85.4	2.8	1.8	100.0	8.2	38.8	387
Mother's age at birth									
Less than 20 years	(5.1)	(11.1)	(78.3)	(0.0)	(5.5)	100.0	(8.5)	(58.0)	52
20-34 years	2.6	5.8	86.7	3.3	1.5	100.0	7.6	37.5	277
35-49 years	7.6	4.0	85.8	2.6	0.0	100.0	10.5	28.1	59
Birth order									
1	4.3	10.2	78.2	1.6	5.6	100.0	8.1	56.6	62
2-3	1.5	5.7	90.7	1.3	0.8	100.0	7.2	35.9	145
4-5	2.2	5.7	88.7	1.8	1.6	100.0	7.5	33.5	104
6+	9.4	4.9	76.8	7.7	1.2	100.0	10.9	37.5	77
Area									
Urban	2.7	4.0	88.4	2.9	1.9	100.0	7.8	62.1	199
Rural	4.7	8.7	82.3	2.6	1.7	100.0	8.6	14.1	188
Mother's education									
None	3.9	7.3	84.9	1.9	2.0	100.0	8.3	26.6	268
Primary	1.3	5.8	88.2	2.6	2.1	100.0	6.9	58.2	82
Secondary+	(7.5)	(0.0)	(83.0)	(9.5)	(0.0)	100.0	(10.3)	(83.6)	38
Wealth index quintile									
Poorest	6.3	11.3	80.2	2.2	0.0	100.0	9.3	9.5	69
Second	6.5	6.7	85.3	0.0	1.5	100.0	9.8	4.7	86
Middle	1.9	5.7	90.2	1.1	1.1	100.0	7.5	40.8	84
Fourth	1.4	3.6	86.7	2.5	5.8	100.0	6.9	67.4	85
Richest	2.6	4.5	83.1	9.8	0.0	100.0	7.4	76.5	63
Ethnicity of household head									
Turkana	3.9	6.8	85.1	3.0	1.2	100.0	8.3	30.0	327
Other ethnic group	2.8	3.2	87.2	1.7	5.2	100.0	7.7	88.1	59
¹ MICS indicator 2.20 - Low-birthweight infants									
² MICS indicator 2.21 - Infants weighed at birth									
() Figures that are based on 25-49 unweighted cases									

4.2 Breastfeeding and Infant and Young Child Feeding

Proper feeding of infants and young children can increase their chances of survival; it can also promote optimal growth and development, especially in the critical window from birth to two years of age. Breastfeeding for the first two years of life protects children from infection, provides an ideal source of nutrients, and is economical and safe. However, many mothers do not start to breastfeed early enough, do not breastfeed exclusively for the recommended 6 months or stop breastfeeding too soon. There are often pressures to switch to infant formula, which can contribute to growth faltering and micronutrient deficiency. In addition, it can be unsafe if hygienic conditions, including safe drinking water are not readily available. Studies have shown that, in addition to continued breastfeeding, consumption of appropriate, adequate and safe solid, semi-solid and soft foods from the age of 6

months onwards leads to better health and growth outcomes, with potential to reduce stunting during the first two years of life.²⁸

UNICEF and WHO recommend that infants be initiated to breastfeeding within one hour of birth, breastfed exclusively for the first six months of life and continue to be breastfed up to 2 years of age and beyond.²⁹ Starting at 6 months, breastfeeding should be combined with safe, age-appropriate feeding of solid, semi-solid and soft foods.³⁰ A summary of key guiding principles^{31, 32} for feeding 6-23 month olds is provided in the Table NU2 below along with proximate measures for these guidelines collected in this survey.

The guiding principles for which proximate measures and indicators exist are:

- (i) continued breastfeeding;
- (ii) appropriate frequency of meals (but not energy density); and
- (iii) appropriate nutrient content of food.

Feeding frequency is used as proxy for energy intake, requiring children to receive a minimum number of meals/snacks (and milk feeds for non-breastfed children) for their age. Dietary diversity is used to ascertain the adequacy of the nutrient content of the food (not including iron) consumed. For dietary diversity, seven food groups were created for which a child consuming at least four of these is considered to have a better quality diet. In most populations, consumption of at least four food groups means that the child has a high likelihood of consuming at least one animal-source food and at least one fruit or vegetable, in addition to a staple food (grain, root or tuber).³³

These three dimensions of child feeding are combined into an assessment of the children who received appropriate feeding, using the indicator of “minimum acceptable diet”. To have a minimum acceptable diet in the previous day, a child must have received:

- (i) the appropriate number of meals/snacks/milk feeds;
- (ii) food items from at least 4 food groups; and
- (iii) breastmilk or at least 2 milk feeds (for non-breastfed children).

²⁸ Bhuta, Z. et al. 2013. *Evidence-based interventions for improvement of maternal and child nutrition: what can be done and at what cost?* The Lancet June 6, 2013.

²⁹ WHO. 2003. *Implementing the Global Strategy for Infant and Young Child Feeding*. Meeting Report Geneva, 3-5 February, 2003.

³⁰ WHO. 2003. *Global Strategy for Infant and Young Child Feeding*.

³¹ PAHO. 2003. *Guiding principles for complementary feeding of the breastfed child*.

³² WHO. 2005. *Guiding principles for feeding non-breastfed children 6-24 months of age*.

³³ WHO. 2008. *Indicators for assessing infant and young child feeding practices. Part 1: Definitions*.

Table NU.2: Guiding Principles for Feeding children age 6 – 23 months

Guiding Principle (age 6-23 months)	Proximate measures	Table
Continue frequent, on-demand breastfeeding for two years and beyond	Breastfed in the last 24 hours	NU.4
Appropriate frequency and energy density of meals	<p>Breastfed children Depending on age, two or three meals/snacks provided in the last 24 hours</p> <p>Non-breastfed children Four meals/snacks <u>and/or milk feeds</u> provided in the last 24 hours</p>	NU.6
Appropriate nutrient content of food	Four food groups ³⁴ eaten in the last 24 hours	NU.6
Appropriate amount of food	No standard indicator exists	na
Appropriate consistency of food	No standard indicator exists	na
Use of vitamin-mineral supplements or fortified products for infant and mother	No standard indicator exists	na
Practice good hygiene and proper food handling	While it was not possible to develop indicators to fully capture programme guidance, one standard indicator does cover part of the principle: Not feeding with a bottle with a nipple	NU.9
Practice responsive feeding, applying the principles of psycho-social care	No standard indicator exists	na

Table NU.3 is based on mothers' reports of what their last-born child, born in the last two years, was fed in the first few days of life. It indicates the proportion who were ever breastfed, those who were first breastfed within one hour and one day of birth, and those who received a prelacteal feed.³⁵

Ninety-seven percent of the children were ever breastfed (Table NU.3). However, although a very important step in management of lactation and establishment of a physical and emotional relationship between the baby and the mother, only 54 percent of babies were breastfed for the first time within one hour of birth and 73 percent of the newborns in Turkana County started breastfeeding within one day of birth. Babies delivered in a health facility were more likely to be breastfed within one hour of delivery compared to those delivered at home, (62 percent and 50 percent, respectively). Ten percent of the babies received prelacteal feed. Babies were more likely to receive prelacteal feed when delivered in a rural area, or delivered at home. Figure NU.1 shows the initiation of breastfeeding in Turkana County by urban rural areas.

³⁴ Food groups used for assessment of this indicator are 1) Grains, roots and tubers, 2) legumes and nuts, 3) dairy products (milk, yogurt, cheese), 4) flesh foods (meat, fish, poultry and liver/organ meats), 5) eggs, 6) vitamin-A rich fruits and vegetables, and 7) other fruits and vegetables.

³⁵ Prelacteal feed refers to the provision of any liquid or food, other than breastmilk, to a newborn during the period when breastmilk flow is generally being established (estimated here as the first 3 days of life).

Table NU.3: Initial breastfeeding

Percentage of last live-born children in the last two years who were ever breastfed, breastfed within one hour of birth, and within one day of birth, and percentage who received a prelacteal feed, Turkana County MICS, 2013/14

	Percentage who were ever breastfed ¹	Percentage who were first breastfed:		Percentage who received a prelacteal feed	Number of last live-born children in the last two years
		Within one hour of birth ²	Within one day of birth		
Total	97.2	54.0	72.9	9.8	387
Area					
Urban	97.6	53.4	79.4	7.6	199
Rural	96.8	54.6	66.1	12.2	188
Months since last birth					
0-11 months	97.7	55.4	72.8	7.9	208
12-23 months	96.6	52.5	73.0	12.1	179
Assistance at delivery					
Skilled attendant	95.6	63.1	80.1	9.0	136
Traditional birth attendant	(*)	(*)	(*)	(*)	15
Other	99.4	49.0	69.4	10.4	165
No one/Missing	94.6	47.8	65.1	11.7	71
Place of delivery					
Home	98.7	49.5	69.3	10.9	247
Health facility	95.6	62.3	80.3	8.4	135
Mother's education					
None	98.2	53.5	72.2	9.1	268
Primary	93.8	59.6	77.1	7.9	82
Secondary+	(97.3)	(45.9)	(69.0)	(19.6)	38
Wealth index quintile					
Poorest	98.2	44.2	53.2	12.7	69
Second	99.2	58.4	72.5	12.1	86
Middle	96.2	61.5	78.2	10.6	84
Fourth	95.6	47.4	76.6	6.0	85
Richest	96.8	57.8	82.9	7.7	63
Ethnicity of household head					
Turkana	98.1	52.8	71.0	10.7	327
Other ethnic group	92.1	60.5	83.0	5.5	59


¹ MICS indicator 2.5 - Children ever breastfed

² MICS indicator 2.6 - Early initiation of breastfeeding

() Figures that are based on 25-49 unweighted cases

(*) Figures that are based on fewer than 25 unweighted cases

Figure NU.1: Initiation of breastfeeding, Turkana County MICS, 2013/14



The set of Infant and Young Child Feeding indicators reported in Tables NU.4 through NU.8 are based on the mother’s report of consumption of food and fluids during the day or night prior to being interviewed. Data are subject to a number of limitations, some related to the mother’s ability to provide a full report on the child’s liquid and food intake due to recall errors as well as lack of knowledge in cases where the child was fed by other individuals.

In Table NU.4, breastfeeding status is presented for both *Exclusively breastfed* and *Predominantly breastfed*; referring to infants age less than 6 months who are breastfed, distinguished by *the former* only allowing vitamins, mineral supplements, and medicine and *the latter* allowing also plain water and non-milk liquids. The table also shows continued breastfeeding of children at 12-15 and 20-23 months of age.

Table NU.4: Breastfeeding

Percentage of living children according to breastfeeding status at selected age groups, Turkana County MICS, 2013/14							
	Children age 0-5 months			Children age 12-15 months		Children age 20-23 months	
	Percent exclusively breastfed ¹	Percent predominantly breastfed ²	Number of children	Percent breastfed (Continued breastfeeding at 1 year) ³	Number of children	Percent breastfed (Continued breastfeeding at 2 years) ⁴	Number of children
Total	68.5	86.1	107	91.8	68	60.5	61
Sex							
Male	74.9	88.7	58	(93.2)	29	(58.4)	36
Female	61.0	83.1	49	(90.8)	39	(63.5)	25
Area							
Urban	(70.9)	(92.9)	52	(*)	32	(56.3)	36
Rural	66.3	79.7	55	(90.1)	37	(66.5)	25
	¹ MICS indicator 2.7 - Exclusive breastfeeding under 6 months						
	² MICS indicator 2.8 - Predominant breastfeeding under 6 months						
	³ MICS indicator 2.9 - Continued breastfeeding at 1 year						
	⁴ MICS indicator 2.10 - Continued breastfeeding at 2 years						
	() Figures that are based on 25-49 unweighted cases						
	(*) Figures that are based on fewer than 25 unweighted cases						

Table NU.5 shows the median duration of breastfeeding by selected background characteristics. Among children under age 3 years, the median duration for ever breastfed was 22 months, 5 months for exclusive breastfeeding, and 7 months for predominant breastfeeding.

Table NU.5: Duration of breastfeeding

Median duration of any breastfeeding, exclusive breastfeeding, and predominant breastfeeding among children age 0-35 months, Turkana County MICS, 2013/14

	Median duration (in months) of:			Number of children age 0-35 months
	Any breastfeeding ¹	Exclusive breastfeeding	Predominant breastfeeding	
Median	21.8	4.6	7.0	627
Sex				
Male	22.6	5.2	7.2	318
Female	21.6	3.7	6.6	309
Area				
Urban	21.2	4.6	6.6	321
Rural	23.2	4.5	7.6	306
Mother's education				
None	22.4	4.5	7.2	442
Primary	19.6	5.1	6.8	130
Secondary+	(22.7)	(0.6)	(6.0)	54
Wealth index quintile				
Poorest	24.7	4.9	9.5	125
Second	22.3	4.9	7.5	143
Middle	21.6	4.8	7.5	134
Fourth	20.5	5.0	6.6	130
Richest	20.9	1.9	4.1	94
Ethnicity of household head				
Turkana	22.1	4.9	7.2	528
Other ethnic group	19.3	2.7	5.5	98
Mean	22.5	6.0	10.0	627
¹ MICS indicator 2.11 - Duration of breastfeeding				
() Figures that are based on 25-49 unweighted cases				

The age-appropriateness of breastfeeding of children under age 24 months is provided in Table NU.6. Different criteria of feeding were used depending on the age of the child. For infants age 0-5 months, exclusive breastfeeding was considered as age-appropriate feeding, while children age 6-23 months were considered to be appropriately fed if they were receiving breastmilk and solid, semi-solid or soft food. As a result of feeding patterns in Turkana County, only 52 percent of children age 0-23 months are being appropriately breastfed and age-appropriate breastfeeding among all children age 6-23 months is 47 percent. Age appropriate breastfeeding is 54 percent for girls and 51 percent for boys.

Table NU.6: Age-appropriate breastfeeding

Percentage of children age 0-23 months who were appropriately breastfed during the previous day, Turkana County MICS, 2013/14

	Children age 0-5 months		Children age 6-23 months		Children age 0-23 months	
	Percent exclusively breastfed ¹	Number of children	Percent currently breastfeeding and receiving solid, semi-solid or soft foods	Number of children	Percent appropriately breastfed ²	Number of children
Total	68.5	107	46.9	315	52.3	422
Sex						
Male	74.9	58	41.6	160	50.5	218
Female	61.0	49	52.3	155	54.4	204
Area						
Urban	(70.9)	52	46.2	164	52.1	216
Rural	66.3	55	47.6	151	52.6	206
Mother's education						
None	65.2	77	47.7	215	52.3	293
Primary	(*)	25	41.9	67	52.2	92
Secondary+	(*)	5	(54.1)	31	(54.9)	36
Wealth index quintile						
Poorest	(65.3)	18	35.8	60	42.5	78
Second	(71.4)	28	55.8	65	60.4	93
Middle	(*)	26	41.0	71	49.2	97
Fourth	(*)	25	52.8	63	56.9	88
Richest	(*)	11	(49.0)	55	51.2	66
Ethnicity of household head						
Turkana	71.2	92	47.0	265	53.2	357
Other ethnic group	(*)	15	(45.8)	50	47.2	65
¹ MICS indicator 2.7 - Exclusive breastfeeding under 6 months						
² MICS indicator 2.12 - Age-appropriate breastfeeding						
() Figures that are based on 25-49 unweighted cases						
(*) Figures that are based on fewer than 25 unweighted cases						

Overall, 31 percent of infants age 6-8 months received solid, semi-solid, or soft foods at least once during the previous day (Table NU.7).³⁶ The same percentage was noted among currently breastfeeding infants.

Table NU.7: Introduction of solid, semi-solid, or soft foods

Percentage of infants age 6-8 months who received solid, semi-solid, or soft foods during the previous day, Turkana County MICS, 2013/14

	Currently breastfeeding		All	
	Percent receiving solid, semi-solid or soft foods	Number of children age 6-8 months	Percent receiving solid, semi-solid or soft foods ¹	Number of children age 6-8 months
Total	31.2	56	31.2	56
¹ MICS indicator 2.13 - Introduction of solid, semi-solid or soft foods				

³⁶ Descriptions by rural/urban areas and sex of child were not done due to small numbers of respondents in those categories.

Overall, only 18 percent of the children age 6-23 months were receiving solid, semi-solid and soft foods the minimum number of times (Table NU.8).³⁷ The proportion of children receiving the minimum dietary diversity, or foods from at least four food groups, was much lower than that for the minimum meal frequency, indicating the need to focus on improving diet quality and nutrient intake among this vulnerable group. The overall assessment using the indicator of minimum acceptable diet revealed that only three percent were benefitting from a diet sufficient in both diversity and frequency.

Table NU.8: Infant and young child feeding (IYCF) practices

Percentage of children age 6-23 months who received appropriate liquids and solid, semi-solid, or soft foods the minimum number of times or more during the previous day, by breastfeeding status, Turkana County MICS, 2013/14

	All			
	Percent of children who received:			
	Minimum dietary diversity ^{4, a}	Minimum meal frequency ^{5, b}	Minimum acceptable diet ^c	Number of children age 6-23 months
Total	9.5	18.1	3.4	315
Sex				
Male	11.9	20.3	4.3	160
Female	7.0	15.8	2.4	155
Age				
6-8 months	0.0	20.6	0.0	56
9-11 months	8.8	20.2	4.3	64
12-17 months	10.4	18.0	3.6	105
18-23 months	14.7	15.2	4.5	90
Area				
Urban	13.5	16.6	4.6	164
Rural	5.1	19.8	2.1	151
Mother's education				
None	7.1	17.1	2.7	215
Primary	16.0	20.9	4.6	67
Secondary+	(12.3)	(20.3)	(5.4)	31
Wealth index quintile				
Poorest	3.4	13.0	0.0	60
Second	3.5	20.0	2.6	65
Middle	8.3	13.7	2.0	71
Fourth	15.1	23.1	7.5	63
Richest	(18.3)	(21.4)	(5.1)	55
Ethnicity of household head				
Turkana	8.5	16.8	3.4	265
Other ethnic group	(13.8)	(24.3)	(2.2)	50
¹ MICS indicator 2.17a - Minimum acceptable diet (breastfed)				
² MICS indicator 2.17b - Minimum acceptable diet (non-breastfed)				
³ MICS indicator 2.14 - Milk feeding frequency for non-breastfed children				
⁴ MICS indicator 2.16 - Minimum dietary diversity				
⁵ MICS indicator 2.15 - Minimum meal frequency				

³⁷ Note that a comparison between children 6-23 months currently breastfeeding and those currently not breastfeeding was removed from Table NU.8 because a high proportion of children were currently breastfeeding.

^a Minimum dietary diversity is defined as receiving foods from at least 4 of 7 food groups: 1) Grains, roots and tubers, 2) legumes and nuts, 3) dairy products (milk, yogurt, cheese), 4) flesh foods (meat, fish, poultry and liver/organ meats), 5) eggs, 6) vitamin-A rich fruits and vegetables, and 7) other fruits and vegetables.

^b Minimum meal frequency among currently breastfeeding children is defined as children who also received solid, semi-solid, or soft foods 2 times or more daily for children age 6-8 months and 3 times or more daily for children age 9-23 months. For non-breastfeeding children age 6-23 months it is defined as receiving solid, semi-solid or soft foods, or milk feeds, at least 4 times.

^c The minimum acceptable diet for breastfed children age 6-23 months is defined as receiving the minimum dietary diversity and the minimum meal frequency, while it for non-breastfed children further requires at least 2 milk feedings and that the minimum dietary diversity is achieved without counting milk feeds.

() Figures that are based on 25-49 unweighted cases

The continued practice of bottle-feeding is a concern because of the possible contamination due to unsafe water and lack of hygiene in preparation. Table NU.9 shows that bottle-feeding is practised for children under two years of age in Turkana County. About 10 percent of children under 6 months are fed using a bottle with a nipple. The findings indicate that bottle feeding becomes prevalent mainly from 6 months of age and older.

Table NU.9: Bottle feeding		
Percentage of children age 0-23 months who were fed with a bottle with a nipple during the previous day, Turkana County MICS, 2013/14		
	Percentage of children age 0-23 months fed with a bottle with a nipple ¹	Number of children age 0-23 months
Total	9.7	422
Sex		
Male	6.6	218
Female	13.1	204
Age		
0-5 months	3.1	107
6-11 months	13.7	120
12-23 months	10.9	196
Area		
Urban	11.6	216
Rural	7.8	206
Mother's education		
None	9.4	293
Primary	9.6	92
Secondary+	13.0	36
Wealth index quintile		
Poorest	8.1	78
Second	5.2	93
Middle	7.5	97
Fourth	10.5	88
Richest	20.3	66
Ethnicity of household head		
Turkana	8.8	357
Other ethnic group	14.7	65
¹ MICS indicator 2.18 - Bottle feeding		

4.3 Salt Iodization

Iodine Deficiency Disorders (IDD) is the world's leading cause of preventable mental retardation and impaired psychomotor development in young children. In its most extreme form, iodine deficiency causes cretinism. It also increases the risks of stillbirth and miscarriage in pregnant women. Iodine deficiency is most commonly and visibly associated with goitre. IDD takes its greatest toll in impaired mental growth and development, contributing in turn to poor school performance, reduced intellectual ability, and impaired work performance. The indicator is the percentage of households consuming adequately iodized salt (≥ 15 parts per million).

The IDD legislation passed in Kenya in 1978 (revised in 1988) covers all salt produced for human consumption. Specifications for edible salt are reviewed regularly (latest revision was in September 2000) by the Kenya Bureau of Standards. Iodization of salt is mandatory. The mandated level of iodization is 168.5 mg/kg of salt, or 100ppm.³⁸ The Ministry of Health monitors IDD in the country.

In 64 percent of households in Turkana County, salt used for cooking was tested for iodine content by using salt test kits and testing for the presence of potassium iodate content. Table NU.10 shows that in 36 percent of households, there was no salt available. These households were included in the denominator of the indicator. In 63 percent of households, salt was found to contain at least 15 parts per million (ppm) or more of iodine. The percentage of households using iodized salt is low in Turkana compared to Bungoma and Kakamega Counties as per MICS5. The Kenya Demographic and Health Survey 2014 also indicates that there was no salt in 34 percent of households in Turkana.

A cost of diet analysis conducted by Save the Children³⁹ under their food security program in 2013 indicates that Turkana community largely depend on readily available local food (milk, blood and meat mostly taken raw or roasted) due to their pastoral/nomadic lifestyle, and they incur minimal expenditure on purchased food or other items including sugar and salt.

³⁸ <http://www.tulane.edu/~internut/Countries/Kenya/kenyaiodine.html>


³⁹ Save the Children with support from Oxfam, 2013. A Cost of the Diet analysis in Turkana district of Kenya. Kerio Agropastoral livelihood zone, Turkana. Funded by ECHO.

Table NU.10: Iodized salt consumption

Percent distribution of households by consumption of iodized salt, Turkana County MICS, 2013/14								
	Percentage of households in which salt was tested	Number of households	Percent of households with:				Total	Number of households in which salt was tested or with no salt
			No salt	Salt test result				
				Not iodized 0 PPM	>0 and <15 PPM	15+ PPM ¹		
Total	63.6	1,277	35.6	0.1	0.9	63.4	100	1,260
Area								
Urban	76.0	684	22.7	0.0	1.0	76.3	100	672
Rural	49.2	593	50.2	0.2	0.8	48.7	100	587
Wealth index quintile								
Poorest	29.8	256	69.9	0.2	0.0	29.9	100	254
Second	48.5	282	51.3	0.1	0.4	48.1	100	281
Middle	65.5	255	33.7	0.2	1.6	64.5	100	252
Fourth	84.3	239	12.9	0.0	1.9	85.2	100	232
Richest	93.9	245	4.8	0.0	0.9	94.3	100	242

¹ MICS indicator 2.19 - Iodized salt consumption

The consumption of adequately iodized salt is graphically presented in Figure NU.2 together with the percentage of salt containing less than 15 ppm. In urban areas, 76 percent of households had salt that contained at least 15ppm compared to 49 percent in rural areas. The proportion of households with salt that contained at least 15ppm ranged from 30 percent in the poorest households to 94 percent in the richest.

Figure NU.2: Consumption of iodized salt, Turkana County MICS, 2013/14

5. Child Health

Kenya has acceded and ratified a number of major international and regional conventions some of which aim at ensuring child survival, growth and development. In 1990, Kenya ratified the United Nations Convention on the rights of the Child (CRC).^{40, 41} Article 6 of the CRC refers to the right to life, survival and development. The term ‘development’ in this context refers to physical, mental, emotional, cognitive, social and cultural development. Further, Article 24 states that ‘children have the right to good quality health care – the best health care possible – to safe drinking water, nutritious food, a clean and safe environment, and information to help them stay healthy’.⁴² The United Nations Millennium Declaration, signed in September 2000, commits world leaders to combat poverty, hunger, disease, illiteracy, environmental degradation, and discrimination against women. The objective of one of the Millennium Development Goals (MDGs) – MDG 4 - is to reduce child mortality by two thirds between 1990 and 2015. The Constitution of Kenya (2010) states that every person has the right to the highest attainable standard of health, which includes the right to health care services, including reproductive health care.

Chapter Five focuses on the following subtopics: vaccinations; neonatal tetanus protection; and care of illnesses (diarrhoea, acute respiratory infections, malaria/fever); and use of solid fuels.

5.1 Vaccinations

Immunization plays a key part in reducing preventable child diseases and mortality. The Global Vaccine Action Plan (GVAP) was endorsed by the 194 Member States of the World Health Assembly in May 2012 to achieve the Decade of Vaccines vision by delivering universal access to immunization. Immunization has saved the lives of millions of children in the four decades since the launch of the Expanded Programme on Immunization (EPI) in 1974. Worldwide there are still millions of children not reached by routine immunization and as a result, vaccine-preventable diseases cause more than 2 million deaths every year.

The WHO Recommended Routine Immunizations for Children⁴³ states that all children to be vaccinated against tuberculosis, diphtheria, pertussis, tetanus, polio, measles, hepatitis B, haemophilus influenza type b, pneumonia/meningitis, rotavirus, and rubella.

All doses in the primary series are recommended to be completed before the child’s first birthday, although depending on the epidemiology of disease in a country, the first doses of measles and rubella containing vaccines may be recommended at 12 months or later. The recommended number and timing of most other doses also vary slightly with local epidemiology and may include booster doses later in childhood.

⁴⁰Kenya Human Rights Commission. 2010. Towards Equality and Anti-Discrimination: An Overview of International and Domestic Law an Anti-discrimination in Kenya.

⁴¹The Kenyan Section of the International Commission of Jurists. 2004. International Human Rights Standards: Reporting Obligations – The Convention of the Rights of the Child.

⁴²The United Nations General Assembly. 1989. The Convention on the Rights of the Child.

⁴³<http://www.who.int/immunization/diseases/en>. Table 2 includes recommendations for all children and additional antigens recommended only for children residing in certain regions of the world or living in certain high-risk population groups.

The Kenya Expanded Programme on Immunization (KEPI) was established in 1980 and is integrated within the Department of Preventive and Promotive Health Services of the Ministry of Health as part of the Essential Health Package (EHP). KEPI is now known as the Division of Vaccine and Immunisation (DVI). The Kenya National Immunization Programme immunization schedule is shown below. All vaccines should be received during the first year of life except the second dose of measles given at 18 months. Yellow fever is given at 9 months to children in selected sub-counties in the former Rift Valley province.⁴⁴

Child Immunization Schedule in Kenya^{45, 46}

Vaccine	Age	Remarks
BCG Vaccine: at birth		Intra-dermal left forearm; BCG Scar checked
Dose: (0.05mls)	Below 1 year	
Dose: (0.1mls)	Above 1 year	
Oral Polio Vaccine (OPV)		2 drops (orally)
Birth dose: OPV 0	At birth or within 2 weeks	
1 st dose: OPV 1	At 6 weeks	
2 nd dose: OPV 2	At 10 weeks	
3 rd dose: OPV 3	At 14 weeks	
Diphtheria/Pertussis/Tetanus/Hepatitis B/haemophilus influenzae Type b		0.5mls (intra-muscular left outer thigh)
1 st dose	6 weeks	
2 nd dose	10 weeks	
3 rd dose	14 weeks	
Pneumococcal Vaccine		0.5mls (intra-muscular right outer thigh)
1 st dose	6 weeks	
2 nd dose	10 weeks	
3 rd dose	14 weeks	
Rota Virus (Rotarix)		1.5mls (orally)
1 st dose	6 weeks	
2 nd dose	10 weeks	
Measles Vaccine at 6 months: in the event of measles outbreak or HIV exposed children (HEI)	6 months	0.5mls (Subcutaneously right upper arm)
Measles Vaccine	9 months	
Measles Vaccine	18 months	
Yellow Fever	9 months	0.5mls (Intra-muscular left upper deltoid)
Other Vaccines		Other vaccines refer to those not in the usual KEPI schedule and may include MMR, Typhoid, etc.

⁴⁴ MICS 2013/14 collected data on Yellow Fever but further analysis is required before the findings can be shared.

⁴⁵Ministry of Health, 2013. Mother and Child Health Booklet. Republic of Kenya

⁴⁶Kenya is planning to carry out a Measles-Rubella (MR) and IPV Campaign in 2016, and subsequently include MR in the child immunization schedule in 2017.

In Turkana County, the MICS collected data on immunization coverage for all children under three years of age. All mothers or caretakers were asked to provide vaccination cards. If the immunization card for a child was available, interviewers copied vaccination information from the cards onto the MICS questionnaire. If no immunization card was available for the child, the interviewer proceeded to ask the mother to recall whether or not the child had received each of the vaccines as per the schedule. The final immunization coverage estimates are based on information obtained from the immunization card and/or the mother's report.

The percentage of children age 12-23 months and 24-35 months who had received each of the specific vaccines by source of information (immunization card and mother's recall) is shown in Table CH.1 and Figure CH.1. The denominators for the table are comprised of children age 12-23 months and 24-35 months and only children in these age groups are counted. In the first three columns in each panel of the table, the numerator includes all children who were vaccinated at any time before the survey according to the immunization card or the mother's report. In the last column in each panel, only those children who were fully immunized before their first birthday, as recommended, were included. The proportion of children immunized before the first birthday but without immunization card/record was assumed to be the same as for those with vaccination cards/records.

Most children age 12-23 months had been vaccinated against BCG and measles by the age of 12 months (96 and 81 percent, respectively), and had received the first dose of DPT, HepB, and Hib vaccines (96 percent, 94 percent and 94 percent, respectively). The percentages decline for the second and third doses of DPT, HepB, and Hib. Similarly, 96 percent of children age 12-23 months had received Polio 1 by age 12 months and this declined to 84 percent by the third dose. As a result, the percentage of children 12-23 months of age who had been fully vaccinated by their first birthday was 64 percent. The proportion of children fully vaccinated by 12 months of age was lower for children age 24-35 months (42 percent). The individual coverage figures for children age 24-35 months are generally lower to those age 12-23 months suggesting that immunization coverage has been on average improving in Turkana County between 2011 and 2013.

Table CH.1: Vaccinations in the first years of life

Percentage of children age 12-23 months and 24-35 months vaccinated against vaccine preventable childhood diseases at any time before the survey and by their first birthday, Turkana County MICS, 2013/14

	Children age 12-23 months:				Children age 24-35 months:			
	Vaccinated at any time before the survey according to:			Vaccinated by 12 months of age ^a	Vaccinated at any time before the survey according to:			Vaccinated by 12 months of age
	Vaccination card	Mother's report	Either		Vaccination card	Mother's report	Either	
Antigen								
BCG ¹	63.3	34.1	97.4	96.4	41.0	53.8	94.8	85.9
Polio								
At birth	62.3	25.2	87.5	84.7	39.7	44.4	84.1	73.6
1	64.6	32.2	96.7	95.8	41.5	53.0	94.4	87.0
2	64.2	27.5	91.7	89.4	41.5	47.0	88.5	79.9
3 ²	64.2	22.7	86.9	84.1	41.5	31.5	73.0	66.2
DPT								
1	64.3	32.5	96.8	95.8	41.8	52.7	94.5	87.1
2	64.0	31.8	95.8	93.4	41.8	52.3	94.1	84.9
3 ³	64.0	29.5	93.5	90.5	41.8	47.1	88.9	80.6
HepB								
At birth	61.7	21.0	82.6	77.8	40.0	31.8	71.8	62.5
1	64.3	30.6	95.0	94.0	41.8	50.3	92.1	84.9
2	64.0	31.0	95.0	92.6	41.8	47.0	88.8	80.2
3 ⁴	64.0	18.3	82.2	79.6	41.8	29.1	70.9	64.3
Hib								
1	64.7	31.8	96.5	94.0	42.0	53.0	95.0	87.0
2	64.3	30.2	94.6	91.0	42.0	52.2	94.2	84.4
3 ⁵	64.3	27.7	92.0	86.2	42.0	47.5	89.5	77.0
Measles (MCV1) ⁷	60.8	31.7	92.5	81.3	42.2	49.3	91.5	67.5
Fully vaccinated ^{8, b}	64.3	14.5	78.8	63.6	42.7	21.3	64.0	42.1
No vaccinations	0.0	2.6	2.6	3.6	0.0	4.5	4.5	8.3
Number of children	196	196	196	196	205	205	205	205

¹ MICS indicator 3.1 - Tuberculosis immunization coverage

² MICS indicator 3.2 - Polio immunization coverage

³ MICS indicator 3.3 - Diphtheria, pertussis and tetanus (DPT) immunization coverage

⁴ MICS indicator 3.5 - Hepatitis B immunization coverage

⁵ MICS indicator 3.6 - Haemophilus influenzae type B (Hib) immunization coverage

⁶ MICS indicator 3.7 - Yellow fever immunization coverage⁴⁷

⁷ MICS indicator 3.4; MDG indicator 4.3 - Measles immunization coverage

⁸ MICS indicator 3.8 - Full immunization coverage

^aAll MICS indicators refer to results in this column

^b Includes: BCG, Polio3, DPT3, HepB3, Hib3, and Measles (MCV1) as per the vaccination schedule in Kenya

⁴⁷ Yellow fever immunization coverage was not included in the analysis

Figure CH.1: Vaccinations by age 12 months (measles by 24 months), Turkana County MICS, 2013/14




Table CH.2 presents vaccination coverage estimates among children age 12-23 and 24-35 months by background characteristics. The figures indicate children receiving the vaccinations at any time up to the date of the survey, and are based on information from both the vaccination cards and mothers'/caretakers' reports. Vaccination cards were seen by the interviewers for only 64 percent of children age 12-23 months. Overall, 79 percent of children age 12-23 months were fully immunized against vaccine preventable childhood diseases.

Table CH.2: Vaccinations by background characteristics

Percentage of children age 12-23 months currently vaccinated against vaccine preventable childhood diseases, Turkana County MICS, 2013/14

	Percentage of children who received:																		Percentage with vaccination card seen	Number of children age 12-23 months
	Polio					DPT			HepB			Hib			Measles (MCV1)	Full ^a	None			
	BCG	At birth	1	2	3	1	2	3	At birth	1	2	3	1	2				3		
Total	97.4	87.5	96.7	91.7	86.9	96.8	95.8	93.5	82.6	95.0	95.0	82.2	96.5	94.6	92.0	92.5	78.8	2.6	64.2	196
Sex																				
Male	97.8	87.3	96.5	90.0	87.1	96.7	95.5	92.1	80.0	93.6	93.6	81.9	96.7	93.6	89.8	92.7	78.7	2.2	58.7	101
Female	96.9	87.7	96.9	93.4	86.8	96.9	96.2	95.0	85.5	96.4	96.4	82.6	96.4	95.6	94.4	92.3	78.9	3.1	70.0	95
Area																				
Urban	100.0	95.1	100.0	94.1	93.0	100.0	100.0	98.6	87.5	98.2	98.2	88.1	100.0	98.2	95.9	96.0	83.0	0.0	65.4	105
Rural	94.3	78.9	93.0	89.0	80.0	93.1	91.0	87.7	77.0	91.2	91.2	75.4	92.6	90.4	87.6	88.5	74.0	5.7	62.7	91

^a Includes: BCG, Polio3, DPT3, HepB3, Hib3, and Measles (MCV1) as per the vaccination schedule in Kenya

5.2 Neonatal Tetanus Protection

The goal of MDG 5 is to reduce by three quarters the maternal mortality ratio, with one strategy to eliminate maternal tetanus. Following on the 42nd and 44th World Health Assembly calls for elimination of neonatal tetanus, the global community continues to work to reduce the incidence of neonatal tetanus to less than one case per 1,000 live births in every sub-county by 2015.

The strategy for preventing maternal and neonatal tetanus is to ensure that all pregnant women receive at least two doses of tetanus toxoid vaccine. If a woman has not received at least two doses during a particular pregnancy, the mother and child are also considered to be protected against tetanus if the woman:

- Received at least two doses of tetanus toxoid vaccine, the last within the previous 3 years;
- Received at least 3 doses, the last within the previous 5 years;
- Received at least 4 doses, the last within the previous 10 years;
- Received 5 or more doses anytime during her life.

To assess the status of tetanus vaccination coverage in Turkana County, women who had a live birth during the two years before the survey were asked if they had received tetanus toxoid injections during the pregnancy for their most recent birth, and if so, how many. Women who did not receive two or more tetanus toxoid vaccinations during this recent pregnancy were then asked about tetanus toxoid vaccinations they may have previously received. Interviewers also asked women to present their vaccination card on which dates of tetanus toxoid are recorded and referred to information from the cards when available.

Table CH.3 shows the protection status from tetanus of women age 15-49 years who have had a live birth within the last two years preceding the survey. In Turkana, 45 percent of these women were protected against neonatal tetanus.

Table CH.3: Neonatal tetanus protection							
Percentage of women age 15-49 years with a live birth in the last 2 years protected against neonatal tetanus, Turkana County MICS, 2013/14							
	Percentage of women who received at least 2 doses during last pregnancy	Percentage of women who did not receive two or more doses during last pregnancy but received:				Protected against tetanus ¹	Number of women with a live birth in the last 2 years
		2 doses, the last within prior 3 years	3 doses, the last within prior 5 years	4 doses, the last within prior 10 years	5 or more doses during lifetime		
Total	29.5	15.0	0.0	0.2	0.0	44.7	387
Area							
Urban	30.7	15.1	0.0	0.0	0.0	45.8	199
Rural	28.2	14.9	0.0	0.5	0.0	43.5	188
Education							
None	27.0	13.7	0.0	0.3	0.0	41.1	268
Primary	28.5	17.4	0.0	0.0	0.0	45.9	82
Secondary+	(48.8)	(19.1)	(0.0)	(0.0)	(0.0)	(67.9)	38
Wealth index quintile							
Poorest	20.2	14.2	0.0	1.2	0.0	35.7	69
Second	24.2	15.9	0.0	0.0	0.0	40.1	86
Middle	27.7	10.2	0.0	0.0	0.0	37.9	84
Fourth	46.0	11.6	0.0	0.0	0.0	57.7	85
Richest	27.0	25.5	0.0	0.0	0.0	52.5	63
Ethnicity of household head							
Turkana	29.9	14.6	0.0	0.3	0.0	44.7	327
Other ethnic group	26.7	17.5	0.0	0.0	0.0	44.3	59
¹ MICS indicator 3.9 - Neonatal tetanus protection							
() Figures that are based on 25-49 unweighted cases							

5.3 Care of Illness

A key strategy for accelerating progress toward MDG 4 is to tackle the diseases that are the leading causes of morbidity and mortality of children under-5 years. Diarrhoea and pneumonia are two such diseases. The Global Action Plan for the Prevention and Control of Pneumonia and Diarrhoea (GAPPD) aims to end preventable pneumonia and diarrhoea death by reducing mortality from pneumonia to 3 deaths per 1,000 live births and mortality from diarrhoea to 1 death per 1,000 live births by 2025. Malaria is also a major cause of mortality of children under-5 years, leading to about 1,200 deaths children every day, especially in sub-Saharan Africa.⁴⁸

Table CH.4 presents the percentage of children under-5 years of age who were reported to have had an episode of diarrhoea, symptoms of acute respiratory infection (ARI), or fever during the two weeks preceding the survey. These results measure period-prevalence of those illnesses over a two-week time window.

⁴⁸UNICEF Fact sheet http://www.unicef.org/media/media_81674.html

The definition of a case of diarrhoea or fever, in this survey, was the mother's or caretaker's report that the child had such symptoms over the specified period; no other evidence was sought beside the opinion of the mother. A child was considered to have had an episode of ARI if the mother or caretaker reported that the child had, over the specified period, an illness with a cough with rapid or difficult breathing, and whose symptoms were perceived to be due to a problem in the chest or both a problem in the chest and a blocked nose. While this approach is reasonable in the context of a MICS, these basically simple case definitions must be kept in mind when interpreting the results, as well as the potential for reporting and recall biases. Further, diarrhoea, fever and ARI are not only seasonal but are also characterized by the often rapid spread of localized outbreaks from one area to another at different points in time.

Table CH.4: Reported disease episodes				
Percentage of children age 0-59 months for whom the mother/caretaker reported an episode of diarrhoea, symptoms of acute respiratory infection (ARI), and/or fever in the last two weeks, Turkana County MICS, 2013/14				
	Percentage of children who in the last two weeks had:			Number of children age 0-59 months
	An episode of diarrhoea	Symptoms of ARI	An episode of fever	
Total	17.4	5.1	18.6	1,067
Sex				
Male	17.6	4.7	17.6	537
Female	17.2	5.6	19.5	530
Area				
Urban	19.0	5.2	20.8	546
Rural	15.7	5.0	16.2	521
Age				
0-11 months	17.9	7.8	17.8	227
12-23 months	26.2	5.5	26.5	196
24-35 months	20.1	2.5	17.4	205
36-47 months	11.9	4.1	17.0	222
48-59 months	12.0	5.5	14.8	217
Mother's education				
None	17.2	5.0	17.2	758
Primary	18.2	3.9	21.5	207
Secondary+	16.1	8.9	23.3	99
Wealth index quintile				
Poorest	18.3	3.8	13.7	216
Second	16.5	5.3	15.4	244
Middle	16.3	6.7	22.1	217
Fourth	16.4	5.9	20.7	227
Richest	20.3	3.4	22.1	163
Ethnicity of household head				
Turkana	17.4	5.3	18.2	898
Other ethnic group	17.5	4.2	20.7	167

In Turkana, 17 percent of children under-5 years of age are reported to have had diarrhoea in the two weeks preceding the survey, five percent symptoms of ARI, and 19 percent an episode of fever (Table CH.4). About 19 percent of children under-5 years in urban areas had experienced an episode of diarrhoea compared to 16 percent in rural areas. Reported episodes of fever were 21 percent in urban areas and 16 percent in rural areas.

5.3.1 Diarrhoea

Diarrhoea is one of the leading causes of death among children under five worldwide.⁴⁹ Most diarrhoea-related deaths in children are due to dehydration from loss of large quantities of water and electrolytes from the body in liquid stools. Management of diarrhoea – either through oral rehydration salts (ORS) or a recommended home fluid (RHF) – can prevent many of these deaths. In addition, provision of zinc supplements has been shown to reduce the duration and severity of the illness as well as the risk of future episodes within the next two or three months. Preventing dehydration and malnutrition by increasing fluid intake and continuing to feed the child are also important strategies for managing diarrhoea.

During the survey, mothers or caretakers were asked whether their child under five years had an episode of diarrhoea in the two weeks prior to the survey. In cases where mothers reported that the child had diarrhoea, a series of questions were asked about the treatment of the illness, including what the child had been given to drink and eat during the episode and whether this was more or less than what was usually given to the child.

The overall period-prevalence of diarrhoea in children under-5 years of age is 17 percent (Table CH.4). The highest period-prevalence is seen among children age 12-23 months (26 percent).

Table CH.5 shows the percentage of children with diarrhoea in the two weeks preceding the survey for whom advice or treatment was sought and where. Overall, a health facility or provider was seen in 70 percent of cases, predominantly in public health facilities (62 percent). However, no advice or treatment was sought in 27 percent of the cases.

⁴⁹WHO, 2013. Fact Sheet number 330.

Table CH.5: Care-seeking during diarrhoea

Percentage of children age 0-59 months with diarrhoea in the last two weeks for whom advice or treatment was sought, by source of advice or treatment, Turkana County MICS, 2013/14

	Percentage of children with diarrhoea for whom:						Number of children age 0-59 months with diarrhoea in the last two weeks
	Advice or treatment was sought from:						
	Health facilities or providers		Other source	A health facility or provider ^{1, b}	No advice or treatment sought		
	Public	Private					
Total	62.0	9.0	1.7	70.1	27.3	186	
Sex							
Male	59.4	7.4	3.4	65.1	29.7	94	
Female	64.6	10.7	0.0	75.3	24.7	91	
Area							
Urban	52.9	14.2	2.8	65.5	30.1	104	
Rural	73.5	2.4	0.4	76.0	23.6	82	
¹ MICS indicator 3.10 - Care-seeking for diarrhoea							
^a Community health providers includes both public (<i>Community health worker and Mobile/Outreach clinic</i>) and private (<i>Mobile clinic</i>) health facilities ⁵⁰							
^b Includes all public and private health facilities and providers, but excludes private pharmacy							

Table CH.6 provides information on drinking and feeding practices during diarrhoea. Overall, 14 percent of under-5 children who experienced an episode of diarrhoea in the last two weeks preceding the survey were given more than usual to drink while 15 percent were given about the same. About 27 percent were given somewhat less, but 40 percent were given much less than usual.

About six percent of children under five years of age who had an episode of diarrhoea in the last two weeks preceding the survey were given more to eat than usual while 11 percent were given about the same quantity of food. Thirty-six percent were given somewhat less to eat and 32 percent were given much less during this period.

⁵⁰Community health worker category removed for lack of cases

Table CH.6: Feeding practices during diarrhoea

Percent distribution of children age 0-59 months with diarrhoea in the last two weeks by amount of liquids and food given during episode of diarrhoea, Turkana County MICS, 2013/14

	Drinking practices during diarrhoea						Eating practices during diarrhoea						Number of children age 0-59 months with diarrhoea in the last two weeks
	Child was given to drink:						Child was given to eat:						
	Much less	Somewhat less	About the same	More	Nothing	Total	Much less	Somewhat less	About the same	More	Nothing	Total	
Total	39.7	26.5	15.4	13.8	4.5	100.0	31.5	36.2	11.4	5.7	15.2	100.0	186
Sex													
Male	44.6	24.8	14.0	11.4	5.2	100.0	35.2	35.6	12.2	6.3	10.7	100.0	94
Female	34.6	28.3	16.9	16.4	3.8	100.0	27.7	36.7	10.6	5.1	19.9	100.0	91
Area													
Urban	38.9	20.8	17.4	17.7	5.3	100.0	35.3	31.1	14.2	5.8	13.6	100.0	104
Rural	40.6	33.8	12.9	9.0	3.6	100.0	26.6	42.6	8.0	5.6	17.2	100.0	82

Table CH.7 and Figure CH.2 show the percentage of children age 0-59 months with diarrhoea in the last two weeks preceding the survey, who received oral rehydration salts (ORS), recommended homemade fluids, and zinc during an episode of diarrhoea. Since children may have been given more than one type of liquid, the percentages do not necessarily add to 100. About 52 percent received fluids from ORS packets or pre-packaged ORS fluids and 47 percent received recommended homemade fluids (cereal gruel – uji; fresh fruit juice; soups; fresh or fermented milk). Approximately 72 percent of children with diarrhoea received one or more of the recommended home treatments (i.e., were treated with ORS or any recommended homemade fluid), while about half of them received zinc. In addition, 27 percent received ORS and zinc.

Table CH.7: Oral rehydration solutions, recommended homemade fluids, and zinc

Percentage of children age 0-59 months with diarrhoea in the last two weeks, and treatment with oral rehydration salts (ORS), recommended homemade fluids, and zinc, Turkana County MICS, 2013/14

	Percentage of children with diarrhoea who received:													Number of children age 0-59 months with diarrhoea in the last two weeks
	Oral rehydration salts (ORS)			Recommended homemade fluids						Zinc				
	Fluid from packet	Pre-packaged fluid	Any ORS	Cereal Gruel(Uji)	Fresh or Fermented Milk	Fresh fruit juices	Soups	Any recommended homemade fluid	ORS or any recommended homemade fluid	Tablet	Syrup	Any zinc	ORS and zinc ¹	
Total	40.0	23.0	52.4	21.8	24.7	9.7	20.2	47.4	72.0	28.9	35.3	50.2	27.3	186
Sex														
Male	42.2	24.2	54.0	20.2	24.9	12.6	19.2	48.2	72.5	28.7	33.9	47.0	28.0	94
Female	37.9	21.8	50.8	23.4	24.5	6.7	21.2	46.6	71.5	29.2	36.8	53.4	26.6	91
Area														
Urban	33.3	19.5	49.0	18.3	18.0	12.5	17.9	45.5	66.3	27.7	34.4	51.9	25.6	104
Rural	48.6	27.4	56.8	26.2	33.3	6.1	23.1	49.8	79.3	30.5	36.5	48.0	29.4	82

¹ MICS indicator 3.11 - Diarrhoea treatment with oral rehydration salts (ORS) and zinc

Figure CH.2: Children under-5 with diarrhoea who received ORS or recommended homemade liquids, Turkana County MICS, 2013/14




Table CH.8 and Figure CH.3 provide the proportion of children age 0-59 months with diarrhoea in the last two weeks preceding the survey who received oral rehydration therapy with continued feeding, and the percentage of children with diarrhoea who received other treatments. Overall, 59 percent of children with diarrhoea received ORS or increased fluids and 74 percent received ORT (ORS or recommended homemade fluids or increased fluids). Combining the information in Table CH.6 with that of Table CH.7 on oral rehydration therapy, it was observed that 43 percent of children received ORT and, at the same time, feeding was continued, as is recommended. Table CH.8 also shows the percentage of children having had diarrhoea in the two weeks preceding the survey who were given various forms of treatment, leaving 13 percent of them without any treatment or drug.

Table CH.8: Oral rehydration therapy with continued feeding and other treatments

Percentage of children age 0-59 months with diarrhoea in the last two weeks who were given oral rehydration therapy with continued feeding and percentage who were given other treatments, Turkana County MICS, 2013/14

	Children with diarrhoea who were given:														Number of children age 0-59 months with diarrhoea in the last two weeks	
	Zinc	ORS or increased fluids	ORT (ORS or recommended homemade fluids or increased fluids)	ORT with continued feeding ¹	Other treatments								Not given any treatment or drug			
					Pill or syrup				Injection							
					Anti-biotic	Anti-motility	Other	Unknown	Anti-biotic	Non-antibiotic	Unknown	Intra-venous		Home remedy, herbal medicine		Other
Total	50.2	59.3	74.1	43.2	10.6	0.0	3.4	1.9	1.3	0.0	0.6	0.0	0.5	0.6	13.4	186
Sex																
Male	47.0	60.2	73.4	44.3	12.6	0.0	2.5	0.9	2.5	0.0	0.0	0.0	1.0	1.2	15.3	94
Female	53.4	58.3	74.8	42.0	8.6	0.0	4.3	2.9	0.0	0.0	1.2	0.0	0.0	0.0	11.5	91
Area																
Urban	51.9	57.1	70.0	36.0	13.2	0.0	4.2	2.0	1.8	0.0	0.0	0.0	0.0	0.0	15.7	104
Rural	48.0	61.9	79.3	52.4	7.4	0.0	2.4	1.7	0.6	0.0	1.3	0.0	1.2	1.4	10.5	82

¹ MICS indicator 3.12 - Diarrhoea treatment with oral rehydration therapy (ORT) and continued feeding

Figure CH.3: Children under-5 with diarrhoea receiving oral rehydration therapy (ORT) and continued feeding, Turkana County MICS, 2013/14




Table CH.9 provides information on the source of ORS and zinc for children who benefitted from these treatments. For both ORS and zinc, the main source was a health facility or a health provider, predominantly from public health facilities.

Table CH.9: Source of ORS and zinc

Percentage of children age 0-59 months with diarrhoea in the last two weeks who were given ORS, and percentage given zinc, by the source of ORS and zinc, Turkana County MICS, 2013/14

Percentage of children who were given as treatment for diarrhoea:		Number of children age 0-59 months with diarrhoea in the last two weeks	Percentage of children for whom the source of ORS was:						Number of children age 0-59 months who were given ORS as treatment for diarrhoea in the last two weeks	Percentage of children for whom the source of zinc was:				Number of children age 0-59 months who were given zinc as treatment for diarrhoea in the last two weeks
			Health facilities or providers							Health facilities or providers				
ORS	zinc		Public	Private	Community health provider ^a	Other source	Missing /DK	A health facility or provider ^b		Public	Private	Other source	A health facility or provider ^b	
52.4	50.2	186	81.3	11.0	0.8	3.3	4.4	92.3	97	80.0	12.6	7.5	92.5	93

^a Community health provider includes both public (*Community health worker and Mobile/Outreach clinic*) and private (*Mobile clinic*) health facilities^b Includes all public and private health facilities and providers

5.3.2 Acute Respiratory Infections

Symptoms of ARI were collected during the Turkana County MICS to capture pneumonia disease, which is a leading cause of death in children under five. Once diagnosed, pneumonia is treated effectively with antibiotics. Studies have shown a limitation in the survey approach of measuring pneumonia because many of the suspected cases identified through surveys are in fact, not true pneumonia.⁵¹ While this limitation does not affect the level and patterns of care-seeking for suspected pneumonia, it limits the validity of the level of treatment of pneumonia with antibiotics, as reported through household surveys. The treatment indicator described in this report must therefore be taken with caution, keeping in mind that the accurate level is likely higher.

Table CH.10 presents the percentage of children with symptoms of ARI in the two weeks preceding the survey for whom care was sought, by source of care and the percentage who received antibiotics. Thirty-nine percent of children with ARI in the two weeks preceding the survey were treated with antibiotics. Advice and treatment was sought from a health facility or a provider (78 percent), mainly the public health facility (66 percent).

Table CH.10 was supposed to further present the results on the use of antibiotics for the treatment of children under-5 years with symptoms of ARI by sex, age, area, wealth index, and to show the point of treatment among children with symptoms of ARI who were treated with antibiotics and the source of antibiotics. However, analysis was not possible due to few cases.

Table CH.10: Care-seeking for and antibiotic treatment of symptoms of acute respiratory infection (ARI)

Percentage of children age 0-59 months with symptoms of ARI in the last two weeks for whom advice or treatment was sought, by source of advice or treatment, and percentage of children with symptoms who were given antibiotics, Turkana County MICS, 2013/14

	Percentage of children with symptoms of ARI for whom:						Percentage of children with symptoms of ARI in the last two weeks who were given antibiotics ²	Number of children age 0-59 months with symptoms of ARI in the last two weeks
	Advice or treatment was sought from:							
	Health facilities or providers			A health facility or provider ¹ ^b	No advice or treatment sought			
	Public	Private	Community health provider ^a			Other source		
Total	65.8	12.9	1.1	0.0	77.8	22.2	39.3	55
¹ MICS indicator 3.13 - Care-seeking for children with acute respiratory infection (ARI) symptoms								
² MICS indicator 3.14 - Antibiotic treatment for children with ARI symptoms								
^a Community health providers includes both public (<i>Community health worker and Mobile/Outreach clinic</i>) and private (<i>Mobile clinic</i>) health facilities								
^b Includes all public and private health facilities and providers, but excludes private pharmacy								
^c Includes all public and private health facilities and providers								

⁵¹Campbell, H. et al. 2013. *Measuring Coverage in MNCH: Challenges in Monitoring the Proportion of Young Children with Pneumonia Who Receive Antibiotic Treatment*. PLoS Med 10(5): e1001421. doi:10.1371/journal.pmed.1001421

Mothers' knowledge of danger signs is an important determinant of care-seeking behaviour. In the MICS, mothers or caretakers were asked to report symptoms that would cause them to take a child under-five years for care immediately at a health facility. Issues related to knowledge of danger signs of pneumonia are presented in Table CH.11. Overall, 60 percent of women know at least one of the two danger signs of pneumonia – fast and/or difficult breathing. The most commonly identified symptom for taking a child to a health facility is when a child develops a fever (77 percent); when a child becomes sicker (72 percent), difficulty in breathing (52 percent), and fast breathing (47 percent).

Table CH.11: Knowledge of the two danger signs of pneumonia

Percentage of women age 15-49 years who are mothers or caretakers of children under age 5 by symptoms that would cause them to take a child under age 5 immediately to a health facility, and percentage of mothers who recognize fast or difficult breathing as signs for seeking care immediately, Turkana County MICS, 2013/14

	Percentage of mothers/caretakers of children age 0-59 months who think that a child should be taken immediately to a health facility if the child:								Mothers/caretakers who recognize at least one of the two danger signs of pneumonia (fast and/or difficult breathing)	Number of women age 15-49 years who are mothers/caretakers of children under age 5
	Is not able to drink or breastfeed	Becomes sicker	Develops a fever	Has fast breathing	Has difficult breathing	Has blood in stool	Is drinking poorly	Has other symptoms		
Total	39.7	71.5	76.5	47.1	51.5	42.6	35.3	24.6	60.2	589
Area										
Urban	42.3	64.9	81.3	44.7	54.2	43.8	37.3	27.7	59.1	309
Rural	36.8	78.7	71.2	49.8	48.6	41.3	33.1	21.2	61.5	280
Education										
None	36.9	72.9	74.4	49.2	49.9	43.3	33.9	22.7	60.5	403
Primary	43.7	70.1	79.8	40.6	56.3	39.2	37.5	30.3	61.0	126
Secondary+	(50.0)	(64.5)	(83.2)	(47.0)	(52.6)	(45.3)	(40.4)	(25.7)	(56.6)	60
Wealth index quintile										
Poorest	31.4	76.7	69.5	46.2	47.3	41.4	27.8	27.4	55.8	112
Second	36.5	73.9	73.4	46.7	51.3	39.0	35.9	15.3	63.6	130
Middle	39.2	75.3	75.8	50.5	48.9	45.2	41.5	28.4	59.9	114
Fourth	46.1	69.7	79.0	48.7	54.5	46.4	32.7	22.7	60.0	138
Richest	45.0	60.1	85.9	42.8	55.8	40.3	39.7	32.2	61.6	96
Ethnicity of household head										
Turkana	37.5	70.6	77.0	46.4	50.9	41.1	35.0	22.7	59.8	500
Other ethnic group	52.2	76.2	73.5	51.1	55.5	51.2	36.9	35.4	62.6	88

() Figures that are based on 25-49 unweighted cases

5.3.3 Solid Fuel Use

More than 3 billion people around the world rely on solid fuels for their basic energy needs, including cooking and heating. Solid fuels include biomass fuels, such as wood, charcoal, crops or other agricultural waste, dung, shrubs and straw, and coal. Cooking and heating with solid fuels leads to high levels of indoor smoke which contains a complex mix of health-damaging pollutants. The main problem with the use of solid fuels is their incomplete combustion, which produces toxic elements such as carbon monoxide, polyaromatic hydrocarbons, and sulphur dioxide (SO₂), among others. Use of solid fuels increases the risks

of incurring acute respiratory illness, pneumonia, chronic obstructive lung disease, cancer, and possibly tuberculosis, asthma, or cataracts, and may contribute to low birth weight of babies born to pregnant women exposed to smoke. The primary indicator for monitoring use of solid fuels is the proportion of the population using solid fuels as the primary source of domestic energy for cooking, shown in Table CH.12.

Almost every household (99 percent) in Turkana County uses solid fuels for cooking, consisting mainly of wood (73 percent). Use of solid fuels is universal in both urban and rural areas, by education of head of household and by household wealth.

Table CH.12: Solid fuel use

Percent distribution of household members according to type of cooking fuel mainly used by the household, and percentage of household members living in households using solid fuels for cooking, Turkana County MICS, 2013/14

	Percentage of household members in households using:										
	Solid fuels						No food cooked in the household	Other	Total	Solid fuels for cooking ¹	Number of household members
	Natural Gas	Biogas	Kerosene	Char-coal	Wood	Straw/Shrubs/Grass					
Total	0.0	0.3	0.2	25.6	73.0	0.7	0.1	0.0	100.0	99.3	6,594
Area											
Urban	0.1	0.6	0.3	43.9	54.5	0.3	0.2	0.1	100.0	98.8	3,598
Rural	0.0	0.0	0.0	3.6	95.3	1.1	0.0	0.0	100.0	100.0	2,996
Education of household head											
None	0.0	0.0	0.0	10.0	89.0	1.0	0.0	0.0	100.0	100.0	4,269
Primary	0.0	0.0	0.0	37.4	62.3	0.2	0.1	0.0	100.0	99.9	1,067
Secondary+	0.3	1.9	0.9	67.9	28.5	0.0	0.4	0.2	100.0	96.4	1,187
Wealth index quintile											
Poorest	0.0	0.0	0.0	0.1	99.9	0.0	0.0	0.0	100.0	100.0	1,319
Second	0.0	0.0	0.0	0.7	97.0	2.3	0.0	0.0	100.0	100.0	1,321
Middle	0.0	0.0	0.0	7.8	91.2	1.0	0.0	0.0	100.0	100.0	1,321
Fourth	0.0	0.0	0.2	34.1	65.3	0.0	0.3	0.0	100.0	99.5	1,314
Richest	0.2	1.7	0.6	85.5	11.6	0.0	0.1	0.1	100.0	97.2	1,319
Ethnicity of household head											
Turkana	0.1	0.3	0.0	22.9	75.9	0.8	0.0	0.0	100.0	99.6	5,483
Other ethnic group	0.0	0.6	1.0	38.6	59.3	0.0	0.4	0.2	100.0	97.8	1,093

¹ MICS indicator 3.15 - Use of solid fuels for cooking

Solid fuel use by place of cooking is depicted in Table CH.13. The presence and extent of indoor pollution are dependent on cooking practices, places used for cooking, as well as types of fuel used. According to the Turkana County MICS, 39 percent of the population living in households using solid fuels for cooking, cook food in a separate room that is used as a kitchen. The percentage that had food cooked in a separate room used as a kitchen within the dwelling unit is slightly higher in urban (41 percent) than in rural areas (37 percent).

Table CH.13: Solid fuel use by place of cooking

Percent distribution of household members in households using solid fuels by place of cooking, Turkana County MICS, 2013/14								
	Place of cooking:						Total	Number of household members in households using solid fuels for cooking
	In the house		In a separate building	Outdoors	Other place			
	In a separate room used as kitchen	Elsewhere in the house						
Total	39.2	13.1	23.3	23.9	0.2	0.2	100.0	6,550
Area								
Urban	40.8	17.2	14.9	26.4	0.4	0.3	100.0	3,554
Rural	37.2	8.4	33.3	21.0	0.1	0.1	100.0	2,996
Education of household head								
None	37.1	11.2	27.6	23.9	0.2	0.1	100.0	4,269
Primary	42.0	17.4	17.1	23.4	0.0	0.0	100.0	1,066
Secondary+	43.8	16.7	14.6	23.2	0.7	1.0	100.0	1,144
Wealth index quintile								
Poorest	35.9	9.6	29.1	25.2	0.0	0.2	100.0	1,319
Second	38.7	7.3	26.5	27.5	0.0	0.0	100.0	1,321
Middle	33.6	18.3	26.1	21.5	0.5	0.0	100.0	1,321
Fourth	39.9	11.6	25.6	22.9	0.0	0.0	100.0	1,307
Richest	48.2	19.0	8.9	22.4	0.6	0.9	100.0	1,281
Ethnicity of household head								
Turkana	40.5	11.5	23.8	23.9	0.1	0.2	100.0	5,462
Other ethnic group	32.4	21.6	21.6	23.7	0.7	0.0	100.0	1,070

5.3.4 Malaria/Fever

Malaria is a major cause of death of children under five years worldwide. In Kenya, malaria accounts for about 31 percent of outpatient consultations and five percent of hospital admissions.⁵² The results of the Kenya Malaria Indicator Survey 2010 showed that children age 5–14 years had the highest prevalence of malaria (13 percent). The prevalence in children below five years increased from four percent in 2007 to eight percent in 2010. Malaria prevalence was also nearly three times as high in rural areas (12 percent) as in urban areas (5 percent).⁵³ Malaria transmission and infection risk in Kenya is determined largely by altitude, rainfall patterns and temperature. Preventive measures and treatment with an effective antimalarial can dramatically reduce malaria mortality rates among children.

In areas where malaria is common, WHO recommends indoor residual spraying (IRS), use of insecticide treated bednets (ITNs) and prompt treatment of cases with recommended anti-malarial drugs.

⁵² President's Malaria Initiative – *Kenya Malaria Operational Plan FY 2014*

⁵³Division of Malaria Control [Ministry of Public Health and Sanitation], Kenya National Bureau of Statistics, and ICF Macro. 2011. *2010 Kenya Malaria Indicator Survey*. Nairobi, Kenya: DOMC, KNBS and ICF Macro.

In 2010 the WHO issued a recommendation for universal use of diagnostic testing to confirm malaria infection and apply appropriate treatment based on the results. According to the guidelines, treatment solely on the basis of clinical suspicion should only be considered when a parasitological diagnosis is not accessible. This recommendation was based on studies that showed substantial reduction in the proportion of fever that are associated with malaria to a low level.⁵⁴ This recommendation implies that the indicator on proportion of children with fever that received antimalarial treatment is no longer an acceptable indicator of the level of treatment of malaria in the population of children under age five. However, as it remains the MDG indicator and for purposes of comparisons, as well as assessment of patterns across socio-demographic characteristics, the indicator remains a standard MICS indicator.

Children with severe malaria symptoms, such as fever and/or convulsions, should be taken to a health facility. Further, children recovering from malaria should be given extra liquids and food, and younger children should continue breastfeeding.

In Kenya, the Division of Malaria Control (DOMC) and President's Malaria Initiative (PMI), have put in place the following interventions for malaria control and case management: indoor residual spraying (IRS); distribution of insecticide-treated nets; intermittent preventive treatment of pregnant women (IPTp); provision of prompt diagnosis and effective treatment at all levels of the health care system; advocacy, communication and social mobilisation through Behaviour Change Communication (BCC); monitoring and evaluation; and health systems strengthening and integration. The Malaria Control Programme is guided by the National Malaria Communication Strategy 2010 – 2013; Kenya National Malaria Strategy 2009 – 2017: Towards a Malaria-free Kenya; and the National Guidelines for the Diagnosis, Treatment and Prevention of Malaria in Kenya 2010.

Insecticide-treated mosquito nets, or ITNs, if used properly, are very effective in offering protection against mosquitos and other insects. The use of ITNs is one of the main health interventions implemented to reduce malaria transmission in Kenya. The questionnaire incorporated questions on the availability and use of bed nets, both at household level and among children under five years of age and pregnant women. In addition, all households in Turkana County were asked whether the interior dwelling walls were sprayed with an insecticide to kill or repel mosquitoes that spread malaria during the 12 months preceding the survey.

In Turkana County, the survey results indicate that 37 percent of households had at least one insecticide treated net (Table CH.14), and 13 percent have at least one ITN for every two household members. Further two percent of households received indoor residual spraying during the last 12 months, and 37 percent have at least one ITN for every two household members and/or received IRS during the last 12 months.

⁵⁴D'Acremont, V et al. 2010. *Reduction in the proportion of fevers associated with Plasmodium falciparum parasitaemia in Africa: a systematic review*. Malaria Journal 9(240).

Table CH.14: Household availability of insecticide treated nets and protection by a vector control method

Percentage of households with at least one mosquito net, one insecticide treated net (ITN), and one long-lasting treated net, percentage of households with at least one mosquito net, one insecticide treated net (ITN) per two people, and one long-lasting treated net, percentage of households with at least one ITN and/or indoor residual spraying (IRS) in the last 12 months, and percentage of households with at least one ITN per two people and/or with indoor residual spraying (IRS) in the last 12 months, Turkana County MICS, 2013/14

	Percentage of households with at least one mosquito net:			Percentage of households with at least one net for every two persons ^a :			Percentage of households with IRS in the past 12 months	Percentage of households with at least one ITN and/or IRS during the last 12 months ³	Percentage of households with at least one ITN for every 2 persons and/or received IRS during the last 12 months ⁴	Number of households
	Any mosquito net	Insecticide treated mosquito net (ITN) ¹	Long-lasting insecticidal treated net (LLIN)	Any mosquito net	Insecticide treated mosquito net (ITN) ²	Long-lasting insecticidal treated net (LLIN)				
Total	40.6	36.6	34.0	14.8	13.4	11.9	2.3	37.3	14.8	1,277
Area										
Urban	53.4	48.2	44.3	21.6	19.4	16.8	3.1	49.0	21.5	684
Rural	25.8	23.3	22.1	7.0	6.4	6.3	1.5	23.9	7.1	593
Education of household head										
None	28.5	25.6	24.6	6.9	6.2	6.1	1.5	26.4	7.5	822
Primary	57.4	52.9	47.7	21.1	20.1	17.0	4.9	53.4	23.1	211
Secondary+	66.4	60.8	54.6	36.1	31.7	27.1	2.5	61.3	32.5	234
Wealth index quintile										
Poorest	18.7	17.4	16.9	4.1	4.1	4.1	0.0	17.4	4.1	256
Second	24.0	21.4	19.2	4.5	3.2	2.7	0.0	21.4	3.2	282
Middle	31.3	27.5	27.0	9.3	8.6	8.6	3.4	28.9	11.5	255
Fourth	50.8	44.9	43.4	14.4	13.0	12.3	6.8	47.2	17.7	239
Richest	82.1	75.7	66.7	43.8	40.0	33.8	1.9	75.7	40.0	245
Ethnicity of household head										
Turkana	35.5	31.2	28.9	8.9	7.7	6.6	0.3	31.2	7.7	1,046
Other ethnic group	63.5	60.9	56.5	41.6	39.1	35.8	11.9	64.8	47.4	226
	¹ MICS indicator 3.16a - Household availability of insecticide-treated nets (ITNs) - One+									
	² MICS indicator 3.16b - Household availability of insecticide-treated nets (ITNs) - One+ per 2 people									
	³ MICS indicator 3.17a - Households covered by vector control - One+ ITNs									
	⁴ MICS indicator 3.17b - Households covered by vector control - One+ ITNs per 2 people									

^a The numerators are based on number of usual (de jure) household members and does not take into account whether household members stayed in the household last night. MICS does not collect information on visitors to the household

Tables CH.15 and CH.16 provide further insight on access to ITNs. Overall, five percent of individuals are estimated to have access to ITNs, i.e. they sleep under an ITN if each ITN in the household is used by two people. Access is slightly higher in urban (8 percent) than in rural (2 percent) areas. Access to an ITN ranges from one percent in the poorest households to 17 percent in the richest households.

Table CH.15: Access to an insecticide treated net (ITN) - number of household members

Percentage of household population with access to an ITN in the household, Turkana County MICS, 2013/14												
	Number of ITNs owned by household:								Total	Percentage with access to an ITN ^a	Number of household members ^b	
	0	1	2	3	4	5	6	8 or more				
Total	63.4	18.4	10.4	5.9	0.8	0.4	0.5	0.2	100.0		5.0	6,594
Number of household members												
1	70.5	25.8	0.9	0.8	0.0	1.9	0.0	0.0	100.0		29.5	91
2	64.3	24.0	10.9	0.9	0.0	0.0	0.0	0.0	100.0		11.7	250
3	60.7	22.5	8.2	8.6	0.0	0.0	0.0	0.0	100.0		16.9	404
4	64.7	19.2	10.4	5.2	0.0	0.0	0.5	0.0	100.0		5.7	820
5	63.2	15.5	12.4	5.4	1.6	1.5	0.4	0.0	100.0		8.9	966
6	57.4	19.8	15.0	7.8	0.0	0.0	0.0	0.0	100.0		0.0	1,030
7	67.6	15.8	10.4	5.7	0.6	0.0	0.0	0.0	100.0		0.6	973
8 or more	62.4	12.1	9.9	9.2	2.7	0.5	2.4	0.9	100.0		3.3	2,061


^a Percentage of household population who could sleep under an ITN if each ITN in the household were used by up to two people

^b The denominator is number of usual (de jure) household members and does not take into account whether household members stayed in the household last night. MICS does not collect information on visitors to the household

Table CH.16: Access to an insecticide treated net (ITN) - background characteristics

Percentage of household population with access to an ITN in the household, Turkana County MICS, 2013/14		
	Percentage with access to an ITN ^a	Number of household members ^b
Total	5.0	6,594
Area		
Urban	7.5	3,598
Rural	2.0	2,996
Wealth index quintile		
Poorest	0.9	1,319
Second	0.6	1,321
Middle	2.3	1,321
Fourth	4.0	1,314
Richest	17.2	1,319
Ethnicity of household head		
Turkana	3.1	5,483
Other ethnic group	14.2	1,093
^a Percentage of household population who could sleep under an ITN if each ITN in the household were used by up to two people		
^b The denominator is number of usual (de jure) household members and does not take into account whether household members stayed in the household last night. MICS does not collect information on visitors to the household		

Figure CH.4: Percentage of household population with access to an ITN in the household, Turkana County MICS, 2013/14



Overall, 76 percent of ITNs were used during the night preceding the survey. The percentage of ITNs used by anyone the night preceding the survey is higher in urban areas (81 percent) than in rural areas (63 percent). The proportion is also higher among the wealthiest compared to the poorest households.

Table CH.17: Use of ITNs

Percentage of insecticide treated nets (ITNs) that were used by anyone last night, Turkana County MICS, 2013/14

	Percentage of ITNs used last night	Number of ITNs
Total	76.2	856
Area		
Urban	80.5	644
Rural	63.0	212
Wealth index quintile		
Poorest	47.8	59
Second	57.8	82
Middle	72.0	103
Fourth	82.6	189
Richest	81.7	424
Ethnicity of household head		
Turkana	71.6	552
Other ethnic group	84.3	298

As for children under the age of five years, who constitute an important vulnerable group, 25 percent slept under an ITN the night preceding the survey. This figure increases to 66 percent considering only children living in a household with at least one ITN. Disparities in ITN use among children under-5 years are noted by urban/rural areas. The percentage of children who slept under an ITN the night before the survey was higher in urban areas than in rural areas (37 compared to 12 percent). Similarly, in households with at least one ITN, a higher proportion (75 percent) slept under an ITN in urban areas, compared to 46 percent in rural areas. Some differences are also apparent in regard to the education level of the mother, and household wealth, with the proportion of children sleeping under an ITN being higher among children of mothers with secondary or higher education (53 percent) compared to children of mothers with no education (17 percent), and among children in the richer households.

Table CH.18: Children sleeping under mosquito nets

Percentage of children age 0-59 months who slept under a mosquito net last night, by type of net, Turkana County MICS, 2013/14

	Percentage of children age 0-59 who spent last night in the interviewed households	Number of children age 0-59 months	Percentage of children under age five who the previous night slept under:				Number of children age 0-59 months who spent last night in the interviewed households	Percentage of children who slept under an ITN last night in households with at least one ITN	Number of children age 0-59 living in households with at least one ITN
			Any mosquito net	An insecticide treated net (ITN) ¹	A Long-lasting insecticidal treated net (LLIN)	An ITN or in a dwelling sprayed with IRS in the past 12 months			
Total	98.5	1,067	27.2	24.7	23.3	25.4	1,051	65.7	395
Sex									
Male	98.0	537	25.8	23.9	22.7	24.6	526	60.3	208
Female	99.0	530	28.6	25.5	23.9	26.3	525	71.7	187
Area									
Urban	98.5	546	40.9	37.1	35.0	38.0	538	75.0	266
Rural	98.5	521	12.9	11.6	11.0	12.3	513	46.4	129
Age									
0-11 months	99.7	227	28.6	27.1	25.1	27.6	226	73.7	83
12-23 months	97.3	196	33.9	29.4	27.4	30.3	190	71.0	79
24-35 months	99.6	205	24.2	24.0	23.1	25.0	204	67.4	73
36-47 months	99.0	222	27.0	23.6	22.6	24.4	220	58.3	89
48-59 months	96.7	217	22.9	19.7	18.6	20.2	210	58.0	71
Mother's education									
None	99.0	758	18.1	16.8	16.3	17.7	751	54.4	231
Primary	98.4	207	48.2	41.3	38.9	41.3	204	79.5	106
Secondary+	94.3	99	55.2	52.9	46.1	54.0	93	(85.6)	58
Wealth index quintile									
Poorest	98.3	216	7.5	6.3	5.8	6.3	213	30.9	44
Second	98.5	244	11.2	11.0	11.0	11.0	241	50.0	53
Middle	98.8	217	23.5	21.0	20.0	23.2	214	67.0	67
Fourth	99.3	227	39.8	34.4	34.4	35.9	225	76.6	101
Richest	97.0	163	65.4	61.4	54.3	61.4	158	74.6	130
Ethnicity of household head									
Turkana	98.3	898	22.6	20.1	18.7	20.1	883	59.2	300

Other ethnic group	99.3	167	52.2	49.5	48.1	54.2	166	86.2	95
¹ MICS indicator 3.18; MDG indicator 6.7 - Children under age 5 sleeping under insecticide-treated nets (ITNs)									

Table CH.19 gives further insight into the use of mosquito nets by household members of any age, 20 percent of whom slept under an ITN the night prior to the survey. This figure increases to 54 percent considering only household members living in a household with at least one ITN. Overall, 21 percent of household members slept under an ITN the previous night or in a dwelling which had IRS in the past 12 months. The percentage of household members who slept under an ITN the night prior to the survey is 29 percent in urban areas and nine percent in rural areas. Variations are also noted by education level of the household head, with those with no education at 12 percent, those with primary education at 33 percent, while those with secondary education or higher is at 39 percent.

Table CH.19: Use of mosquito nets by the household population							
Percentage of household members who slept under a mosquito net last night, by type of net, Turkana County MICS, 2013/14							
	Percentage of household members who the previous night slept under:				Number of household members who spent the previous night in the interviewed households	Percentage of household members who slept under an ITN last night in households with at least one ITN	Number of household members in households with at least one ITN
	Any mosquito net	An insecticide treated net (ITN) ¹	A Long-lasting insecticidal treated net (LLIN)	An ITN or in a dwelling sprayed with IRS in the past 12 months			
Total	21.7	19.8	18.1	21.0	6,065	53.7	2,237
Sex							
Male	19.3	17.6	16.0	18.7	2,958	47.3	1,101
Female	24.0	21.9	20.1	23.2	3,107	59.9	1,135
Area							
Urban	31.6	28.6	25.9	30.1	3,287	59.4	1,583
Rural	10.0	9.4	8.9	10.2	2,778	39.9	653
Age							
0-4 ^a	27.1	24.5	23.1	25.3	1,051	65.2	396
5-14	17.7	15.9	14.1	16.9	2,012	46.2	692
15-34	23.1	21.4	19.0	23.3	1,826	51.7	757
35-49	27.7	25.4	24.2	26.1	602	65.3	234
50+	15.4	13.9	13.4	14.5	574	50.5	158
Education of household head							
None	12.5	11.5	11.1	12.6	3,970	42.7	1,064
Primary	35.7	33.1	28.6	34.8	989	63.0	520
Secondary+	42.9	38.9	34.3	39.5	1,043	63.9	634
Wealth index quintile							
Poorest	4.9	4.5	4.4	4.5	1,229	27.0	207
Second	9.3	8.9	8.0	8.9	1,226	38.5	282
Middle	13.5	12.2	11.6	15.5	1,212	44.7	330
Fourth	27.7	24.2	23.5	26.7	1,207	57.6	508
Richest	54.3	50.1	43.7	50.1	1,191	65.6	909
Ethnicity of household head							
Turkana	16.9	15.1	13.5	15.1	5,033	45.3	1,679
Other ethnic group	45.4	42.5	40.4	49.5	1,014	78.6	548
¹ MICS indicator 3.19 - Population that slept under an ITN							
^a The results of the age group 0-4 years do not match those in Table CH.18, which is based on completed under-5 interviews only. The two tables are computed with different sample weights							

Table CH.20 provides information on care-seeking behaviour during an episode of fever in the last two weeks preceding the survey. As shown in Table CH.20, advice was sought from a health facility or a qualified health care provider for 67 percent of children with fever; these services were provided mainly by the public health facility (49 percent). However, no advice or treatment was sought in 27 percent of the cases.

Table CH.20: Care-seeking during fever							
Percentage of children age 0-59 months with fever in the last two weeks for whom advice or treatment was sought, by source of advice or treatment, Turkana County MICS, 2013/14							
	Percentage of children for whom:						Number of children with fever in last two weeks
	Advice or treatment was sought from:						
	Health facilities or providers			Other source	A health facility or provider ^{1, b}	No advice or treatment sought	
Public	Private	Community health provider ^a					
Total	48.7	19.0	1.8	5.2	66.9	27.1	198
Sex							
Male	54.0	20.3	3.2	6.0	71.6	19.7	94
Female	43.9	17.9	0.6	4.5	62.5	33.8	103
Area							
Urban	46.2	24.7	0.9	5.3	67.5	23.8	114
Rural	52.0	11.4	3.1	5.1	66.0	31.5	84
¹ MICS indicator 3.20 - Care-seeking for fever							
^a Community health providers include both public (<i>Community health worker</i> and <i>Mobile/Outreach clinic</i>) and private (<i>Mobile clinic</i>) health facilities							
^b Includes all public and private health facilities and providers as well as shops							

Mothers were asked to report all of the medicines given to a child to treat the fever, including both medicines given at home and medicines given or prescribed at a health facility. Artemisinin-based Combination therapy (ACT) is the first line antimalarial recommended by the World Health Organization and used in the country. In addition, confirmation of malaria is done on all fever cases through a malaria test.

Table CH.21 presents the results of children age 0-59 months who had a fever in the last two weeks preceding the survey, by type of medicine given for the illness. About nine percent of children with fever during this period were treated with an artemisinin-based combination therapy (ACT) and 10 percent received other antimalarials including Fansidar, amodiaquine and quinine. The proportion of children who received ACT in urban areas is 13 percent while it is three percent in rural areas.

Table CH.21: Treatment of children with fever

Percentage of children age 0-59 months who had a fever in the last two weeks, by type of medicine given for the illness, Turkana County MICS, 2013/14

	Children with a fever in the last two weeks who were given:													Number of children with fever in last two weeks
	Anti-malarials						Other medications							
	SP/ Fansidar	Chloroquine	Amodia- quine	Quinine	Artemisinin- based Combination Therapy (ACT)	Other anti- malarial	Antibiotic pill or syrup	Antibiotic injection	Paracetamol/ Panadol/ Acetaminophen	Aspirin	Ibuprofen	Other	Missing/DK	
Total	3.5	0.0	2.0	1.5	8.8	3.1	20.7	7.1	44.2	0.8	3.9	14.0	4.0	198
Sex														
Male	2.6	0.0	4.2	2.6	9.8	3.9	25.9	5.3	42.4	0.6	5.1	9.8	4.3	94
Female	4.4	0.0	0.0	0.4	7.8	2.4	15.9	8.9	45.8	0.9	2.8	17.8	3.8	103
Area														
Urban	4.1	0.0	3.5	0.0	12.7	4.6	24.0	5.9	47.4	0.0	4.1	17.5	3.1	114
Rural	2.7	0.0	0.0	3.4	3.4	1.1	16.2	8.8	39.9	1.8	3.6	9.3	5.3	84

Overall, 45 percent of children with a fever in the previous two weeks preceding the survey had blood taken from a finger or heel for testing. Eighteen percent of children who had fever in the two weeks preceding the survey were treated with any antimalarial drugs. Of these, half of them were treated with ACT.

Table CH.22: Diagnostics and anti-malarial treatment of children

Percentage of children age 0-59 months who had a fever in the last two weeks who had a finger or heel stick for malaria testing, who were given Artemisinin-combination Treatment (ACT) and any anti-malarial drugs, and percentage who were given ACT among those who were given anti-malarial drugs, Turkana County MICS, 2013/14								
	Percentage of children who:					Number of children age 0-59 months with fever in the last two weeks	Treatment with Artemisinin-based Combination Therapy (ACT) among children who received anti-malarial treatment ³	Number of children age 0-59 months with fever in the last two weeks who were given any antimalarial drugs
	Had blood taken from a finger or heel for testing ¹	Were given:						
	Artemisinin-combination Treatment (ACT)	ACT the same or next day	Any antimalarial drugs ²	Any antimalarial drugs same or next day				
Total	45.4	8.8	8.1	17.5	14.8	198	(50.3)	35
Sex								
Male	48.0	12.7	11.6	22.5	19.4	114	(*)	26
Female	42.0	3.4	3.4	10.7	8.6	84	(*)	9
Area								
Urban	48.0	12.7	11.6	22.5	19.4	114	(*)	26
Rural	42.0	3.4	3.4	10.7	8.6	84	(*)	9
¹ MICS indicator 3.21 - Malaria diagnostics usage								
² MICS indicator 3.22; MDG indicator 6.8 - Anti-malarial treatment of children under age 5								
³ MICS indicator 3.23 - Treatment with Artemisinin-based Combination Therapy (ACT) among children who received anti-malarial treatment								
() Figures that are based on 25-49 unweighted cases								
(*) Figures that are based on fewer than 25 unweighted cases								

Table CH.23 presents the source of antimalarial for children under-5 years who were treated with an antimalarial. Eighteen percent of children were given antimalarial.

Table CH.23: Source of anti-malarial

Percentage of children age 0-59 months with fever in the last two weeks who were given anti-malarial by the source of anti-malarial, Turkana County MICS, 2013/14								
	Percentage of children who were given anti-malarial	Number of children age 0-59 months with fever in the last two weeks	Percentage of children for whom the source of anti-malarial was:					Number of children age 0-59 months who were given anti-malarial as treatment for fever in the last two weeks
			Health facilities or providers			Other source	A health facility or provider ^b	
			Public	Private	Community health provider ^a			
Total	17.5	198	(48.9)	(43.3)	(5.0)	(7.8)	(97.7)	35
^a Community health providers include both public (<i>Community health worker and Mobile/Outreach clinic</i>) and private (<i>Mobile clinic</i>) health facilities								
^b Includes all public and private health facilities and providers as well as shops								
() Figures that are based on 25-49 unweighted cases								

Pregnant women living in places where malaria is highly prevalent are highly vulnerable to malaria. Once infected, pregnant women risk anaemia, premature delivery and stillbirth. Their babies are at

increased risk of low birth weight, which carries an increased risk of dying in infancy.⁵⁵ For this reason, steps are taken to protect pregnant women by distributing insecticide-treated mosquito nets and mobilizing for their consistent use; and treatment during antenatal check-ups with drugs that prevent malaria infection (Intermittent preventive treatment/IPT). WHO recommends that in areas of moderate-to-high malaria transmission, all pregnant women be provided an intermittent preventive treatment with Sulfadoxine-Pyrimethamine (SP) at every scheduled ANC visit.

During the Turkana County MICS, women were asked of the medicines they had received to prevent malaria in their last pregnancy during the 2 years preceding the survey. Women were considered to have received intermittent preventive therapy if they had received at least 3 doses of SP/Fansidar during the pregnancy, at least one of which was taken during ANC.

Table CH.24 presents the proportion of pregnant women who slept under a mosquito net during the previous night.⁵⁶ Thirty-seven percent of pregnant women slept under any mosquito net the night prior to the survey and 34 percent slept under an insecticide-treated net.

Table CH.24: Pregnant women sleeping under mosquito nets									
Percentage of pregnant women age 15-49 years who slept under a mosquito net last night, by type of net, Turkana County MICS, 2013/14									
Percentage of pregnant women who spent last night in the interviewed households	Number of pregnant women age 15-49 years	Percentage of pregnant women age 15-49 years who the previous night slept under:					Number of pregnant women who spent last night in the interviewed households	Percentage of pregnant women who slept under an ITN last night in households with at least one ITN	Number of pregnant women age 15-49 years living in households with at least one ITN
		Any mosquito net	An insecticide treated net (ITN) ¹	A Long-lasting insecticidal treated net (LLIN)	An ITN or in a dwelling sprayed with IRS in the past 12 months				
Total	100.0	67	37.3	33.5	31.7	36.0	67	(76.1)	29
¹ MICS indicator 3.24 - Pregnant women who slept under an insecticide treated net (ITN)									
() Figures that are based on 25-49 unweighted cases									

Intermittent preventive treatment for malaria in pregnant women who gave birth in the two years preceding the survey is presented in Table CH.25. Overall, 93 percent of women age 15-49 years who had a live birth during the two years preceding the survey received antenatal care. Ninety-five percent of women received any medicine to prevent malaria at any ANC visit during the pregnancy. About 30 percent of the women received SP/Fansidar at least three or more times during an ANC visit. The proportion in urban areas that received SP/Fansidar three or more times during ANC is 26 percent and 35 percent in rural areas.

⁵⁵Shulman, CE and Dorman, EK. 2003. *Importance and prevention of malaria in pregnancy*. Trans R Soc Trop Med Hyg 97(1): 30–55.

⁵⁶All background variables were removed from the table due to small number of cases reported.

Table CH.25: Intermittent preventive treatment for malaria

Percentage of women age 15-49 years who had a live birth during the two years preceding the survey and who received intermittent preventive treatment (IPT) for malaria during pregnancy at any antenatal care visit, Turkana County MICS, 2013/14

	Percentage of women who received antenatal care (ANC)	Number of women with a live birth in the last two years	Percentage of pregnant women:				Number of women with a live birth in the last two years and who received antenatal care	
			Who took any medicine to prevent malaria at any ANC visit during pregnancy	who took SP/Fansidar at least once during an ANC visit and in total took:				
				At least once	Two or more times	Three or more times ¹	Four or more times	
Total	93.4	387	95.4	91.1	58.7	30.1	7.3	362
Area								
Urban	98.3	199	96.5	92.8	54.0	26.2	4.2	196
Rural	88.2	188	94.3	89.2	64.3	34.8	10.8	166
Education								
None	90.5	268	95.1	91.5	62.1	31.9	8.8	242
Primary	100.0	82	94.4	88.0	46.6	19.3	2.8	82
Secondary+	(100.0)	38	(100.0)	(95.2)	(63.0)	(42.3)	(6.9)	38
Wealth index quintile								
Poorest	84.1	69	94.7	91.3	65.0	32.0	10.2	58
Second	91.2	86	95.2	90.2	62.3	35.1	7.8	79
Middle	96.1	84	93.5	87.8	52.5	34.2	10.3	81
Fourth	97.6	85	96.6	91.5	62.3	22.0	3.7	83
Richest	97.5	63	97.5	95.9	51.5	27.8	4.5	62
Ethnicity of household head								
Turkana	92.5	327	95.3	91.0	60.6	30.8	7.4	303
Other ethnic group	98.3	59	(96.3)	(91.6)	(48.2)	(25.8)	(6.4)	58
¹ MICS indicator 3.25 - Intermittent preventive treatment for malaria								
() Figures that are based on 25-49 unweighted cases								

6. Water and Sanitation

Safe drinking water is a basic necessity for good health. Unsafe drinking water can be a significant determinant of diseases such as cholera, typhoid, and schistosomiasis. Drinking water can also be contaminated with chemical and physical contaminants with harmful effects on human health. In addition to preventing disease, improved access to drinking water may be particularly important for women and children, especially in rural areas, who bear the primary responsibility for carrying water, often for long distances.⁵⁷

Inadequate disposal of human excreta and personal hygiene is associated with a range of diseases including diarrhoeal diseases and polio and is an important determinant for stunting. Improved sanitation can reduce diarrheal disease by more than a third,⁵⁸ and can significantly lessen the adverse health impacts of other disorders responsible for death and disease among millions of children in developing countries.

The goal of MDG 7 is to reduce by half, between 1990 and 2015, the proportion of people without sustainable access to safe drinking water and basic sanitation.

For more details on water and sanitation and to access some reference documents, please visit data.unicef.org⁵⁹ or the website of the WHO/UNICEF Joint Monitoring Programme for Water Supply and Sanitation.⁶⁰

The Kenya National Water Policy of 2012 was developed in response to the mandate, vision and mission of the ministry responsible for water affairs in the country. The policy takes into account requirements of the Constitution of Kenya 2010;⁶¹ the Kenya Vision 2030; the Millennium Development Goals (MDGs), and other national policies and strategies.⁶²

6.1 Use of Improved Water Sources

The distribution of the population by main source of drinking water is shown in Table WS.1. The population using improved sources of drinking water are those using any of the following types of supply: piped water (into dwelling, compound, yard or plot, to neighbour, public tap/standpipe), tube well/borehole, protected well, protected spring, and rainwater collection. Bottled water is considered as an improved water source only if the household is using an improved water source for handwashing and cooking.

In Turkana, 72 percent of the population uses an improved source of drinking water – 86 percent in urban areas and 55 percent in rural areas (Table WS.1). There is a positive correlation between the proportion of the population using an improved source of drinking water and the education level of

⁵⁷WHO/UNICEF. 2012. *Progress on Drinking water and Sanitation: 2012 update*.

⁵⁸Cairncross, S et al. 2010. *Water, sanitation and hygiene for the prevention of diarrhoea*. International Journal of Epidemiology 39: i193-i205

⁵⁹<http://data.unicef.org/water-sanitation>

⁶⁰<http://www.wssinfo.org>

⁶¹Constitution of Kenya of 2010 [Promulgated on 25th August 2010]

⁶²Ministry of Water and Irrigation. 2012. *The National Water Policy 2012*

the head of household. The proportion increases from 63 percent for heads of households with no education, to 82 percent for those with primary education, and further to 93 percent for those with secondary and higher education. The improved drinking water sources for the population varied strongly by urban/rural area. In urban areas, 35 percent of the population used drinking water that was from a public tap/standpipe, 35 percent used piped water into their dwelling or into their yard or plot and 12 percent used piped water to neighbour. In rural areas the improved drinking water sources mainly used were a public tap/standpipe (44 percent).

Table WS.1: Use of improved water sources

Percent distribution of household population according to main source of drinking water and percentage of household population using improved drinking water sources, Turkana County MICS, 2013/14


	Main source of drinking water										Total	Percentage using improved sources of drinking water ¹	Number of household members
	Improved sources					Unimproved sources							
	Piped water			Public tap/stand-pipe	Tube-well/bore-hole	Protected well	Unprotected well	Unprotected spring	Surface water	Other			
	Into dwelling	Into yard/plot	To neighbour										
Total	2.6	17.6	8.2	39.1	3.4	1.0	5.5	0.6	21.7	0.4	100.0	71.9	6,594
Area													
Urban	4.7	30.4	12.3	34.6	2.8	0.7	2.5	0.0	11.2	0.8	100.0	85.5	3,598
Rural	0.0	2.2	3.2	44.4	4.3	1.3	9.1	1.2	34.2	0.0	100.0	55.4	2,996
Education of household head													
None	0.2	7.0	7.7	42.6	4.2	1.4	7.3	0.6	28.7	0.4	100.0	63.1	4,269
Primary	3.6	21.9	12.5	40.8	2.1	0.6	3.7	1.0	13.5	0.4	100.0	81.5	1,067
Secondary+	10.3	48.5	6.4	25.6	2.3	0.0	1.0	0.0	5.1	0.7	100.0	93.1	1,187
Wealth index quintile													
Poorest	0.0	0.0	0.0	24.6	2.9	1.8	15.0	0.4	55.3	0.0	100.0	29.3	1,319
Second	0.0	0.0	1.1	53.2	5.4	1.1	7.0	2.1	28.8	1.2	100.0	60.8	1,321
Middle	0.0	1.3	15.3	55.8	3.9	2.2	4.1	0.2	16.8	0.3	100.0	78.5	1,321
Fourth	4.9	17.1	12.7	54.2	3.4	0.0	1.3	0.0	6.4	0.0	100.0	92.2	1,314
Richest	8.0	69.4	11.8	7.6	1.6	0.0	0.0	0.0	0.9	0.6	100.0	98.5	1,319
Ethnicity of household head													
Turkana	2.9	16.1	9.5	33.3	3.7	1.2	6.5	0.7	25.8	0.4	100.0	66.7	5,483
Other ethnic group	0.5	25.0	1.7	68.1	2.4	0.0	0.6	0.0	0.9	0.8	100.0	97.7	1,093

¹ MICS indicator 4.1; MDG indicator 7.8 - Use of improved drinking water sources

^aHouseholds using bottled water as the main source of drinking water are classified into improved or unimproved drinking water users according to the water source used for other purposes such as cooking and handwashing. There were no cases for bottled water as a source under 'unimproved sources'.

The sources of drinking water used in Turkana County are depicted in Figure WS.1. The majority of the population (39 percent) used a public tap/stand-pipe, followed by surface water (22 percent) and piped water into dwelling, yard or plot (20 percent).

Figure WS.1: Percent distribution of household members by source of drinking water, Turkana County MICS, 2013/14



Use of household water treatment is presented in Table WS.2. Households were asked about ways they may be treating water at home to make it safer to drink. Boiling water, adding bleach or chlorine, using a water filter, and using solar disinfection are considered as effective treatment of drinking water. The table shows water treatment by all household members and the percentage of those living in households using unimproved water sources but using appropriate water treatment methods. Ninety-six percent do not use any method to treat water in order to make it safer to drink and only three percent of household members in households using unimproved drinking water sources are using an appropriate water treatment method, nine percent in urban areas and one percent rural areas. There are noted variations by education level of head of household as it is two percent for those with no education, three percent for those with primary education and 29 percent for those with secondary/higher education.

Table WS.2: Household water treatment

Percentage of household population by drinking water treatment method used in the household, and for household members living in households where an unimproved drinking water source is used, the percentage who are using an appropriate treatment method, Turkana County MICS, 2013/14

	Water treatment method used in the household					Number of household members	Percentage of household members in households using unimproved drinking water sources and using an appropriate water treatment method ¹	Number of household members in households using unimproved drinking water sources
	None	Boil	Add bleach/chlorine	Let it stand and settle	Other			
Total	95.7	2.3	1.7	0.2	0.3	6,594	3.3	1,856
Area								
Urban	93.9	3.1	2.7	0.0	0.6	3,598	8.9	521
Rural	97.7	1.3	0.5	0.5	0.0	2,996	1.1	1,336
Main source of drinking water								
Improved	95.5	2.5	1.8	0.0	0.4	4,738	na	na
Unimproved	96.0	1.8	1.5	0.7	0.0	1,856	3.3	1,856
Education of household head								
None	97.7	1.5	0.5	0.3	0.0	4,269	1.9	1,577
Primary	94.6	2.9	1.8	0.0	0.7	1,067	3.2	198
Secondary+	89.0	4.6	6.3	0.0	1.1	1,187	29.2	82
Wealth index quintile								
Poorest	98.6	0.8	0.0	0.6	0.0	1,319	0.7	933
Second	97.7	1.5	0.3	0.5	0.0	1,321	3.0	517
Middle	97.2	1.9	0.9	0.0	0.0	1,321	5.3	284
Fourth	95.1	3.2	1.7	0.0	0.0	1,314	14.4	102
Richest	89.6	4.0	5.7	0.0	1.6	1,319	(46.0)	20
Ethnicity of household head								
Turkana	95.9	2.4	1.5	0.3	0.1	5,483	3.3	1,828
Other ethnic group	94.8	1.4	2.7	0.0	1.2	1,093	(*)	25
¹ MICS indicator 4.2 - Water treatment								
na: not applicable								

The amount of time it takes to obtain water is presented in Table WS.3 and the person who usually collects the water in Table WS.4. Note that for Table WS.3, household members using water on premises are also shown in this table and for others, the results refer to one roundtrip from home to drinking water source. Information on the number of trips made in one day was not collected.

Table WS.3 shows that 29 percent of the household population have drinking water source on premises. The availability of water on premises is associated with greater use, better family hygiene and better health outcomes. For a water collection round trip of 30 minutes or more it has been observed that households carry progressively less water and are likely to compromise on the minimal basic drinking water needs of the household.⁶³ For 22 percent of the household population in the survey, it takes 30 minutes or more to get to the water source and bring water from an improved water source. About 23 percent of those using an unimproved drinking water source spend 30 minutes or

⁶³Cairncross, S and Cliff, JL. 1987. *Water use and Health in Mueda, Mozambique*. Transactions of the Royal Society of Tropical Medicine and Hygiene 81: 51-4.

more per round trip. In rural areas a higher percentage of household members live in households that spend more time in collecting water compared to those in urban areas especially for drinking water from an unimproved water source.

Table WS.3: Time to source of drinking water

Percent distribution of household population according to time to go to source of drinking water, get water and return, for users of improved and unimproved drinking water sources, Turkana County MICS, 2013/14

	Time to source of drinking water								Number of household members
	Users of improved drinking water sources				Users of unimproved drinking water sources				
	Water on premises	Less than 30 minutes	30 minutes or more	Missing/DK	Less than 30 minutes	30 minutes or more	Missing/DK	Total	
Total	28.6	20.3	22.3	0.7	4.8	23.0	0.3	100.0	6,594
Area									
Urban	47.4	18.7	18.4	1.0	3.3	10.7	0.4	100.0	3,598
Rural	5.9	22.3	26.9	0.3	6.6	37.7	0.2	100.0	2,996
Education of household head									
None	15.0	22.2	25.0	0.8	6.0	30.7	0.2	100.0	4,269
Primary	38.0	22.4	20.7	0.4	2.1	16.0	0.4	100.0	1,067
Secondary+	66.4	12.1	14.3	0.4	3.3	2.9	0.7	100.0	1,187
Wealth index quintile									
Poorest	0.0	11.3	17.5	0.4	8.1	62.4	0.2	100.0	1,319
Second	1.1	26.1	32.7	0.9	9.4	29.3	0.5	100.0	1,321
Middle	16.9	32.5	28.0	1.1	3.9	17.3	0.3	100.0	1,321
Fourth	35.4	27.9	28.1	0.8	2.1	5.7	0.0	100.0	1,314
Richest	89.6	3.9	5.0	0.0	0.7	0.2	0.6	100.0	1,319
Ethnicity of household head									
Turkana	28.7	16.4	20.9	0.6	5.8	27.3	0.2	100.0	5,483
Other ethnic group	27.4	40.5	28.9	0.9	0.1	1.4	0.8	100.0	1,093

Table WS.4 shows that for the majority of households (80 percent), an adult female usually collects drinking water when the source is not on the premises. Adult men collect water in only nine percent of cases, while for the rest of the households, female (7 percent), or male children (3 percent) under 15 years collected water.

Table WS.4: Person collecting water

Percentage of households without drinking water on premises, and percent distribution of households without drinking water on premises according to the person usually collecting drinking water used in the household, Turkana County MICS, 2013/14

	Percentage of households without drinking water on premises	Number of households	Person usually collecting drinking water						Total	Number of households without drinking water on premises
			Adult woman	Adult man	Female child under age 15	Male child under age 15	Missing/DK			
Total	71.3	1,277	80.0	8.6	7.1	3.0	1.2	100.0	911	
Area										
Urban	52.7	684	77.1	10.7	5.7	4.8	1.7	100.0	360	
Rural	92.8	593	82.0	7.2	8.0	1.8	0.9	100.0	550	
Education of household head										
None	85.5	822	81.1	6.4	8.4	3.2	0.9	100.0	703	
Primary	61.4	211	79.9	13.2	0.9	3.3	2.8	100.0	129	
Secondary+	32.1	234	71.6	20.6	4.6	1.6	1.6	100.0	75	
Wealth index quintile										
Poorest	100.0	256	84.3	5.6	7.3	2.4	0.5	100.0	256	
Second	99.0	282	79.7	6.0	10.0	3.5	0.9	100.0	279	
Middle	80.2	255	81.2	10.4	4.6	1.7	2.1	100.0	204	
Fourth	60.6	239	71.2	14.7	6.2	5.8	2.2	100.0	145	
Richest	10.8	245	(*)	(*)	(*)	(*)	(*)	100.0	26	
Ethnicity of household head										
Turkana	73.7	1,046	81.7	6.6	7.5	3.0	1.3	100.0	771	
Other ethnic group	60.9	226	70.4	20.0	5.2	3.4	0.9	100.0	138	

(*) Figures that are based on fewer than 25 unweighted cases

6.2 Use of Improved Sanitation

Inadequate disposal of human excreta and personal hygiene are associated with a range of diseases including diarrhoeal diseases and polio and are important determinants of stunting. Improved sanitation can reduce diarrhoeal disease by more than a third⁶⁴, and can substantially lessen the adverse health impacts of other disorders among millions of children in many countries.

An improved sanitation facility is defined as one that hygienically separates human excreta from human contact. Improved sanitation facilities for excreta disposal include flush or pour flush to a piped sewer system, septic tank, or pit latrine; ventilated improved pit latrine, pit latrine with slab, and use of a composting toilet. The data on the use of improved sanitation facilities in Turkana County are provided in Table WS.5.

A quarter of the population were living in households using improved sanitation facilities (Table WS.5). The percentage is 42 percent in urban areas and five percent in rural areas. The table indicates that use of improved sanitation facilities increases with the education level of the head of the household. The household population in urban areas primarily use pit latrines with slabs. Sixty-three percent of

⁶⁴Cairncross, S. 2010. *Water, sanitation and hygiene for the prevention of diarrhoea*. Int. J. Epidemiology 39: i193-i205.

household population practises open defecation (no facility, bush/field) – 90 percent in rural areas and 41 percent in urban areas.

Table WS.5: Types of sanitation facilities

Percent distribution of household population according to type of toilet facility used by the household, Turkana County MICS, 2013/14

	Type of toilet facility used by household												Total	Number of household members
	Improved sanitation facility						Unimproved sanitation facility					Open defecation (no facility, bush, field)		
	Flush/Pour flush to:			Ventilated improved pit latrine	Pit latrine with slab	Composting toilet	Flush/Pour flush to somewhere else	Pit latrine without slab/open pit	Hanging toilet/latrine	Other	Missing/DK			
Piped sewer system	Septic tank	Pit latrine												
Total	0.1	0.2	2.2	4.9	15.2	2.7	0.0	11.1	0.1	0.2	0.3	63.0	100.0	6,594
Area														
Urban	0.2	0.0	4.1	8.0	25.1	4.7	0.0	16.5	0.2	0.3	0.5	40.5	100.0	3,598
Rural	0.0	0.4	0.0	1.1	3.3	0.3	0.0	4.7	0.0	0.0	0.1	90.1	100.0	2,996
Education of household head														
None	0.0	0.1	0.5	2.4	7.0	0.9	0.0	8.8	0.0	0.2	0.1	80.1	100.0	4,269
Primary	0.0	0.2	4.9	5.1	29.6	2.3	0.1	11.1	0.0	0.5	0.0	46.2	100.0	1,067
Secondary+	0.7	0.5	6.3	13.8	30.9	9.3	0.0	19.5	0.6	0.0	1.4	17.1	100.0	1,187
Wealth index quintile														
Poorest	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	100.0	100.0	1,319
Second	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.4	0.0	0.0	0.2	98.4	100.0	1,321
Middle	0.0	0.3	0.0	1.1	7.1	3.3	0.0	14.0	0.0	0.6	0.0	73.6	100.0	1,321
Fourth	0.0	0.0	3.1	3.7	29.4	2.5	0.0	26.9	0.5	0.4	0.0	33.4	100.0	1,314
Richest	0.6	0.6	8.1	19.5	39.5	7.5	0.1	13.3	0.0	0.0	1.2	9.5	100.0	1,319
Ethnicity of household head														
Turkana	0.0	0.1	2.0	4.4	9.9	1.7	0.0	6.2	0.1	0.2	0.3	74.9	100.0	5,483
Other ethnic group	0.8	0.5	3.3	6.2	41.8	7.7	0.1	35.7	0.0	0.0	0.0	3.8	100.0	1,093

The MDGs and the WHO/UNICEF Joint Monitoring Programme (JMP) for Water Supply and Sanitation classify otherwise acceptable sanitation facilities which are public or shared between two or more households as unimproved. Therefore, “use of improved sanitation” is used both in the context of this report and as an MDG indicator to refer to improved sanitation facilities, which are not public or shared. Data on the use of improved sanitation are presented in Tables WS.6 and WS.7.

Table WS.6 shows that 25 percent of the household population is using an improved sanitation facility (12 percent not shared and 13 percent public or shared with other households). Figure WS.2 presents the distribution of the survey population by use and sharing of sanitation facilities.


Table WS.6: Use and sharing of sanitation facilities

Percent distribution of household population by use of private and public sanitation facilities and use of shared facilities, by users of improved and unimproved sanitation facilities, Turkana County MICS, 2013/14

	Users of improved sanitation facilities					Users of unimproved sanitation facilities					Open defecation (no facility, bush, field)	Total	Number of household members
	Not shared ¹	Public facility	Shared by		Missing/DK	Not shared	Public facility	Shared by					
			5 households or less	More than 5 households				5 households or less	More than 5 households				
Total	12.3	1.6	7.4	3.6	0.4	6.1	1.3	2.9	1.5	63.0	100.0	6,594	
Area													
Urban	20.0	2.1	12.9	6.2	0.7	8.3	2.2	4.8	2.1	40.5	100.0	3,598	
Rural	2.9	1.0	0.7	0.5	0.0	3.4	0.2	0.5	0.7	90.1	100.0	2,996	
Education of household head													
None	5.6	0.6	3.9	0.5	0.3	5.4	1.4	1.6	0.6	80.1	100.0	4,269	
Primary	15.1	4.6	13.2	7.7	1.4	5.7	0.3	3.6	2.1	46.2	100.0	1,067	
Secondary+	33.1	2.5	14.5	11.4	0.0	9.1	1.9	7.0	3.5	17.1	100.0	1,187	
Wealth index quintile													
Poorest	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	100.0	100.0	1,319	
Second	0.0	0.0	0.0	0.0	0.0	1.0	0.4	0.2	0.0	98.4	100.0	1,321	
Middle	5.4	0.9	5.4	0.3	0.0	8.6	2.2	3.0	0.7	73.6	100.0	1,321	
Fourth	16.5	6.8	10.7	3.6	1.1	12.4	3.4	7.8	4.3	33.4	100.0	1,314	
Richest	39.5	0.4	20.8	14.4	0.8	8.3	0.5	3.3	2.5	9.5	100.0	1,319	
Ethnicity of household head													
Turkana	9.2	1.4	5.7	1.7	0.3	3.4	0.5	2.2	0.9	74.9	100.0	5,483	
Other ethnic group	27.8	2.9	15.9	12.7	1.0	19.6	5.4	6.4	4.4	3.8	100.0	1,093	

¹ MICS indicator 4.3; MDG indicator 7.9 - Use of improved sanitation

Figure WS.2: Percent distribution of household members by use and sharing of sanitation facilities, Turkana County MICS, 2013/14



Having access to both an improved drinking water source and an improved sanitation facility brings the largest public health benefits to a household.⁶⁵ In its 2008 report,⁶⁶ the JMP developed a new way of presenting the access figures, by disaggregating and refining the data on drinking-water and sanitation and reflecting them in "ladder" format. This ladder allows a disaggregated analysis of trends in a three rung ladder for drinking-water and a four-rung ladder for sanitation. For sanitation, this gives an understanding of the proportion of population with no sanitation facilities at all – who revert to open defecation, of those reliant on technologies defined by JMP as "unimproved," of those sharing sanitation facilities of otherwise acceptable technology, and those using "improved" sanitation facilities.

Table WS.7 presents the percentages of household population by these drinking water and sanitation ladders. The table also shows the percentage of household members using both improved sources of

⁶⁵Wolf, J et al. 2014. *Systematic review: Assessing the impact of drinking water and sanitation on diarrhoeal disease in low- and middle-income settings: systematic review and meta-regression*. Tropical Medicine and International Health 2014.

DfID. 2013. *Water, Sanitation and Hygiene: Evidence Paper*. DfID: <http://r4d.dfid.gov.uk/pdf/outputs/sanitation/WASH-evidence-paper-april2013.pdf>

⁶⁶WHO/UNICEF JMP. 2008. *MDG assessment report*. http://www.wssinfo.org/fileadmin/user_upload/resources/1251794333-JMP_08_en.pdf

drinking water⁶⁷ and an improved sanitary means of excreta disposal. Approximately, 72 percent of household members use an improved drinking water source. The use of improved water sources is higher in urban than rural areas and improves with the education level and household wealth of the head of the household.

Twelve percent of household members use improved sanitation while 13 percent use shared improved sanitation facilities. About 12 percent of the household population uses both improved drinking water sources and improved sanitation facilities (20 percent in urban and three percent in rural areas). These results are presented by household wealth quintiles in Figure WS.3 and by urban/rural areas in Figure WS.4.

⁶⁷Those indicating bottled water as the main source of drinking water are distributed according to the water source used for other purposes such as cooking and handwashing.

Table WS.7: Drinking water and sanitation ladders

Percentage of household population by drinking water and sanitation ladders, Turkana County MICS, 2013/14

	Percentage of household population using:										Number of household members
	Improved drinking water ^{1, a}				Unimproved sanitation					Improved drinking water sources and improved sanitation	
	Piped into dwelling, plot or yard	Other improved	Unimproved drinking water	Total	Improved sanitation ²	Shared improved facilities	Unimproved facilities	Open defecation	Total		
Total	20.1	51.7	28.1	100.0	12.3	13.0	11.7	63.0	100.0	12.2	6,594
Area											
Urban	35.1	50.5	14.5	100.0	20.0	22.0	17.5	40.5	100.0	19.9	3,598
Rural	2.2	53.2	44.6	100.0	2.9	2.3	4.8	90.1	100.0	2.9	2,996
Education of household head											
None	7.2	55.9	36.9	100.0	5.6	5.2	9.0	80.1	100.0	5.6	4,269
Primary	25.5	56.0	18.5	100.0	15.1	26.9	11.7	46.2	100.0	15.1	1,067
Secondary+	58.7	34.4	6.9	100.0	33.1	28.3	21.5	17.1	100.0	32.8	1,187
Wealth index quintile											
Poorest	0.0	29.3	70.7	100.0	0.0	0.0	0.0	100.0	100.0	0.0	1,319
Second	0.0	60.8	39.2	100.0	0.0	0.0	1.6	98.4	100.0	0.0	1,321
Middle	1.3	77.2	21.5	100.0	5.4	6.5	14.5	73.6	100.0	5.4	1,321
Fourth	22.0	70.2	7.8	100.0	16.5	22.2	27.8	33.4	100.0	16.2	1,314
Richest	77.4	21.0	1.5	100.0	39.5	36.4	14.7	9.5	100.0	39.5	1,319
Ethnicity of household head											
Turkana	19.0	47.7	33.3	100.0	9.2	9.0	6.9	74.9	100.0	9.1	5,483
Other ethnic group	25.5	72.2	2.3	100.0	27.8	32.5	35.8	3.8	100.0	27.8	1,093
	¹ MICS indicator 4.1; MDG indicator 7.8 - Use of improved drinking water sources										
	² MICS indicator 4.3; MDG indicator 7.9 - Use of improved sanitation										
	^a Those indicating bottled water as the main source of drinking water are distributed according to the water source used for other purposes such as cooking and handwashing.										

Figure WS.3: Use of Improved drinking water sources and Improved sanitation facilities by household members, Turkana County MICS, 2013/14





Figure 4: Use of Improved water and sanitation in urban and rural areas, Turkana, 2013/14



Safe disposal of a child’s faeces is disposing of the stool, by the child using a toilet or by rinsing the stool into a toilet or latrine. Putting disposable diapers with solid waste, a very common practice throughout the world has thus far been classified as an inadequate means of disposal of child faeces for concerns about poor disposal of solid waste itself. This classification is currently under review. Disposal of faeces of children 0-2 years of age is presented in Table WS.8. In 27 percent of the cases, children’s stool is disposed of safely (46 percent in urban areas and 6 percent in rural areas).

Table WS.8: Disposal of child's faeces

Percent distribution of children age 0-2 years according to place of disposal of child's faeces, and the percentage of children age 0-2 years whose stools were disposed of safely the last time the child passed stools, Turkana County MICS, 2013/14

	Place of disposal of child's faeces							Total	Percentage of children whose last stools were disposed of safely ¹	Number of children age 0-2 years
	Put/rinsed into toilet or latrine	Put/rinsed into drain or ditch	Thrown into garbage	Buried	Left in the open	Other	Missing/DK			
Total	26.8	2.4	23.6	40.3	1.4	2.5	3.1	100.0	26.8	632
Type of sanitation facility used by household members										
Improved	81.8	1.1	4.9	9.6	0.0	0.0	2.5	100.0	81.8	143
Unimproved	71.4	0.9	13.8	10.3	0.0	0.0	3.6	100.0	71.4	65
Open defecation	1.6	3.0	31.3	55.2	2.0	3.7	3.2	100.0	1.6	425
Area										
Urban	46.0	1.7	20.1	26.2	0.2	3.0	2.7	100.0	46.0	325
Rural	6.4	3.0	27.3	55.3	2.6	1.9	3.5	100.0	6.4	307
Mother's education										
None	16.6	2.2	26.2	47.1	1.8	2.1	4.0	100.0	16.6	446
Primary	44.7	3.8	19.9	28.2	0.4	3.1	0.0	100.0	44.7	130
Secondary+	(68.3)	(0.0)	(11.5)	(12.8)	(0.0)	(4.3)	(3.1)	100.0	(68.3)	55
Wealth index quintile										
Poorest	0.0	2.1	25.5	64.1	1.7	2.3	4.3	100.0	0.0	125
Second	0.4	3.0	28.3	57.1	4.5	2.6	4.1	100.0	0.4	144
Middle	24.1	1.8	33.7	35.7	0.0	2.9	1.7	100.0	24.1	134
Fourth	49.5	2.3	16.8	24.8	0.0	3.4	3.2	100.0	49.5	133
Richest	73.8	2.6	9.0	12.0	0.0	0.8	1.8	100.0	73.8	95
Ethnicity of household head										
Turkana	15.5	2.6	26.1	47.7	1.6	3.0	3.4	100.0	15.5	533
Other ethnic group	(*)	(*)	(*)	(*)	(*)	(*)	(*)	100.0	(*)	98
¹ MICS indicator 4.4 - Safe disposal of child's faeces										
() Figures that are based on 25-49 unweighted cases										
(*) Figures that are based on fewer than 25 unweighted cases										

6.3 Handwashing

Handwashing with water and soap is the most cost effective health intervention to reduce the incidence of both diarrhoea and pneumonia in children under five.⁶⁸ It is most effective when done using water and soap after visiting a toilet or cleaning a child, before eating or handling food and, before feeding a child. Monitoring correct handwashing behaviour at these critical times is challenging. A reliable alternative to observations or self-reported behaviour is assessing the likelihood that correct handwashing behaviour takes place by asking if a household has a specific place where people wash their hands and, if yes, observing whether water and soap (or other local cleansing materials) are available at this place.⁶⁹

⁶⁸Cairncross, S and Valdmanis, V. 2006. *Water supply, sanitation and hygiene promotion* Chapter 41 in *Disease Control Priorities in Developing Countries*. 2nd Edition, Edt. Jameson et al. The World Bank.

⁶⁹Ram, P et al. editors. 2008. *Use of a novel method to detect reactivity to structured observation for measurement of handwashing behavior*. American Society of Tropical Medicine and Hygiene.

In Turkana County, the percentage of households where a place for handwashing was observed is 4 percent. Eighty-seven percent of the households had no specific place for handwashing in the dwelling, yard, or plot (Table WS.9). The percentage of households where a place for handwashing was observed, and where both water and soap (or another cleansing agent) were present at the place for handwashing, was only three percent. The percentage of households with a specific handwashing place and water (but no soap) present at the place for handwashing, was less than one percent, while the percentage of households with a handwashing place and soap (but no water), was one percent. Finally, the percentage of households with a place for handwashing, but with neither water nor soap available at the specific place for handwashing, was less than one percent. Differentials were observed by urban/rural areas and by education of head of household and wealth category.

Table WS.9: Water and soap at place for handwashing

Percentage of households where place for handwashing was observed, percentage with no specific place for handwashing, and percent distribution of households by availability of water and soap at specific place for handwashing, Turkana County MICS, 2013/14

	<u>Percentage of households:</u>			<u>Place for handwashing observed</u>						Percentage of households with a specific place for handwashing where water and soap or other cleansing agent are present ¹	Number of households where place for handwashing was observed or with no specific place for handwashing in the dwelling, yard, or plot		
	Where place for handwashing was observed	With no specific place for handwashing in the dwelling, yard, or plot	Number of households	<u>Water is available and:</u>			<u>Water is not available and:</u>						
				Soap present	<u>No soap:</u>		Soap present	<u>No soap:</u>					
			Ash, mud, or sand present		No other cleansing agent present			Ash, mud, or sand present	No other cleansing agent present	No specific place for handwashing in the dwelling, yard, or plot	Total		
Total	4.3	87.0	1,277	3.1	0.3	0.6	0.4	0.1	0.3	95.2	100.0	3.3	1,167
Area													
Urban	5.9	85.0	684	5.4	0.0	0.8	0.2	0.0	0.0	93.5	100.0	5.4	621
Rural	2.6	89.3	593	0.4	0.5	0.3	0.7	0.3	0.5	97.2	100.0	0.9	545
Education of household head													
None	3.1	87.9	822	1.2	0.4	0.5	0.7	0.2	0.4	96.6	100.0	1.6	747
Primary	6.2	84.0	211	5.1	0.0	1.8	0.0	0.0	0.0	93.1	100.0	5.1	190
Secondary+	7.3	85.9	234	7.9	0.0	0.0	0.0	0.0	0.0	92.1	100.0	7.9	218
Wealth index quintile													
Poorest	1.8	85.3	256	0.0	0.7	0.0	0.7	0.3	0.3	97.9	100.0	0.7	223
Second	4.1	88.6	282	1.3	0.6	1.0	0.8	0.4	0.4	95.6	100.0	1.8	261
Middle	2.8	90.0	255	1.9	0.0	1.1	0.0	0.0	0.0	96.9	100.0	1.9	236
Fourth	5.5	87.2	239	4.2	0.0	0.8	0.5	0.0	0.5	94.0	100.0	4.2	222
Richest	7.7	83.7	245	8.4	0.0	0.0	0.0	0.0	0.0	91.6	100.0	8.4	224
Ethnicity of household head													
Turkana	3.2	87.9	1,046	2.3	0.3	0.2	0.4	0.2	0.2	96.5	100.0	2.6	953
Other ethnic group	9.7	82.7	226	6.9	0.0	2.5	0.6	0.0	0.5	89.5	100.0	6.9	209

¹ MICS indicator 4.5 - Place for handwashing

Table WS.10 presents the percent distribution of households by availability of soap or other cleansing agent in the dwelling. The percentage of households with soap or other cleansing agent anywhere in the dwelling was 34 percent. In households where a place of handwashing was observed, 30 percent were not able or did not want to show soap or other cleansing agent. In households where a place of handwashing was not observed, 12 percent of households were not able or were unwilling to show soap or other cleansing agent in the dwelling. The percentage of households with soap or other cleansing agent anywhere in the house was higher in urban areas (45 percent) than rural areas (21 percent). The percentage was highest for households in the richest wealth quintile (71 percent) compared to the poorest quintile (6 percent).

Table WS.10: Availability of soap or other cleansing agent

Percent distribution of households by availability of soap or other cleansing agent in the dwelling, Turkana County MICS, 2013/14

	Place for handwashing observed				Place for handwashing not observed			Total	Percentage of households with soap or other cleansing agent anywhere in the dwelling ¹	Number of households
	Soap or other cleansing agent observed	Soap or other cleansing agent not observed at place for handwashing			No soap or other cleansing agent in household	Not able/Does not want to show soap or other cleansing agent	Total			
		Soap or other cleansing agent shown	No soap or other cleansing agent in household	Not able/Does not want to show soap or other cleansing agent						
Total	3.6	0.1	0.7	30.2	53.2	12.3	100.0	33.8	1,277	
Area										
Urban	5.1	0.0	0.8	39.9	38.1	16.2	100.0	45.0	684	
Rural	1.8	0.2	0.7	18.9	70.7	7.8	100.0	20.9	593	
Education of household head										
None	2.3	0.1	0.7	20.7	66.8	9.5	100.0	23.1	822	
Primary	4.6	0.0	1.6	41.4	37.0	15.4	100.0	46.0	211	
Secondary+	7.3	0.0	0.0	53.5	20.9	18.3	100.0	60.8	234	
Wealth index quintile										
Poorest	1.5	0.0	0.3	4.9	92.5	0.8	100.0	6.4	256	
Second	2.8	0.3	1.0	16.5	70.4	8.9	100.0	19.6	282	
Middle	1.8	0.0	1.0	28.3	51.8	17.1	100.0	30.1	255	
Fourth	4.3	0.0	1.2	41.2	35.9	17.4	100.0	45.6	239	
Richest	7.7	0.0	0.0	63.3	10.8	18.2	100.0	71.0	245	
Ethnicity of household head										
Turkana	2.9	0.1	0.3	26.9	58.3	11.6	100.0	29.9	1,046	
Other ethnic group	6.9	0.0	2.8	45.1	30.2	15.1	100.0	52.0	226	

¹ MICS indicator 4.6 - Availability of soap or other cleansing agent

7. Reproductive Health

The 1994 International Conference on Population and Development (ICPD) affirmed that respect, protection, promotion and fulfilment of human rights are necessary preconditions for improving the dignity and well-being of women and adolescent girls and for empowering them to exercise their reproductive rights; and that sexual and reproductive health and rights and understanding the implications of population dynamics are foundational to sustainable development.⁷⁰ Kenya is signatory to a number of international and regional conventions that aim to address sexual and reproductive rights of men, women, boys and girls including the ICPD 1994 and Campaign on Accelerated Reduction of Maternal Mortality in Africa (CARMMA) (2009).

Notable policies and strategies developed since the 1994 Cairo meeting include the Contraceptive Policy and Strategy (2002-2006); the Adolescent Reproductive Health and Development Policy, 2003; the Contraceptive Commodities Procurement Plan (2003-2006); National Reproductive Health Policy, 2007; the Contraceptive Commodities Security Strategy (2007-2012); the National Reproductive Health Policy Enhancing Reproductive Health Status for all Kenyans, 2007; the National Reproductive Health and HIV and AIDS integration Strategy-August 2009; the HIV and AIDS Strategic Plan (2009/10-2012/13); the National Condom Policy and Strategy (2009-2014); the National Road Map for Accelerating the Attainment of the MDGs Related to Maternal and Newborn Health in Kenya, August 2010; the National Reproductive Health Strategy 2009-2015; the Constitution of Kenya 2010 that for the first time guarantees the right to health care including reproductive health; the School Health Policy 2009⁷¹; and the Kenya National Population Policy 2012.⁷²

This chapter presents results on the following topics: fertility; contraception; unmet need for contraception; antenatal care (ANC); assistance at and place of delivery; and post-natal checks (PNC).

7.1 Fertility

Measures of current fertility are presented in Table RH.1 for the three-year period preceding the survey. The Turkana MICS used birth history to derive current fertility rates. The main shortcomings associated with birth histories besides possible sampling errors, are response errors (e.g. age misstatements, misdating of events and omissions of births and deaths).⁷³ A three-year period was chosen for calculating these rates to provide the most current information while also allowing the rates to be calculated for a sufficient number of cases so as not to compromise the statistical precision of the estimates. Age-specific fertility rates (ASFRs), expressed as the number of live births per 1,000 women in a specified age group, show the age pattern of fertility. Numerators for ASFRs are calculated by identifying live births that occurred in the three-year period preceding the survey classified

⁷⁰Framework of Actions for the follow - up to the Programme of Action of the International Conference on Population and Development Beyond 2014

⁷¹ Government of Kenya. National School Health Policy. Ministry of Public Health and Sanitation and Ministry of Education. Nairobi: Republic of Kenya; 2009.

⁷²Kenya National Commission for Human Rights. 2012. Realising Sexual and Reproductive Health Rights in Kenya: A myth or reality? A Report of the Public Inquiry into Violations of Sexual and Reproductive Health Rights in Kenya April 2012.

⁷³ Samuel Gaisie. Fertility Trend in Ghana. African Population Studies Vol. 20 N°2/Etude de la population africaine vol. 20 n° 2

according to the age of the mother (in five-year age groups) at the time of the child's birth. The denominators of the rates represent the number of woman-years lived by the survey respondents in each of the five-year age groups during the specified period.

The total fertility rate (TFR) is a measure that denotes the number of live births a woman would have if she were subject to the current age-specific fertility rates throughout her reproductive years (15-49 years).

The general fertility rate (GFR) is the number of live births occurring during the specified period per 1,000 women age 15-49 years.

The crude birth rate (CBR) is the number of live births per 1,000 population during the specified period.

Table RH.1 shows current fertility in Turkana County according to the type of place of residence. The TFR for the three years preceding the survey is 6 births per woman. Fertility is considerably higher in rural areas (7.5 births per woman) than in urban areas (4.7 births per woman).

Table RH.1: Fertility rates			
Adolescent birth rate, age-specific and total fertility rates, the general fertility rate, and the crude birth rate for the three-year period preceding the survey, by area, Turkana County MICS, 2013/14			
	Urban	Rural	Total
Age			
15-19 ¹	104	(93)	101
20-24	221	259	234
25-29	(217)	357	286
30-34	(247)	(277)	260
35-39	(94)	(238)	166
40-44	(*)	(146)	(91)
45-49	(*)	(*)	(*)
TFR ^a	(*)	(7.5)	6.0
GFR ^b	155.8	243.5	190.2
CBR ^c	35.7	39.7	38.0
¹ MICS indicator 5.1; MDG indicator 5.4 - Adolescent birth rate			
^a TFR: Total fertility rate expressed per woman age 15-49 years			
^b GFR: General fertility rate expressed per 1,000 women age 15-49 years			
^c CBR: Crude birth rate expressed per 1,000 population			
() Figures that are based on 125 to 249 unweighted cases			
(*) Figures that are based on less than 125 unweighted cases			

The overall age pattern of fertility, as reflected in the ASFRs, indicates that childbearing began early. Fertility rates among adolescents at 101 births per 1,000 women, increase to a peak of 286 births per 1,000 among women age 25-29 years, and declines thereafter.

Table RH.2 shows adolescent birth rates and total fertility rates. The adolescent birth rate (age-specific fertility rate for women age 15-19 years) is defined as the number of births to women age 15-19 years during the three-year period preceding the survey, divided by the average number of women age 15-

19 years (number of women-years lived between ages 15 through 19 years, inclusive) during the same period, expressed per 1,000 women.

Table RH.2: Adolescent birth rate and total fertility rate		
Adolescent birth rates and total fertility rates for the three-year period preceding the survey, Turkana County MICS, 2013/14		
	Adolescent birth rate ¹ (Age-specific fertility rate for women age 15-19 years)	Total fertility rate
Total	101	6.0
Education		
None	99	6.8
Primary	(134)	(5.1)
Secondary+	(51)	(2.9)
Ethnicity of household head		
Turkana	115	6.5
Other ethnic group	(51)	(4.3)
¹ MICS indicator 5.1; MDG indicator 5.4 - Adolescent birth rate		
() Figures that are based on 125 to 249 unweighted cases		

Table RH.3 presents some early childbearing⁷⁴ indicators for women age 15-19 years and 20-24 years while Table RH.4 presents the trends for early childbearing. As shown in Table RH.3, 16 percent of women age 15-19 years had begun childbearing, three percent were pregnant with their first child, and three percent have had a live birth before age 15. The table also presents that 29 percent of women age 20-24 years have had a live birth before age 18.

⁷⁴Childbearing is the process of giving birth to children. While early childbearing is defined as having had live births before specific young ages, for the purposes of Table RH.3, women age 15-19 years who have begun childbearing includes those who have had a live birth as well as those who have not had a live birth but are pregnant with their first child.

Table RH.3: Early childbearing

Percentage of women age 15-19 years who have had a live birth, are pregnant with the first child, have begun childbearing, and who have had a live birth before age 15, and percentage of women age 20-24 years who have had a live birth before age 18, Turkana County MICS, 2013/14

	Percentage of women age 15-19 years who:				Number of women age 15-19 years	Percentage of women age 20-24 years who have had a live birth before age 18 ¹	Number of women age 20-24 years
	Have had a live birth	Are pregnant with first child	Have begun childbearing	Have had a live birth before age 15			
Total	14.6	1.7	16.4	2.7	252	29.2	209
Area							
Urban	14.4	2.3	16.7	3.3	192	30.1	153
Rural	15.3	0.0	15.3	0.8	60	26.7	56
Education							
None	19.3	4.8	24.1	3.5	65	25.8	87
Primary	12.7	0.9	13.7	1.5	128	41.3	68
Secondary+	(13.7)	(0.0)	(13.7)	(4.4)	58	(19.5)	54
Ethnicity of household head							
Turkana	16.7	2.2	18.9	3.4	198	34.7	152
Other ethnic group	(6.9)	(0.0)	(6.9)	(0.0)	53	(14.4)	57
¹ MICS indicator 5.2 - Early childbearing							
() Figures that are based on 25-49 unweighted cases							

In the county six percent of women age 15-19 years have had a live birth before age 15 (Table RH.4). The proportion of women who had a live birth before age 15 is seven percent in urban areas and five percent in rural areas.

Table RH.4: Trends in early childbearing

Percentage of women who have had a live birth, by age 15 and 18, by area and age group, Turkana County MICS, 2013/14

	Urban				Rural				All			
	Percentage of women with a live birth before age 15	Number of women age 15-49 years	Percentage of women with a live birth before age 18	Number of women age 20-49 years	Percentage of women with a live birth before age 15	Number of women age 15-49 years	Percentage of women with a live birth before age 18	Number of women age 20-49 years	Percentage of women with a live birth before age 15	Number of women age 15-49 years	Percentage of women with a live birth before age 18	Number of women age 20-49 years
Total	6.5	683	29.0	491	5.4	421	23.3	361	6.1	1,104	26.6	852
Age												
15-19	3.3	192	na	na	0.8	60	na	na	2.7	252	na	na
20-24	5.5	153	30.1	153	3.3	56	26.7	56	4.9	209	29.2	209
25-29	19.9	111	40.2	111	5.3	99	23.3	99	13.0	210	32.2	210
30-34	6.2	71	30.1	71	8.9	73	32.1	73	7.6	144	31.1	144
35-39	1.2	75	17.9	75	5.7	59	13.5	59	3.2	134	15.9	134
40-44	(3.9)	38	(25.4)	38	3.9	44	22.3	44	3.9	82	23.7	82
45-49	(1.8)	44	(17.9)	44	(11.6)	31	(17.0)	31	5.8	75	17.5	75

na: not applicable

7.2 Contraception

Appropriate family planning is important to the health of women and children by: 1) preventing pregnancies that are too early or too late; 2) extending the period between births; and 3) limiting the total number of children. Access by all couples to information and services to prevent pregnancies that are too early, too closely spaced, too late or too many is critical.

The level of contraceptive use by women who are currently married or in union⁷⁵ is 14 percent (Table RH.5 and Figure RH.1). The most popular method is injectables which is used by one in ten married women in the county. Eighty-six percent of currently married/in union women are not using any method of contraception.

Twenty-three percent of married women in urban and five percent in rural areas use a method of contraception. Use of contraception according to the type of place and level of education are depicted in Figure RH.1. Use of contraception increases with both education and household wealth.

⁷⁵ All references to “married women” in this chapter include women in marital union as well.

Table RH.5: Use of contraception

Percentage of women age 15-49 years currently married or in union who are using (or whose partner is using) a contraceptive method, Turkana County MICS, 2013/14

	Percent of women currently married or in union who are using (or whose partner is using):																		Number of women age 15-49 years currently married or in union	
	No method	Female sterilization	Male sterilization	IUD	Injectables	Implants	Pill	Male condom	Female condom	Diaphragm/Foam/Jelly	LAM	Periodic abstinence	Withdrawal	Other	Missing	Any modern method	Any traditional method	Any method ¹		
Total	85.9	0.4	0.0	0.0	9.8	1.1	1.8	0.3	0.0	0.0	0.0	0.5	0.0	0.1	0.2	13.3	0.5	14.1	615	
Area																				
Urban	77.3	0.8	0.0	0.0	16.0	2.1	2.3	0.5	0.0	0.0	0.0	0.9	0.0	0.0	0.0	21.8	0.9	22.7	320	
Rural	95.3	0.0	0.0	0.0	2.9	0.0	1.2	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.4	4.1	0.2	4.7	295	
Age																				
15-19	(85.0)	(0.0)	(0.0)	(0.0)	(11.7)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(3.3)	(0.0)	(0.0)	(0.0)	(11.7)	(3.3)	(15.0)	34	
20-24	73.0	0.0	0.0	0.0	18.5	3.5	3.5	1.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	27.0	0.0	27.0	110	
25-29	87.6	0.0	0.0	0.0	9.7	1.0	0.6	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.8	11.3	0.3	12.4	166	
30-34	86.1	1.5	0.0	0.0	7.5	1.1	2.3	0.0	0.0	0.0	0.0	1.5	0.0	0.0	0.0	12.4	1.5	13.9	114	
35-39	87.1	1.0	0.0	0.0	9.2	0.0	2.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	12.9	0.0	12.9	95	
40-44	96.1	0.0	0.0	0.0	2.4	0.0	1.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	3.9	0.0	3.9	52	
45-49	(97.9)	(0.0)	(0.0)	(0.0)	(2.1)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(2.1)	(0.0)	(2.1)	43	
Number of living children																				
0	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	23
1	83.2	0.0	0.0	0.0	11.0	3.8	0.6	0.0	0.0	0.0	0.0	1.4	0.0	0.0	0.0	15.4	1.4	16.8	83	
2	78.0	0.8	0.0	0.0	19.4	0.0	1.3	0.0	0.0	0.0	0.0	0.0	0.0	.4	0.0	21.6	0.4	22.0	122	
3	85.8	1.7	0.0	0.0	10.5	1.9	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	14.2	0.0	14.2	99	
4+	89.6	0.0	0.0	0.0	5.8	0.6	3.0	0.0	0.0	0.0	0.0	0.6	0.0	0.0	0.4	9.4	0.6	10.4	288	
Education																				
None	92.0	0.0	0.0	0.0	5.5	0.4	1.6	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.3	7.5	0.1	8.0	408	
Primary	81.6	1.4	0.0	0.0	13.1	0.6	2.3	0.0	0.0	0.0	0.0	0.9	0.0	0.0	0.0	17.4	0.9	18.4	121	
Secondary+	63.1	1.1	0.0	0.0	24.9	5.1	1.9	1.8	0.0	0.0	0.0	2.0	0.0	0.0	0.0	34.9	2.0	36.9	86	
Wealth index quintile																				
Poorest	98.9	0.0	0.0	0.0	0.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.0	0.6	0.5	1.1	115	


Second	95.5	0.0	0.0	0.0	2.3	0.0	1.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.0	3.4	0.0	4.5	123
Middle	90.9	0.0	0.0	0.0	9.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	9.1	0.0	9.1	104
Fourth	77.6	0.0	0.0	0.0	18.3	0.0	3.3	0.0	0.0	0.0	0.0	0.8	0.0	0.0	0.0	21.6	0.8	22.4	138
Richest	70.9	2.0	0.0	0.0	16.1	5.0	3.6	1.2	0.0	0.0	0.0	1.3	0.0	0.0	0.0	27.8	1.3	29.1	135
Ethnicity of household head																			
Turkana	87.3	0.5	0.0	0.0	9.0	1.3	1.0	0.0	0.0	0.0	0.0	0.5	0.0	0.1	0.2	11.8	0.6	12.7	526
Other ethnic group	77.8	0.0	0.0	0.0	14.3	0.0	6.2	1.8	0.0	0.0	0.0	0.0	0.0	0.0	0.0	22.2	0.0	22.2	88

¹ MICS indicator 5.3; MDG indicator 5.3 - Contraceptive prevalence rate

() Figures that are based on 25-49 unweighted cases

(*) Figures that are based on fewer than 25 unweighted cases

Figure RH.1: Differentials in contraceptive use, Turkana County MICS, 2013/14



7.3 Unmet Need

Unmet need for contraception refers to fecund women who are married or in union and are not using any method of contraception, but who wish to postpone the next birth (spacing) or who wish to stop childbearing altogether (limiting). Unmet need is identified in MICS by using a set of questions eliciting current behaviours and preferences pertaining to contraceptive use, fecundity, and fertility preferences.

Unmet need for spacing is defined as the percentage of women who are married or in union and are not using a method of contraception and:

- are not pregnant, and not postpartum amenorrheic⁷⁶, and are fecund⁷⁷, and say they want to wait two or more years for their next birth OR
- are not pregnant, and not postpartum amenorrheic, and are fecund, and unsure whether they want another child OR
- are pregnant, and say that pregnancy was mistimed: would have wanted to wait OR
- are postpartum amenorrheic, and say that the birth was mistimed: would have wanted to wait.

⁷⁶ A woman is postpartum amenorrheic if she had a birth in last two years and is not currently pregnant, and her menstrual period has not returned since the birth of the last child

⁷⁷ A woman is considered infecund if she is neither pregnant nor postpartum amenorrheic, and (1a) has not had menstruation for at least six months, or (1b) never menstruated, or (1c) her last menstruation occurred before her last birth, or (1d) in menopause/has had hysterectomy OR
 (2) She declares that she has had hysterectomy, or that she has never menstruated, or that she is menopausal, or that she has been trying to get pregnant for 2 or more years without result in response to questions on why she thinks she is not physically able to get pregnant at the time of survey OR
 (3) She declares she cannot get pregnant when asked about desire for future birth OR
 (4) She has not had a birth in the preceding 5 years, is currently not using contraception and is currently married and was continuously married during the last 5 years preceding the survey.

Unmet need for limiting is defined as percentage of women who are married or in union and are not using a method of contraception AND

- are not pregnant, and not postpartum amenorrheic, and are fecund, and say they do not want any more children OR
- are pregnant, and say they did not want to have a child OR
- are postpartum amenorrheic, and say that they did not want the birth.

Total unmet need for contraception is the sum of unmet need for spacing and unmet need for limiting. This indicator is also known as unmet need for family planning and is one of the indicators used to track progress toward the MDG 5 of improving maternal health.

Met need for limiting includes women married or in union who are using (or whose partner is using) a contraceptive method,⁷⁸ and who want no more children, are using male or female sterilization, or declare themselves as infecund. Met need for spacing includes women who are using (or whose partner is using) a contraceptive method, and who want to have another child, or are undecided whether to have another child. The total of met need for spacing and limiting adds up to the total met need for contraception.

Using information on contraception and unmet need, the percentage of demand for contraception satisfied is also estimated from the MICS data. The percentage of demand satisfied is defined as the proportion of women currently married or in union who are currently using contraception, over the total demand for contraception. The total demand for contraception includes women who currently have an unmet need (for spacing or limiting), plus those who are currently using contraception.

Table RH.6 shows the levels of met need for contraception, unmet need, and the demand for contraception satisfied. The results show that the total met need is 14 percent, while total unmet need for family planning is 34 percent. Unmet need is equally high among urban women (36 percent) and rural women (32 percent). The table further highlights that the total demand for family planning satisfied is 29 percent.

⁷⁸ In this chapter, whenever reference is made to the use of a contraceptive by a woman, this may refer to her partner using a contraceptive method (such as male condom).

Table RH.6: Unmet need for contraception

Percentage of women age 15-49 years currently married or in union with an unmet need for family planning and percentage of demand for contraception satisfied, Turkana County MICS, 2013/14

	Met need for contraception			Unmet need for contraception			Number of women currently married or in union	Percentage of demand for contraception satisfied	Number of women currently married or in union with need for contraception
	For spacing	For limiting	Total	For spacing	For limiting	Total ¹			
Total	10.3	3.8	14.1	23.5	10.5	34.0	615	29.2	296
Area									
Urban	16.6	6.0	22.7	24.4	11.2	35.5	320	38.9	186
Rural	3.4	1.3	4.7	22.5	9.9	32.3	295	12.7	109
Age									
15-19	(15.0)	(0.0)	(15.0)	(48.6)	(0.0)	(48.6)	34	(*)	22
20-24	24.4	2.5	27.0	25.1	5.2	30.3	110	47.1	63
25-29	9.9	2.5	12.4	30.7	5.5	36.2	166	25.5	81
30-34	6.7	7.2	13.9	23.5	14.7	38.2	114	26.7	59
35-39	6.3	6.6	12.9	16.4	20.0	36.4	95	(26.1)	47
40-44	0.9	3.0	3.9	10.0	22.6	32.7	52	(*)	19
45-49	(2.1)	(0.0)	(2.1)	(3.2)	(5.4)	(8.5)	43	(*)	5
Education									
None	6.4	1.5	8.0	22.6	9.5	32.1	408	19.9	164
Primary	11.6	6.8	18.4	25.7	15.2	41.0	121	31.0	72
Secondary+	26.8	10.1	36.9	24.3	8.9	33.2	86	(52.6)	60
Wealth index quintile									
Poorest	1.1	0.0	1.1	28.5	8.7	37.2	115	2.8	44
Second	4.5	0.0	4.5	22.0	9.4	31.4	123	12.5	44
Middle	7.4	1.7	9.1	19.7	11.9	31.6	104	(22.4)	42
Fourth	16.8	5.6	22.4	29.7	12.7	42.4	138	34.5	89
Richest	19.1	10.0	29.1	17.1	9.8	26.9	135	52.0	76
Ethnicity of household head									
Turkana	9.4	3.4	12.7	23.2	11.1	34.3	526	27.1	247
Other ethnic group	16.0	6.2	22.2	24.7	7.6	32.3	88	(40.8)	48
¹ MICS indicator 5.4; MDG indicator 5.6 - Unmet need									
() Figures that are based on 25-49 unweighted cases									
(*) Figures that are based on fewer than 25 unweighted cases									

7.4 Antenatal Care

The antenatal period presents important opportunities for reaching pregnant women with a number of interventions that may be vital to the health and well-being of both mother and that of their unborn baby. Better understanding of foetal growth and development and its relationship to the mother's health has resulted in increased attention to the potential of antenatal care as an intervention to improve both maternal and newborn health. For example, antenatal care (ANC) can be used to inform women and families about risks and symptoms in pregnancy. In addition, it can inform about the risks of labour and delivery, and therefore it may provide the route for ensuring that pregnant women do in practice, deliver with the assistance of a skilled health care provider. Antenatal visits also provide an opportunity to supply information on birth spacing, which is recognized as an important factor in improving unborn baby survival. Tetanus immunization during pregnancy can be life-saving for both the mother and the unborn baby.

The prevention and treatment of malaria among pregnant women, management of anaemia during pregnancy and treatment of sexually transmitted infections (STIs) can significantly improve foetal outcomes and improve maternal health. Adverse outcomes such as low birth weight can be reduced through a combination of interventions to improve women's nutritional status and prevent infections (e.g., malaria and STIs) during pregnancy. More recently, the potential of the ANC as an entry point for HIV prevention and care, in particular for the prevention of HIV transmission from mother to child, has led to renewed interest in access to and use of antenatal services.

WHO recommends a minimum of four antenatal visits based on a review of the effectiveness of different models of antenatal care. WHO guidelines are specific on the content on ANC visits, which include:

- Blood pressure measurement
- Urine testing for bacteriuria and proteinuria
- Blood testing to detect syphilis and severe anaemia
- Weight/height measurement (optional).

It is of crucial importance for pregnant women to start attending ANC visits as early in pregnancy as possible in order to prevent and detect pregnancy conditions that could affect both the woman and her unborn baby. ANC should therefore, continue throughout the entire pregnancy.

Antenatal care coverage indicators (at least one visit with a skilled provider and 4 or more visits with any providers) are used to track progress toward the MDG 5 of improving maternal health.

The type of personnel providing antenatal care to women age 15-49 years who gave birth in the two years preceding the survey is presented in Table RH.7. The results show that 94 percent of women had received ANC and 94 percent received it from a skilled provider. In Turkana County, most of the ANC was provided by a nurse or midwife (86 percent) while 8 percent of women were attended to by a medical doctor. About 98 percent of women in urban areas and 89 percent in rural areas received ANC from a skilled provider.

Table RH.7: Antenatal care coverage

Percent distribution of women age 15-49 years with a live birth in the last two years by antenatal care provider during the pregnancy for the last birth, Turkana County MICS, 2013/14

	Provider of antenatal care ^a					No antenatal care	Total	Any skilled provider ^{1,b}	Number of women with a live birth in the last two years
	Medical doctor	Nurse/Midwife	Community nurse	Relative/friend	Other/Missing				
Total	7.5	85.9	0.2	0.4	0.3	5.7	100.0	93.6	387
Area									
Urban	9.0	89.3	0.0	0.0	0.0	1.7	100.0	98.3	199
Rural	5.9	82.3	0.4	0.8	0.5	10.0	100.0	88.6	188
Mother's age at birth									
Less than 20	(4.7)	(94.2)	(0.0)	(0.0)	(0.0)	(1.1)	100.0	(98.9)	52
20-34	8.5	84.9	0.3	0.3	0.4	5.7	100.0	93.7	277
35-49	5.3	83.3	0.0	1.3	0.0	10.1	100.0	88.6	59
Education									
None	6.7	83.8	0.3	0.6	0.4	8.3	100.0	90.7	268
Primary	9.9	90.1	0.0	0.0	0.0	0.0	100.0	100.0	82
Secondary+	(8.2)	(91.8)	(0.0)	(0.0)	(0.0)	(0.0)	100.0	(100.0)	38
Wealth index quintile									
Poorest	5.7	78.4	1.0	1.1	0.0	13.8	100.0	85.1	69
Second	3.1	88.0	0.0	0.0	0.0	8.8	100.0	91.2	86
Middle	9.5	86.6	0.0	0.9	0.0	3.0	100.0	96.1	84
Fourth	8.3	89.3	0.0	0.0	1.2	1.2	100.0	97.6	85
Richest	11.7	85.8	0.0	0.0	0.0	2.5	100.0	97.5	63
Ethnicity of household head									
Turkana	7.6	84.9	0.2	0.5	0.3	6.5	100.0	92.7	327
Other ethnic group	7.0	91.3	0.0	0.0	0.0	1.7	100.0	98.3	59

¹ MICS indicator 5.5a; MDG indicator 5.5 - Antenatal care coverage

^aOnly the most qualified provider is considered in cases where more than one provider was reported.

^b Skilled providers include *Medical doctor* and *Nurse/Midwife*.

() Figures that are based on 25-49 unweighted cases

Table RH.8 shows the number of ANC visits during the latest pregnancy that took place within the last two years preceding the survey, regardless of provider, by selected characteristics. Approximately, 87 percent of mothers received ANC more than once and 44 percent of the mothers received ANC at least four times. The percentage of women from urban areas (53 percent) who received ANC four or more times was higher than those from rural areas (34 percent).

The table also provides information about the timing of the first antenatal care visit. Overall, 22 percent of women with a live birth in the last two years preceding the survey had their first ANC visit during the first trimester of their last pregnancy. The median month pregnant women registered for the first ANC visit is five months.

Table RH.8: Number of antenatal care visits and timing of first visit

Percent distribution of women age 15-49 years with a live birth in the last two years by number of antenatal care visits by any provider and by the timing of first antenatal care visits, Turkana County MICS, 2013/14

	<u>Percent distribution of women who had:</u>							<u>Percent distribution of women by number of months pregnant at the time of first antenatal care visit</u>							Number of women with a live birth in the last two years	Median months pregnant at first ANC visit	Number of women with a live birth in the last two years who had at least one ANC visit
	No antenatal care visits	One visit	Two visits	Three visits	4 or more visits ¹	Missing/DK	Total	No antenatal care visits	First trimester	4-5 months	6-7 months	8+ months	DK/Missing	Total			
Total	5.7	3.5	15.3	28.1	44.0	3.4	100.0	5.7	22.2	43.0	24.1	3.9	1.1	100.0	387	5	361
Area																	
Urban	1.7	1.5	10.1	29.8	53.1	3.8	100.0	1.7	22.8	46.3	24.5	3.0	1.7	100.0	199	5	192
Rural	10.0	5.6	20.9	26.3	34.3	2.9	100.0	10.0	21.5	39.5	23.7	4.9	0.5	100.0	188	5	168
Mother's age at birth																	
Less than 20	(1.1)	(0.0)	(28.8)	(36.5)	(30.5)	(3.1)	100.0	(1.1)	(14.4)	(33.8)	(38.9)	(8.7)	(3.1)	100.0	52	(5)	50
20-34	5.7	3.4	12.4	27.8	47.0	3.7	100.0	5.7	24.0	44.7	22.2	2.5	0.9	100.0	277	5	258
35-49	10.1	7.2	17.3	21.8	41.8	1.8	100.0	10.1	20.6	43.2	20.1	6.0	0.0	100.0	59	5	53
Education																	
None	8.3	4.4	15.7	25.6	43.9	2.1	100.0	8.3	25.8	39.5	21.6	3.9	0.9	100.0	268	5	243
Primary	0.0	2.0	18.4	31.9	43.1	4.6	100.0	0.0	13.4	51.8	27.1	5.6	2.1	100.0	82	5	80
Secondary+	(0.0)	(0.0)	(6.2)	(37.5)	(46.8)	(9.6)	100.0	(0.0)	(15.6)	(48.8)	(35.6)	(0.0)	(0.0)	100.0	38	(5)	38
Wealth index quintile																	
Poorest	13.8	3.8	23.1	29.5	24.5	5.3	100.0	13.8	21.7	28.8	30.1	1.9	3.7	100.0	69	5	57
Second	8.8	8.9	15.4	26.0	39.1	1.8	100.0	8.8	18.2	44.7	20.2	8.1	0.0	100.0	86	5	79
Middle	3.0	3.0	16.1	28.8	48.0	1.2	100.0	3.0	32.2	37.2	25.4	2.3	0.0	100.0	84	5	81
Fourth	1.2	.8	10.0	34.6	50.7	2.6	100.0	1.2	26.0	45.6	22.8	4.3	0.0	100.0	85	5	84
Richest	2.5	0.0	12.9	19.7	57.7	7.3	100.0	2.5	9.6	60.4	22.9	2.1	2.6	100.0	63	(5)	60
Ethnicity of household head																	
Turkana	6.5	4.1	17.2	30.1	38.8	3.3	100.0	6.5	19.7	42.3	25.6	4.6	1.3	100.0	327	5	302
Other ethnic group	1.7	0.0	5.3	16.9	72.5	3.6	100.0	1.7	36.1	46.1	16.1	0.0	0.0	100.0	59	(4)	58

¹ MICS indicator 5.5b; MDG indicator 5.5 - Antenatal care coverage

() Figures that are based on 25-49 unweighted cases

The coverage of key services that pregnant women were expected to receive during ANC visits is shown in Table RH.9. Among those women who had a live birth during the last two years preceding the survey, 70 percent had blood pressure checked, and urine and blood samples taken. Ninety percent reported that a blood sample was taken during antenatal care visits, another 90 percent had their blood pressure checked, and 72 percent had a urine specimen taken. Measuring of blood pressure and having urine and blood samples taken was higher for women in urban areas (90 percent) than for those in rural areas (49 percent).

Table RH.9: Content of antenatal care

Percentage of women age 15-49 years with a live birth in the last two years who, at least once, had their blood pressure measured, urine sample taken, and blood sample taken as part of antenatal care, during the pregnancy for the last birth, Turkana County MICS, 2013/14

	Percentage of women who, during the pregnancy of their last birth, had:				Number of women with a live birth in the last two years
	Blood pressure measured	Urine sample taken	Blood sample taken	Blood pressure measured, urine and blood sample taken ¹	
Total	89.8	71.6	90.3	70.2	387
Area					
Urban	98.3	90.4	97.1	90.4	199
Rural	80.7	51.7	83.2	48.8	188
Mother's age at birth					
Less than 20	(97.8)	(79.0)	(91.5)	(79.0)	52
20-34	90.3	74.2	90.9	72.6	277
35-49	80.1	52.6	86.4	50.8	59
Education					
None	85.2	67.2	86.9	65.2	268
Primary	100.0	73.9	96.9	73.9	82
Secondary+	(100.0)	(97.3)	(100.0)	(97.3)	38
Wealth index quintile					
Poorest	78.2	57.6	83.1	54.8	69
Second	79.6	52.8	80.8	49.2	86
Middle	94.8	70.5	92.9	69.9	84
Fourth	98.8	90.4	98.8	90.4	85
Richest	97.5	88.6	96.4	88.6	63
Ethnicity of household head					
Turkana	88.2	66.7	88.8	65.0	327
Other ethnic group	98.3	98.3	98.3	98.3	59
¹ MICS indicator 5.6 - Content of antenatal care					
() Figures that are based on 25-49 unweighted cases					

7.5 Assistance at Delivery

About three quarters of all maternal deaths occur due to direct obstetric causes.⁷⁹ The single most critical intervention for safe motherhood is to ensure that a competent health worker with midwifery skills is present at every birth, and in case of emergency that transport is available to a referral facility for obstetric care. The skilled attendant at delivery is an indicator used to track progress toward the Millennium Development Goal 5 of improving maternal health.

The MICS included a number of questions to assess the proportion of births attended by a skilled attendant. A *skilled attendant* includes a doctor, nurse, or midwife. In Turkana, 35 percent of births occurring in the two years preceding the MICS were delivered by skilled personnel (Table RH.10 and Figure RH.2). Fifty-seven percent and 13 percent of women in urban and rural areas, respectively, were delivered by a skilled attendant.

Table RH.10 also shows information on women who delivered by caesarian section (C-section) and provides additional information on the timing of the decision to conduct a C-section (before labour pains began or after) in order to better assess if such decisions were mostly driven by medical or non-medical reasons. Overall, two percent of women who delivered in the last two years preceding the survey had a C-section and for nearly all of them, the decision was taken after the onset of labour pains.

⁷⁹ Say, L et al. 2014. *Global causes of maternal death: a WHO systematic analysis*. The Lancet Global Health2 (6): e323-33. DOI: 10.1016/S2214-109X(14)70227-X


Table RH.10: Assistance during delivery and caesarean section

Percent distribution of women age 15-49 years with a live birth in the last two years by person providing assistance at delivery, and percentage of births delivered by C-section, Turkana County MICS, 2013/14

	Person assisting at delivery								Total	Delivery assisted by any skilled attendant ^{1,a}	Percent delivered by C-section			Number of women who had a live birth in the last two years
	Medical doctor	Nurse/Midwife	Auxiliary midwife/Clinical Officer	Traditional birth attendant	Community health worker	Relative/Friend	Other/Missing	No attendant			Decided before onset of labour pains	Decided after onset of labour pains	Total ²	
Total	4.8	30.1	0.3	3.8	2.7	38.7	2.5	17.2	100.0	35.1	0.4	1.9	2.3	387
Area														
Urban	8.5	47.9	0.0	0.4	3.1	30.5	2.4	7.1	100.0	56.5	0.8	1.8	2.6	199
Rural	0.8	11.2	0.5	7.4	2.2	47.4	2.6	27.9	100.0	12.6	0.0	1.9	1.9	188
Mother's age at birth														
Less than 20	(9.3)	(44.0)	(0.0)	(0.9)	(0.9)	(37.3)	(1.4)	(6.2)	100.0	(53.3)	(0.0)	(6.8)	(6.8)	52
20-34	3.9	30.1	0.4	3.9	3.0	38.9	2.8	17.0	100.0	34.4	0.6	1.3	1.9	277
35-49	4.8	17.5	0.0	5.7	2.6	39.1	1.9	28.4	100.0	22.3	0.0	0.0	0.0	59
Place of delivery														
Home	0.0	1.1	0.0	5.9	4.2	60.2	2.6	25.9	100.0	1.1	0.0	0.0	0.0	247
Health facility	13.8	84.4	0.8	0.0	0.0	0.0	1.1	0.0	100.0	98.9	1.2	5.4	6.6	135
Education														
None	2.2	22.2	0.4	5.0	2.6	43.6	2.8	21.3	100.0	24.8	0.0	1.0	1.0	268
Primary	7.2	44.4	0.0	0.6	4.2	34.4	2.6	6.6	100.0	51.6	0.0	2.1	2.1	82
Secondary+	(18.1)	(54.9)	(0.0)	(2.1)	(0.0)	(13.4)	(0.0)	(11.5)	100.0	(73.0)	(4.3)	(7.4)	(11.7)	38
Wealth index quintile														
Poorest	0.0	8.9	0.0	5.2	2.2	49.4	2.3	32.1	100.0	8.9	0.0	0.0	0.0	69
Second	0.7	2.4	0.0	9.6	3.1	50.2	3.6	30.4	100.0	3.1	0.0	0.7	0.7	86
Middle	5.5	33.8	0.0	2.4	3.2	43.2	1.1	10.7	100.0	39.3	1.9	0.0	1.9	84
Fourth	8.4	47.6	1.2	0.9	4.1	26.7	3.0	8.2	100.0	57.2	0.0	5.9	5.9	85
Richest	9.9	62.5	0.0	0.0	0.0	21.4	2.3	4.0	100.0	72.4	0.0	2.7	2.7	63
Ethnicity of household head														
Turkana	2.3	24.1	0.0	4.2	3.2	44.2	2.5	19.5	100.0	26.4	0.0	0.9	0.9	327

Other ethnic group	18.5	63.7	1.8	1.3	0.0	7.4	2.4	4.9	100.0	83.9	2.8	7.1	9.8	59
¹ MICS indicator 5.7; MDG indicator 5.2 - Skilled attendant at delivery														
² MICS indicator 5.9 - Caesarean section														
^a Skilled attendants include <i>Medical doctor</i> and <i>Nurse/Midwife</i> .														
() Figures that are based on 25-49 unweighted cases														

Figure RH.2: Person assisting at delivery, Turkana County MICS, 2013/14



7.6 Place of Delivery

Increasing the proportion of births that are delivered in health facilities is an important factor in reducing the health risks to both the mother and the baby. Proper medical attention and hygienic conditions during delivery can reduce the risks of complications and infection that can cause morbidity and mortality to either the mother or the baby. Table RH.11 presents the percent distribution of women age 15-49 years who had a live birth in the two years preceding the survey by place of delivery, and the percentage of births delivered in a health facility, according to background characteristics.

Thirty-five percent of births in Turkana County were delivered in a health facility; 30 percent of deliveries occurred in public health facilities and five percent in private health facilities while 64 percent took place at home. The proportion of women in urban areas who delivered in a health facility was higher than in rural areas (57 percent compared with 12 percent).

Table RH.11: Place of delivery

Percent distribution of women age 15-49 years with a live birth in the last two years by place of delivery of their last birth, Turkana County MICS, 2013/14

	Place of delivery					Total	Delivered in health facility ¹	Number of women with a live birth in the last two years
	Health facility		Home	Other	Missing/DK			
	Public sector	Private sector						
Total	29.6	5.2	63.8	1.0	0.4	100.0	34.8	387
Area								
Urban	48.0	8.9	41.7	1.4	0.0	100.0	56.9	199
Rural	10.1	1.3	87.2	0.5	0.9	100.0	11.5	188
Mother's age at birth								
Less than 20	(48.7)	(4.6)	(45.3)	(1.4)	(0.0)	100.0	(53.3)	52
20-34	28.4	5.5	64.3	1.1	0.6	100.0	34.0	277
35-49	18.2	4.1	77.7	0.0	0.0	100.0	22.3	59
Number of antenatal care visits								
None	(0.0)	(2.6)	(93.2)	(0.0)	(4.2)	100.0	(2.6)	22
1-3 visits	23.6	0.0	75.6	0.9	0.0	100.0	23.6	182
4+ visits	38.8	10.5	49.7	0.6	0.4	100.0	49.3	170
Education								
None	19.7	4.3	74.4	1.0	0.6	100.0	24.0	268
Primary	44.0	8.6	47.4	0.0	0.0	100.0	52.6	82
Secondary+	(68.7)	(4.3)	(24.3)	(2.6)	(0.0)	100.0	(73.0)	38
Wealth index quintile								
Poorest	8.0	0.0	92.0	0.0	0.0	100.0	8.0	69
Second	0.7	0.0	97.5	1.0	0.8	100.0	0.7	86
Middle	33.2	6.0	58.4	1.2	1.1	100.0	39.3	84
Fourth	44.6	12.6	40.7	2.1	0.0	100.0	57.2	85
Richest	67.7	7.0	25.3	0.0	0.0	100.0	74.7	63
Ethnicity of household head								
Turkana	22.2	3.4	73.1	0.8	0.5	100.0	25.6	327
Other ethnic group	71.0	15.4	11.8	1.9	0.0	100.0	86.4	59
¹ MICS indicator 5.8 - Institutional deliveries								
() Figures that are based on 25-49 unweighted cases								

7.7 Post-natal Health Checks

The time of birth and immediately after is a critical window of opportunity to deliver lifesaving interventions for both the mother and newborn. Across the world, approximately 3 million newborns die annually in the first month of life⁸⁰ and the majority of these deaths occur within a day or two of birth⁸¹, which is also the time when the majority of maternal deaths occur.⁸²

Despite the importance of the first few days following birth, large-scale, nationally representative

⁸⁰UN Interagency Group for Child Mortality Estimation. 2013. *Levels and Trends in Child Mortality: Report 2013*

⁸¹ Lawn, JE et al. 2005. *4 million neonatal deaths: When? Where? Why?* Lancet 2005; 365:891–900.

⁸² WHO, UNICEF, UNFPA, The World Bank. 2012. *Trends in Maternal Mortality: 1990-2010*. World Health Organization.

household survey programmes have not systematically included questions on the post-natal period and care for the mother and newborn. In 2008, the Countdown to 2015 initiative, which monitors progress on maternal, newborn and child health interventions, highlighted this data gap. This not only called for post-natal care (PNC) programmes to be strengthened, but also for better data availability and quality.⁸³

Following the establishment and discussions of an Inter-Agency Group on PNC and drawing on lessons learned from earlier attempts of collecting PNC data, a new questionnaire module for MICS was developed and validated. The Post-natal Health Checks (PNHC) module collected information on newborns' and mothers' contact with a provider, but not content of care. The rationale for this is that as PNC programmes scale up, it is important to measure the coverage of that scale up and ensure that the platform for providing essential services is in place. Content is considered more difficult to measure, particularly because the respondent is asked to recall services delivered up to two years preceding the interview.

Table RH.12 presents the percent distribution of women age 15-49 years who gave birth in a health facility in the two years preceding the survey by duration of stay in the facility following the delivery.⁸⁴ Overall, 71 percent of women who gave birth in a health facility stayed 12 hours or more in the facility after delivery.

Table RH.12: Post-partum stay in health facility								
Percent distribution of women age 15-49 years with a live birth in the last two years who had their last birth delivered in a health facility by duration of stay in health facility, Turkana County MICS, 2013/14								
	<u>Duration of stay in health facility</u>					Total	12 hours or more ¹	Number of women who had their last birth delivered in a health facility in the last 2 years
	Less than 6 hours	6-11 hours	12-23 hours	1-2 days	3 days or more			
Total	17.7	11.7	6.4	49.4	14.8	100.0	70.6	135

¹ MICS indicator 5.10 - Post-partum stay in health facility

Safe motherhood programmes have recently increased emphasis on the importance of post-natal care, recommending that all women and newborns receive a health check within two days of delivery. *Health checks following birth* while in facility or at home refer to checks provided by any health provider regardless of timing (column 1). *Post-natal care (PNC) visits* on the other hand, refer to a separate visit to check on the health of the newborn and provide preventive care services. These, therefore, do not include *health checks following birth* while in facility or at home. The indicator *Post-natal health checks* includes any health check after birth received while in the health facility and at home (column 1), regardless of timing, as well as PNC visits within two days of delivery (columns 2, 3, and 4). To assess the extent of post-natal care utilization, women were asked whether they and their newborn received a health check after the delivery, the timing of the first check, and the type of health provider for the woman's last birth in the two years preceding the survey.

⁸³HMN, UNICEF, WHO. 2008. *Countdown to 2015: Tracking Progress in Maternal, Newborn & Child Survival, The 2008 Report*. UNICEF.

⁸⁴ Background characteristics variable were not included due to the small number of cases reported.

Table RH.13 shows the percentage of newborns born in the last two years preceding the survey who received health checks and post-natal care visits from any health provider after birth. Overall, 36 percent of newborns received a health check following birth while in a health facility or at home. With regards to PNC visits, these predominantly occurred on the same day as the delivery (7 percent), during the week after deliver (6 percent) and three percent after the first week following delivery. As a result, a total of 37 percent of all newborns received a post-natal health check. The proportion of urban newborns who received a health check, both following birth (56 percent) and in total including PNC visits (58 percent), was higher than that of their rural counterparts (14 percent and 15 percent, respectively). Among births that took place in a health facility, 94 percent had a health check following birth, and 5 percent for those that took place at home.⁸⁵

⁸⁵ Information on newborns who received the first PNC visit within one week of birth and type of provider of service is not included due to the small number of cases reported.

Table RH.13: Post-natal health checks for newborns

Percentage of women age 15-49 years with a live birth in the last two years whose last live birth received health checks while in facility or at home following birth, percent distribution whose last live birth received post-natal care (PNC) visits from any health provider after birth, by timing of visit, and percentage who received post-natal health checks, Turkana County MICS, 2013/14

	Health check following birth while in facility or at home ^a	PNC visit for newborns ^b						Total	Post-natal health check for the newborn ^{1, c}	Number of last live births in the last two years
		Same day	1 day following birth	2 days following birth	3-6 days following birth	After the first week following birth	No post-natal care visit			
Total	35.7	7.4	2.7	1.2	2.0	2.7	84.1	100.0	37.1	387
Area										
Urban	55.8	11.4	4.6	2.1	3.0	1.5	77.4	100.0	57.7	199
Rural	14.3	3.1	0.6	0.2	0.9	3.9	91.2	100.0	15.3	188
Mother's age at birth										
Less than 20	(52.8)	(14.1)	(0.0)	(0.0)	(0.0)	(0.9)	(85.0)	100.0	(54.1)	52
20-34	34.9	5.9	3.3	1.7	2.0	3.1	84.0	100.0	36.2	277
35-49	23.9	8.1	2.1	0.0	3.6	2.3	83.9	100.0	25.9	59
Place of delivery										
Home	4.7	1.5	1.3	0.6	0.8	1.5	94.4	100.0	6.7	247
Health facility	93.8	17.9	5.4	2.4	4.2	4.9	65.2	100.0	93.8	135
Public	92.7	15.5	6.3	2.8	2.1	3.8	69.5	100.0	92.7	115
Private	(*)	(*)	(*)	(*)	(*)	(*)	(*)	100.0	(*)	20
Education										
None	26.2	5.3	2.2	0.5	1.9	1.5	88.6	100.0	27.5	268
Primary	53.8	9.1	2.4	2.0	3.2	4.3	79.0	100.0	54.7	82
Secondary+	(63.4)	(17.9)	(7.0)	(4.3)	(0.0)	(7.2)	(63.6)	100.0	(66.7)	38
Wealth index quintile										
Poorest	11.8	4.4	0.0	0.0	1.8	1.5	92.3	100.0	12.8	69
Second	5.7	.6	1.4	0.5	0.5	2.9	94.1	100.0	7.0	86
Middle	39.4	4.7	1.9	0.0	2.1	0.0	91.3	100.0	39.4	84
Fourth	57.2	8.9	5.8	1.9	3.8	3.9	75.6	100.0	60.3	85
Richest	68.8	21.3	4.1	4.1	1.6	5.6	63.3	100.0	70.4	63
Ethnicity of household head										
Turkana	27.8	6.6	2.2	0.6	1.1	1.5	88.0	100.0	29.2	327
Other ethnic group	79.8	11.4	5.2	4.4	7.2	9.5	62.3	100.0	81.5	59

¹ MICS indicator 5.11 - Post-natal health check for the newborn

^aHealth checks by any health provider following facility births (before discharge from facility) or following home births (before departure of provider from home).

^bPost-natal care visits (PNC) refer to a separate visit by any health provider to check on the health of the newborn and provide preventive care services. PNC visits do not include health checks following birth while in facility or at home (see note ^a above).

^cPost-natal health checks include any health check performed while in the health facility or at home following birth (see note ^a above), as well as PNC visits (see note ^b above) within two days of delivery.

() Figures that are based on 25-49 unweighted cases

(*) Figures that are based on fewer than 25 unweighted cases

Table RH.14 presents information collected on post-natal health checks for mothers. Overall, 36 percent of mothers received a health check following birth while in a health facility or at home. Health checks following birth occurred mainly in health facility deliveries (96 percent), whereas for mothers delivering at home the figure was four percent.

Table RH.14: Post-natal health checks for mothers

Percentage of women age 15-49 years with a live birth in the last two years who received health checks while in facility or at home following birth, percent distribution who received post-natal care (PNC) visits from any health provider after birth at the time of last birth, by timing of visit, and percentage who received post-natal health checks, Turkana County MICS, 2013/14

	Health check following birth while in facility or at home ^a	PNC visit for mothers ^b								Post-natal health check for the mother ^{1, c}	Number of women with a live birth in the last two years
		Same day	1 day following birth	2 days following birth	3-6 days following birth	After the first week following birth	No post-natal care visit	Missing/DK	Total		
Total	36.1	3.0	0.9	0.9	2.4	4.3	87.7	0.9	100.0	37.1	387
Area											
Urban	56.7	4.5	1.4	1.3	2.9	1.0	88.1	0.8	100.0	57.2	199
Rural	14.4	1.4	0.5	0.5	1.8	7.7	87.2	0.9	100.0	15.7	188
Mother's age at birth											
Less than 20	(52.8)	(4.8)	(0.0)	(0.0)	(1.4)	(1.4)	(92.4)	(0.0)	100.0	(52.8)	52
20-34	35.6	2.5	1.3	1.1	2.3	4.5	87.0	1.2	100.0	36.7	277
35-49	23.9	3.6	0.0	0.9	3.6	5.6	86.3	0.0	100.0	24.7	59
Place of delivery											
Home	4.3	1.4	0.4	0.8	1.6	5.6	89.6	0.7	100.0	5.8	247
Health facility	95.8	6.1	2.0	1.2	3.9	2.0	83.6	1.2	100.0	95.8	135
Education											
None	25.9	2.1	1.4	0.7	1.9	4.8	88.4	0.6	100.0	27.2	268
Primary	53.8	4.3	0.0	0.0	4.1	0.7	90.9	0.0	100.0	53.8	82
Secondary+	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	100.0	(*)	38
Wealth index quintile											
Poorest	11.8	2.5	0.6	0.8	3.4	6.7	83.6	2.5	100.0	14.2	69
Second	4.6	1.1	0.6	0.5	1.2	8.0	88.6	0.0	100.0	5.7	86
Middle	39.4	0.9	1.9	0.0	0.0	1.1	94.0	1.9	100.0	39.4	84
Fourth	57.2	5.7	0.0	0.0	5.5	2.7	86.1	0.0	100.0	57.2	85
Richest	73.1	5.3	1.8	4.1	1.6	2.8	84.4	0.0	100.0	74.7	63
Ethnicity of household head											
Turkana	27.8	2.1	0.6	0.8	1.5	4.4	90.0	0.5	100.0	28.6	327
Other ethnic group	82.8	7.9	2.8	1.7	7.2	3.5	74.3	2.8	100.0	84.4	59

¹ MICS indicator 5.12 - Post-natal health check for the mother

^aHealth checks by any health provider following facility births (before discharge from facility) or following home births (before departure of provider from home).

^bPost-natal care visits (PNC) refer to a separate visit by any health provider to check on the health of the mother and provide preventive care services. PNC visits do not include health checks following birth while in facility or at home (see note ^a above).

^cPost-natal health checks include any health check performed while in the health facility or at home following birth (see note ^a above), as well as PNC visits (see note ^b above) within two days of delivery.

() Figures that are based on 25-49 unweighted cases

(*) Figures that are based on fewer than 25 unweighted cases

With regards to PNC visits, the visits were more or less the same for those that took place on the same day as the delivery, those that happened during the week, and after first week following delivery. A total of 37 percent of all mothers received a post-natal health check. The proportion of urban mothers receiving a health check, both following birth and in total including post-natal health checks (57 percent), was somewhat higher than that of their rural counterparts (56 percent and 58 percent,

respectively). PNC was also much higher for deliveries that happened in a health facility (96 percent) compared to those delivered at home (6 percent).⁸⁶

Table RH.15 and Figure RH.3 present the distribution of women who had a live birth in the two years preceding the survey by receipt of post-natal health checks within two days of birth for the mother and the newborn, thus combining the indicators presented in Tables RH.13 and RH.14. The Turkana County MICS showed that for 36 percent of live births, both the mothers and their newborns received either a post-natal health check following birth or a timely PNC visit. Sixty-two percent of births neither received post-natal health checks or timely visits.


Table RH.15: Post-natal health checks for mothers and newborns

Percent distribution of women age 15-49 years with a live birth in the last two years by post-natal health checks for the mother and newborn, within two days of the most recent birth, Turkana County MICS, 2013/14

	Post-natal health checks within two days of birth for:				Total	Number of women with a live birth in the last two years
	Both mothers and newborns	Mothers only	Newborns only	Neither mother nor newborn		
Total	36.1	0.9	1.0	62.0	100.0	387
Area						
Urban	56.3	0.9	1.3	41.5	100.0	199
Rural	14.7	1.0	0.6	83.6	100.0	188
Mother's age at birth						
Less than 20	(52.8)	(0.0)	(1.4)	(45.9)	100.0	52
20-34	35.6	1.1	0.7	62.6	100.0	277
35-49	23.9	0.9	2.1	73.2	100.0	59
Place of delivery						
Home	5.4	0.4	1.3	92.9	100.0	247
Health facility	93.8	2.0	0.0	4.2	100.0	135
Education						
None	26.8	0.4	0.7	72.1	100.0	268
Primary	53.8	0.0	0.9	45.3	100.0	82
Secondary+	(63.4)	(7.2)	(3.2)	(26.2)	100.0	38
Wealth index quintile						
Poorest	12.8	1.4	0.0	85.8	100.0	69
Second	5.7	0.0	1.4	93.0	100.0	86
Middle	39.4	0.0	0.0	60.6	100.0	84
Fourth	57.2	0.0	3.1	39.7	100.0	85
Richest	70.4	4.3	0.0	25.3	100.0	63
Ethnicity of household head						
Turkana	28.0	0.6	1.2	70.2	100.0	327
Other ethnic group	81.5	2.9	0.0	15.6	100.0	59

⁸⁶ Information on PNC visits for mothers by location and type of provider is not included due to the small number of cases reported.

Figure RH.3: Place of delivery and post-natal health checks, Turkana, 2013/14



8. Early Childhood Development

This chapter focuses on early childhood care and development, quality of care, child support for learning in the home, learning materials available for child use such as reading books and toys, and the developmental status of children under-5 years of age.

8.1 Early Childhood Care and Education

Readiness of children for primary school can be improved through attendance to early childhood education programmes or through pre-school attendance. Early childhood education programmes include programmes for children that have organised learning components as opposed to baby-sitting and day-care which do not typically have organised education and learning.

The Government of Kenya recognizes the importance of early childhood development (ECD) for attainment of Education for All (EFA) and the Millennium Development Goals (MDGs). The first goal of EFA obligates governments to expand early childhood care. In particular, the Government has demonstrated concern for improving the well-being of young children by enacting the Children's Act, 2001, which has managed to amalgamate all the laws of children into one document. The Act is now a legal instrument that not only protects children, but also advocates for them. Furthermore, the Government of Kenya developed Early Childhood Development Service Standard Guidelines and a National Early Childhood Development Policy Framework in 2006 which provide ECD standards, a co-ordination mechanism, and explicitly define the roles of parents, communities, various Government ministries and departments, development partners, and other stakeholders in the provision of ECD services in the country.^{87, 88}

Table CD.1 presents the results on children age 36-59 months who are attending an organized early childhood education programme in Turkana County. About 31 percent of children age 36-59 months are attending an organised early childhood education programme. Thirty-three percent of children who are attending an early childhood education programme are boys compared with 29 percent being girls. The proportions of children attending early childhood education programmes at ages 36-47 months and 48-59 months are 20 percent and 42 percent, respectively. Thirty-four percent of children from the richest households are attending school compared with 24 percent of children from the poorest households.

⁸⁷ Government of Kenya. 2006. National Early Childhood Development Policy Framework 2006

⁸⁸ Government of Kenya. 2006. Early Childhood Development Service Standard Guidelines for Kenya 2006.

Table CD.1: Early childhood education		
Percentage of children age 36-59 months who are attending an organized early childhood education programme, Turkana County MICS, 2013/14		
	Percentage of children age 36-59 months attending early childhood education ¹	Number of children age 36-59 months
Total	30.8	440
Sex		
Male	32.5	218
Female	29.2	221
Area		
Urban	29.5	225
Rural	32.2	214
Age of child		
36-47 months	20.0	222
48-59 months	41.9	217
Mother's education		
None	28.8	316
Primary	38.7	77
Secondary+	(29.9)	45
Wealth index quintile		
Poorest	24.3	91
Second	33.9	101
Middle	33.5	83
Fourth	29.3	96
Richest	34.0	68
¹ MICS indicator 6.1 - Attendance to early childhood education		
() Figures that are based on 25-49 unweighted cases		

8.2 Quality of Care

It is well recognized that a period of rapid brain development occurs in the first 3-4 years of life, and the quality of home care is a major determinant of the child's development during this period.⁸⁹ In this context, engagement of adults in activities with children, presence of books in the home for the child, and the conditions of care are important indicators of quality of home care. As set out in 'A World Fit for Children', "children should be physically healthy, mentally alert, emotionally secure, socially competent and ready to learn."⁹⁰

Information on a number of activities that support early learning was collected in the survey. These included the involvement of adults with children in the following activities: reading books or looking at picture books, telling stories, singing songs, taking children outside the home, compound or yard, playing with children, and spending time with children naming, counting, or drawing things.

⁸⁹ Grantham-McGregor, S et al. 2007. *Developmental Potential in the First 5 Years for Children in Developing Countries*. The Lancet 369: 60-70

Belsky, J et al. 2006. *Socioeconomic Risk, Parenting During the Preschool Years and Child Health Age 6 Years*. European Journal of Public Health 17(5): 511-2.

⁹⁰ UNICEF. 2002. *A World Fit For Children* adopted by the UN General Assembly at the 27th Special Session, 10 May 2002: 2.

Table CD.2 shows that fifty-three percent of children age 36-59 months have an adult household member engaged in four or more activities that promote learning and school readiness during the three days preceding the survey. The mean number of activities that adults engage with children is 3. The father's involvement in such activities was very limited since only one percent of children age 36-59 months have fathers who are involved in four or more activities, with less than one activity as the mean number of activities they are involved in. Mother's engagement in four or more activities that promote learning during the three days preceding the survey is also very limited at three percent, with a mean number of activities at 1.

There are minor differentials by sex in terms of engagement of adults, fathers or mothers in activities with children. A higher proportion of adults engaged in learning and school readiness activities among children living in urban areas (61 percent) than in rural areas (45 percent).

Table CD.2: Support for learning

Percentage of children age 36-59 months with whom adult household members engaged in activities that promote learning and school readiness during the last three days, and engagement in such activities by biological fathers and mothers, Turkana County MICS, 2013/14

	Percentage of children with whom adult household members have engaged in four or more activities ¹	Mean number of activities with adult household members	Percentage of children living with their:		Number of children age 36-59 months	Percentage of children with whom biological fathers have engaged in four or more activities ²	Mean number of activities with biological fathers	Number of children age 36-59 months living with their biological fathers	Percentage of children with whom biological mothers have engaged in four or more activities ³	Mean number of activities with biological mothers	Number of children age 36-59 months living with their biological mothers
			Biological father	Biological mother							
Total	52.9	3.3	64.8	89.2	440	0.8	0.2	285	3.1	0.8	392
Sex											
Male	51.2	3.3	65.6	91.8	218	0.3	0.3	143	5.5	0.8	200
Female	54.5	3.4	64.0	86.6	221	1.3	0.1	142	0.7	0.7	192
Area											
Urban	60.7	3.7	53.8	87.9	225	0.3	0.2	121	4.9	0.9	198
Rural	44.7	3.0	76.4	90.5	214	1.3	0.2	164	1.2	0.6	194
Age											
36-47 months	50.2	3.2	68.7	92.6	222	0.6	0.2	153	3.8	0.8	206
48-59 months	55.6	3.4	60.8	85.7	217	1.0	0.2	132	2.4	0.7	186
Mother's education^a											
None	47.8	3.1	65.3	87.5	316	1.1	0.2	206	1.5	0.6	277
Primary	68.2	4.0	56.3	94.4	77	(0.0)	(0.1)	43	5.3	0.9	73
Secondary+	(61.4)	(4.1)	(74.8)	(91.8)	45	(0.0)	(0.4)	34	(10.4)	(1.4)	41
Father's education											
None	46.7	3.1	100.0	98.3	174	0.8	0.2	174	1.2	0.6	171
Primary	(50.6)	(3.2)	(100.0)	(100.0)	45	(0.0)	(0.2)	45	(1.0)	(0.8)	45
Secondary	(70.0)	(4.0)	(100.0)	(100.0)	58	(3.8)	(0.5)	58	(10.1)	(1.3)	58
Father not in the household	51.7	3.3	0.0	71.2	155	na	na	na	3.4	0.7	110
Wealth index quintile											

Poorest	47.7	3.1	74.5	86.2	91	0.0	0.1	68	1.0	0.7	79
Second	46.6	2.9	62.8	90.5	101	0.0	0.1	63	3.0	0.6	91
Middle	46.9	3.2	49.9	86.7	83	(3.4)	(0.2)	41	0.6	0.8	72
Fourth	61.8	3.6	65.6	94.4	96	0.8	0.2	63	2.4	0.8	91
Richest	63.7	4.1	71.7	86.9	68	(0.0)	(0.3)	49	(10.1)	(1.1)	60
Ethnicity of household head											
Turkana	50.8	3.2	67.4	87.9	370	0.4	0.2	250	2.2	0.7	325
Other ethnic group	63.9	3.9	50.8	95.8	70	(3.1)	(0.4)	35	8.0	1.3	67
¹ MICS indicator 6.2 - Support for learning											
² MICS Indicator 6.3 - Father's support for learning											
³ MICS Indicator 6.4 - Mother's support for learning											
na: not applicable											
^a The background characteristic "Mother's education" refers to the education level of the respondent to the Questionnaire for Children Under Five, and covers both mothers and primary caretakers, who are interviewed when the mother is not listed in the same household. Since indicator 6.4 reports on the biological mother's support for learning, this background characteristic refers to only the educational levels of biological mothers when calculated for the indicator in question.											
() Figures that are based on 25-49 unweighted cases											

Exposure to books in early years not only provides the child with greater understanding of the nature of print, but may also give the child opportunities to see others reading, such as older siblings doing school work. Presence of books is important for later school performance. Mothers/caretakers of all children under-five years were asked about the number of children's books or picture books they have for the child, and the types of playthings that are available at home. In Turkana County, Table CD.3 shows that availability of children's books for those age 0-59 months was very minimal.

The types of playthings included contribute to the development of a child. The types of playthings included in the questionnaire were homemade toys (such as dolls and cars, or other toys made at home), toys that came from a store, and household objects (such as pots and bowls) or objects and materials found outside the home (such as sticks, rocks, animal shells, or leaves). Nineteen percent of children age 0-59 months had two or more types of playthings to play with in their homes.

Forty-six percent of the children age 0-59 play with household objects or objects found outside, 20 percent play with homemade toys, and 10 percent of children play with toys that come from a store.

The proportion of children who have two or more types of playthings to play with is 24 percent for urban children and 15 percent for rural children. The percentage of children who have two or more types of playthings to play with increases with the child's age; 12 percent for children age 0-23 months, and 25 percent for children age 24-59 months. The same pattern is observed for mother's education in which the proportion of children with two or more things to play with increases with an increase in mother's level of education.

Table CD.3: Learning materials

Percentage of children under age 5 by numbers of children's books present in the household, and by playthings that child plays with, Turkana County MICS, 2013/14

	Percentage of children living in households that have for the child:		Percentage of children who play with:				Number of children under age 5
	3 or more children's books ¹	10 or more children's books	Homemade toys	Toys from a shop/manufactured toys	Household objects/objects found outside	Two or more types of playthings ²	
Total	0.3	0.0	20.2	10.1	46.3	19.4	1,067
Sex							
Male	0.2	0.0	19.8	11.1	48.0	19.6	537
Female	0.5	0.0	20.6	9.1	44.5	19.3	530
Area							
Urban	0.7	0.0	23.5	15.8	51.7	24.1	546
Rural	0.0	0.0	16.7	4.1	40.5	14.6	521
Age							
0-23 months	0.0	0.0	13.9	7.2	27.4	11.7	422
24-59 months	0.6	0.0	24.3	12.0	58.6	24.5	645
Mother's education							
None	0.3	0.0	16.6	4.9	42.7	15.2	758
Primary	0.0	0.0	31.3	17.4	52.8	28.9	207
Secondary+	1.1	0.0	24.7	34.6	60.8	33.0	99
Wealth index quintile							
Poorest	0.0	0.0	16.3	1.9	38.0	14.3	216
Second	0.0	0.0	14.3	2.6	37.7	12.5	244
Middle	0.0	0.0	18.2	7.9	45.7	17.7	217
Fourth	0.0	0.0	22.9	11.8	51.3	21.8	227
Richest	2.2	0.0	33.2	32.7	63.7	35.7	163
Ethnicity of household head							
Turkana	0.3	0.0	19.5	8.7	45.8	18.6	898
Other ethnic group	0.7	0.0	23.4	17.6	48.5	23.6	167
¹ MICS indicator 6.5 - Availability of children's books							
² MICS indicator 6.6 - Availability of playthings							

Leaving children alone or in the presence of other young children is known to increase the risk of injuries.⁹¹ In Turkana County MICS, two questions were asked to find out whether children age 0-59 months were left alone during the week preceding the interview, and whether children were left in the care of other children under 10 years of age.

Table CD.4 shows that 53 percent of children age 0-59 months are left in the care of other children, while 26 percent are left alone. Combining the two care indicators, it is estimated that a total of 54 percent of children are left with inadequate care, either by being left alone or in the care of another child. Differences are observed by age of the child. More children age 24-59 months are left with inadequate care (72 percent) than those who are of age 0-23 months (28 percent).

⁹¹ Grossman, DC. 2000. The History of Injury Control and the Epidemiology of Child and Adolescent Injuries. The Future of Children, 10(1): 23-52.

Table CD.4: Inadequate care				
Percentage of children under age 5 left alone or left in the care of another child younger than 10 years of age for more than one hour at least once during the past week, Turkana County MICS, 2013/14				
	Percentage of children under age 5:			Number of children under age 5
	Left alone in the past week	Left in the care of another child younger than 10 years of age in the past week	Left with inadequate care in the past week ¹	
Total	25.6	52.6	54.1	1,067
Sex				
Male	27.4	52.8	54.2	537
Female	23.7	52.4	54.0	530
Area				
Urban	21.9	54.3	55.7	546
Rural	29.4	50.8	52.5	521
Age				
0-23 months	8.4	26.2	27.5	422
24-59 months	36.8	70.0	71.6	645
Mother's education				
None	26.5	50.8	51.9	758
Primary	23.5	53.3	54.8	207
Secondary+	22.4	65.6	69.8	99
Wealth index quintile				
Poorest	28.1	51.3	53.2	216
Second	31.5	50.4	52.2	244
Middle	26.0	52.5	53.0	217
Fourth	20.9	50.2	51.6	227
Richest	19.2	61.4	63.3	163
Ethnicity of household head				
Turkana	28.2	54.6	56.0	898
Other ethnic group	11.5	42.1	44.0	167
¹ MICS indicator 6.7 - Inadequate care				

8.3 Developmental Status of Children

Early childhood development is defined as an orderly, predictable process along a continuous path, in which a child learns to handle more complicated levels of moving, thinking, speaking, feeling and relating to others. Physical growth, literacy and numeracy skills, socio-emotional development and readiness to learn are vital domains of a child's overall development, which is a basis for overall human development.⁹²

A 10-item module was used to calculate the Early Child Development Index (ECDI). The primary purpose of the ECDI is to inform public policy regarding the developmental status of children in Turkana County. The index is based on selected milestones that children are expected to achieve by

⁹² Shonkoff, J and Phillips, D (eds). 2000. *From neurons to neighborhoods: the science of early childhood development*. Committee on Integrating the Science of Early Childhood Development, National Research Council, 2000.

ages 3 and 4. The 10 items are used to determine if children are developmentally on track in four domains:

Literacy-numeracy: Children are identified as being developmentally on track based on whether they can identify/name at least ten letters of the alphabet, whether they can read at least four simple, popular words, and whether they know the name and recognize the symbols of all numbers from 1 to 10. If at least two of these are true, then the child is considered developmentally on track.

Physical: If the child can pick up a small object with two fingers, like a stick or a rock from the ground and/or the mother/caretaker does not indicate that the child is sometimes too sick to play, then the child is regarded as being developmentally on track in the physical domain.

Social-emotional: Children are considered to be developmentally on track if two of the following are true: If the child gets along well with other children, if the child does not kick, bite, or hit other children and if the child does not get distracted easily.

Learning: If the child follows simple directions on how to do something correctly and/or when given something to do, is able to do it independently, then the child is considered to be developmentally on track in this domain.

ECDI is then calculated as the percentage of children who are developmentally on track in at least three of these four domains. The results are presented in Table CD.5.

In Turkana County, half of children age 36-59 months were developmentally on track. ECDI is slightly higher among boys (56 percent) than among girls (44 percent). The ECDI is 54 percent among children age 48-59 months and 47 percent among those age 36-47 months, since children develop more skills with increasing age. A higher ECDI is seen in children attending to an early childhood education programme at 64 percent compared with 44 percent among those who are not attending.

The analysis of four domains of child development shows that 82 percent of children are on track in the physical domain, 71 percent in the learning domain, 69 percent in the social-emotional domain, but much less on track in literacy-numeracy domain (12 percent). In each individual domain (except the physical), the higher score is associated with children attending an early childhood education programme and in older children (except the social-emotional).

Table CD.5: Early child development index

Percentage of children age 36-59 months who are developmentally on track in literacy-numeracy, physical, social-emotional, and learning domains, and the early child development index score, Turkana County MICS, 2013/14

	Percentage of children age 36-59 months who are developmentally on track for indicated domains				Early child development index score ¹	Number of children age 36-59 months
	Literacy-numeracy	Physical	Social-Emotional	Learning		
Total	11.9	82.1	69.0	71.2	50.1	440
Sex						
Male	13.3	82.6	69.9	75.6	56.2	218
Female	10.6	81.7	68.1	66.8	44.1	221
Area						
Urban	16.7	80.2	70.0	77.4	56.8	225
Rural	7.0	84.1	67.9	64.7	43.1	214
Age						
36-47 months	5.8	79.8	71.3	66.0	46.5	222
48-59 months	18.2	84.5	66.6	76.5	53.8	217
Attendance to early childhood education						
Attending	32.5	80.3	72.9	81.4	64.4	136
Not attending	2.8	82.9	67.2	66.6	43.7	304
Mother's education						
None	6.3	82.0	66.5	66.8	44.9	316
Primary	28.8	81.1	76.7	81.0	63.4	77
Secondary+	(22.9)	(83.9)	(72.1)	(84.0)	(62.5)	45
Wealth index quintile						
Poorest	2.1	81.9	69.0	61.8	36.2	91
Second	7.4	83.4	63.0	63.4	42.2	101
Middle	14.3	78.4	74.0	75.6	49.7	83
Fourth	14.2	87.2	74.5	75.6	63.6	96
Richest	25.7	78.0	64.0	83.4	61.7	68
Ethnicity of household head						
Turkana	11.6	81.5	66.7	69.4	46.7	370
Other ethnic group	13.6	85.2	81.3	80.6	67.9	70
¹ MICS indicator 6.8 - Early child development index						
() Figures that are based on 25-49 unweighted cases						

9. Literacy and Education

Kenya is a signatory to several critical instruments for the enhancement of the rights to quality education for its citizens. These include the Universal Declaration on Human Rights (1948); the minimum Age Convention (1973); the convention on the Elimination of all forms of Discrimination Against Women (CEDAW) of 1979; the Convention on the rights of the Child (CRC) of 1989; the International Convention on the Protection of the rights of All Migrant workers and members of their families (1990); the Dakar Framework of Action on EFA (2000); the Millennium Development Goals (MDGs) 2000; and the convention on the Rights of Persons with Disabilities (2006). According to the Constitution of Kenya, Section 43, 1f, every child has the right to education.⁹³

Chapter Nine focuses on literacy among young women, school readiness, primary and secondary school participation and gender parity.

9.1 Literacy among Young Women

The Youth Literacy Rate reflects the outcomes of primary education over the previous 10 years or so. As a measure of the effectiveness of the primary education system, it is often seen as a proxy measure of social progress and economic achievement. Since a men's questionnaire was not administered as part of the Turkana County MICS, the results are based only on females age 15-24 years. Literacy is assessed on the ability of the respondent to read a short simple statement or based on school attendance.

Table ED.1 indicates that 55 percent of young women in Turkana County are literate and that literacy status varies by area (65 percent in urban and 24 percent in rural areas). Among the young women who stated that primary school is their highest level of education, 71 percent are able to read the statement shown to them. Variations are also noted by age and household wealth status. Young women age 15-19 years (64 percent) are more likely to read than women age 20-24 years (43 percent). Women in the richest wealth index quintile (81 percent) are ten more times likely to read than those in the poorest wealth index quintile (8 percent). Women of other ethnic groups (64 percent) are more likely than Turkana women (52 percent) to read.

⁹³ The Constitution of Kenya 2010

Table ED.1: Literacy (young women)			
Percentage of women age 15-24 years who are literate, Turkana County MICS, 2013/14			
	Percentage literate ¹	Percentage not known	Number of women age 15-24 years
Total	54.7	0.4	461
Area			
Urban	64.8	0.6	345
Rural	24.4	0.0	116
Education			
None	0.5	0.0	152
Primary	70.9	1.0	196
Secondary+	100.0	0.0	112
Age			
15-19	64.1	0.8	252
20-24	43.3	0.0	209
Wealth index quintile			
Poorest	6.1	0.0	44
Second	17.7	0.0	45
Middle	48.3	0.0	100
Fourth	59.7	0.0	127
Richest	81.1	1.3	145
Ethnicity of household head			
Turkana	51.6	0.5	350
Other ethnic group	64.3	0.0	110
¹ MICS indicator 7.1; MDG indicator 2.3 - Literacy rate among young women			

9.2 School Readiness

Pre-primary school attendance is important for the readiness of children to education. Table ED.2 shows the proportion of children in the first grade of primary school (regardless of age) who attended pre-primary the previous year.⁹⁴ Overall, 44 percent of children who are currently attending the first grade of primary school attended pre-primary the previous year.

⁹⁴ The computation of the indicator does not exclude repeaters, and therefore is inclusive of both children who are attending primary school for the first time, as well as those who were in the first grade of primary school the previous school year and are repeating. Children repeating may have attended Pre-primary prior to the school year during which they attended the first grade of primary school for the first time; these children are not captured in the numerator of the indicator.

Table ED.2: School readiness		
Percentage of children attending first grade of primary school who attended Pre-primary the previous year, Turkana County MICS, 2013/14		
	Percentage of children attending first grade who attended Pre-primary in previous year ¹	Number of children attending first grade of primary school
Total	43.8	185
Sex		
Male	40.9	103
Female	47.4	82
Area		
Urban	33.0	106
Rural	58.1	79
¹ MICS indicator 7.2 - School readiness		

9.3 Primary and Secondary School Participation

Achievement of universal primary education and by the world's children was one of the Millennium Development Goals. Education is a vital prerequisite for combating poverty, empowering women, protecting children from hazardous and exploitative labour and sexual exploitation, promoting human rights and democracy, protecting the environment, and influencing population growth.

In Kenya, the structure of Early Childhood Development and Education (ECDE) provision is divided into two parts: 0-2 and 3-5 year-old children. Children are expected to enter primary school at age 6 and secondary school at age 14. Primary school has 8 grades (1-8) and secondary school comprises 4 grades (1-4). In primary school level of education, grades are referred to as Standard 1 to Standard 8 after which a Kenya Certificate of Primary Education (KCPE) is attained after sitting examinations. For secondary school level of education, grades are referred to as Form 1 to Form 4, and a Kenya Certificate of Secondary Education (KCSE) is attained after successful completion of the full cycle and sitting examinations. The school year typically runs from January to November.⁹⁵

In Turkana County, 34 percent of children who are of primary school entry age (age 6 years) are attending the first grade of primary school (Table ED.3). The proportion of 6-year old children entering grade 1 is similar for boys (36 percent) and girls (30 percent), and is higher for children in urban areas (38 percent) compared to those in rural areas (28 percent).

⁹⁵ Ministry of Education Science and Technology, 2005. Kenya Education Sector Support Programme 2005-2010.

Table ED.3: Primary school entry		
Percentage of children of primary school entry age entering grade 1 (net intake rate), Turkana County MICS, 2013/14		
	Percentage of children of primary school entry age entering grade 1 ¹	Number of children of primary school entry age
Total	33.6	202
Sex		
Male	36.2	112
Female	30.3	90
Area		
Urban	38.4	105
Rural	28.4	97
¹ MICS indicator 7.3 - Net intake rate in primary education		

Table ED.4 provides the percentage of children of primary school age 6 to 13 years who are attending primary or secondary school⁹⁶ and those who are out of school. Sixty-five percent of children of primary school age are attending school while 33 percent are out of school. Net attendance ratio is 82 percent in urban areas and 47 percent in rural areas. Net attendance increases with age at beginning of school year, mother's level of education, wealth index quintiles and ethnicity. Net attendance ratio to primary school is 89 percent for children whose mothers have secondary or higher education and 58 percent for children whose mothers have no education. Net attendance ratio increases with an increase in household wealth ranging between 28 percent for the poorest households to 92 percent in the richest.

⁹⁶ Ratios presented in this table are "adjusted" since they include not only primary school attendance, but also secondary school attendance in the numerator.

Table ED.4: Primary school attendance and out of school children

Percentage of children of primary school age attending primary or secondary school (adjusted net attendance ratio), percentage attending Pre-primary, and percentage out of school, Turkana County MICS, 2013/14

	Male					Female					Total				
	Percentage of children:					Percentage of children:					Percentage of children:				
	Net attendance ratio (adjusted)	Not attending school or Pre-primary	Attending Pre-primary	Out of school ^a	Number of children	Net attendance ratio (adjusted)	Not attending school or Pre-primary	Attending Pre-primary	Out of school ^a	Number of children	Net attendance ratio (adjusted) ¹	Not attending school or Pre-primary	Attending Pre-primary	Out of school ^a	Number of children
Total	66.2	25.3	6.8	32.0	820	64.5	27.2	6.9	34.1	786	65.4	26.2	6.8	33.0	1,606
Area															
Urban	82.1	7.7	8.6	16.4	437	82.1	7.2	9.5	16.7	407	82.1	7.5	9.1	16.5	844
Rural	48.2	45.2	4.6	49.9	383	45.5	48.8	4.0	52.8	379	46.9	47.0	4.3	51.3	762
Age at beginning of school year															
6	46.5	26.5	24.9	51.4	112	38.7	28.2	29.6	57.8	90	43.1	27.3	27.0	54.3	202
7	46.7	32.4	15.1	47.4	115	60.6	22.3	16.0	38.4	117	53.7	27.3	15.6	42.8	232
8	65.5	28.0	5.0	33.0	104	62.3	29.8	4.0	33.9	76	64.1	28.8	4.6	33.4	180
9	70.6	26.1	3.4	29.4	110	69.4	29.8	0.8	30.6	109	70.0	27.9	2.1	30.0	219
10	78.4	21.0	0.6	21.6	83	71.3	26.5	2.2	28.7	85	74.8	23.8	1.4	25.2	169
11	80.2	18.7	1.1	19.8	89	65.3	28.4	3.1	31.5	84	73.0	23.4	2.1	25.5	173
12	74.5	23.1	0.0	23.1	115	72.4	26.6	0.0	26.6	116	73.4	24.8	0.0	24.8	230
13	75.5	23.7	0.0	23.7	92	72.5	27.5	0.0	27.5	108	73.9	25.8	0.0	25.8	200
Mother's education															
None	59.5	32.0	7.2	39.1	608	56.4	36.0	6.7	42.8	565	58.0	33.9	6.9	40.9	1,172
Primary	85.2	7.0	5.4	12.4	128	86.5	3.8	8.3	12.2	126	85.8	5.5	6.8	12.3	255
Secondary	89.8	3.6	6.6	10.2	78	89.0	2.3	7.4	9.6	76	89.4	2.9	7.0	9.9	154
Cannot be determined ^b	(*)	(*)	(*)	(*)	4	(*)	(*)	(*)	(*)	17	(*)	(*)	(*)	(*)	21
Wealth index quintile															
Poorest	30.3	61.6	7.7	69.3	168	26.0	67.0	6.6	73.6	171	28.1	64.3	7.1	71.5	338
Second	52.8	36.9	7.2	44.1	179	47.7	44.2	5.7	49.9	155	50.4	40.3	6.5	46.8	335
Middle	76.4	13.6	6.4	20.0	175	74.4	14.5	7.5	22.0	154	75.5	14.0	6.9	20.9	329
Fourth	85.9	7.6	6.5	14.1	166	87.8	4.3	7.9	12.2	161	86.8	6.0	7.2	13.2	327

Richest	92.0	0.8	5.9	6.7	132	91.5	1.2	6.6	7.8	145	91.7	1.0	6.3	7.3	277
Ethnicity of household head															
Turkana	62.8	28.2	7.2	35.5	688	60.4	31.1	6.9	38.0	664	61.6	29.6	7.0	36.7	1352
Other ethnic group	84.3	9.3	4.5	13.9	128	87.0	6.0	7.1	13.0	121	85.6	7.7	5.8	13.5	249

¹ 7.S1 - Primary school net attendance ratio (adjusted)

^a The percentage of children of primary school age out of school are those not attending school and those attending Pre-primary

^b Children age 15 or higher at the time of the interview whose mothers were not living in the household

(*) Figures that are based on fewer than 25 unweighted cases

The secondary school net attendance ratio is presented in Table ED.5.⁹⁷ About 19 percent of the children of secondary school age are attending secondary school, 51 percent are attending primary school and 28 percent are out of school. The secondary net attendance ratio is similar for males and females (19 percent for females and 20 percent for males). In urban areas, 24 percent of children of secondary school age are attending secondary school, while 10 percent of them are attending in rural areas. The proportion of secondary school age children out of school is higher in rural areas (54 percent) compared to 15 percent for urban areas.

⁹⁷ Ratios presented in this table are "adjusted" since they include not only secondary school attendance, but also attendance to higher levels in the numerator.

Table ED.5: Secondary school attendance and out of school children

Percentage of children of secondary school age attending secondary school or higher (adjusted net attendance ratio), percentage attending primary school, and percentage out of school, Turkana County MICS, 2013/14

	Male				Female				Total			
	Percentage of children:				Percentage of children:				Percentage of children:			
	Net attendance ratio (adjusted)	Attending primary school	Out of school ^a	Number of children	Net attendance ratio (adjusted)	Attending primary school	Out of school ^a	Number of children	Net attendance ratio (adjusted) ¹	Attending primary school	Out of school ^a	Number of children
Total	19.6	53.5	25.4	347	19.1	46.9	31.7	299	19.4	50.5	28.3	646
Area												
Urban	25.3	64.2	9.0	217	23.2	51.7	21.9	216	24.2	58.0	15.4	433
Rural	10.2	35.6	52.8	130	8.4	34.6	56.9	83	9.5	35.2	54.4	213
Age at beginning of school year												
14	2.9	72.9	20.6	99	7.6	59.1	31.7	74	4.9	67.0	25.3	174
15	15.1	52.8	32.0	77	10.3	60.0	29.7	75	12.8	56.4	30.9	153
16	33.1	45.7	21.2	81	31.6	36.6	31.8	77	32.4	41.3	26.3	157
17	29.9	39.4	28.8	90	26.6	32.0	33.6	73	28.4	36.1	30.9	162
Wealth index quintile												
Poorest	5.0	28.4	66.6	56	(4.1)	(11.8)	(84.1)	32	4.7	22.3	73.0	88
Second	9.9	45.2	41.6	58	(6.0)	(27.3)	(66.7)	27	8.6	39.5	49.7	85
Middle	19.1	61.1	18.4	70	12.7	49.9	33.7	67	16.0	55.6	25.9	137
Fourth	21.6	66.9	11.4	78	9.4	68.6	18.5	77	15.6	67.8	15.0	154
Richest	34.3	57.0	5.9	86	40.0	44.9	13.1	96	37.3	50.6	9.7	182
Ethnicity of household head												
Turkana	20.7	48.2	29.6	271	20.8	40.5	36.2	233	20.7	44.6	32.7	504
Other ethnic group	16.6	71.1	11.0	73	12.8	69.8	15.6	66	14.8	70.5	13.2	140

¹ 7.S2 - Secondary school net attendance ratio (adjusted)

^a The percentage of children of secondary school age out of school are those who are not attending primary, secondary, or higher education

() Figures that are based on 25-49 unweighted cases

The MICS included only questions on school attendance in the current and previous year. Thus, the indicator is calculated by computing the cumulative probability of survival from the first to the last grade of primary school, as opposed to calculating the indicator for a real cohort that needs to be followed from the time a cohort of children entered primary school, up to the time they reached the last grade of primary school. Repeaters are excluded from the calculation of the indicator, because it is not known whether they will eventually graduate. As an example, the probability that a child will move from the first grade to the second grade is computed by dividing the number of children who moved from the first grade to the second grade (during the two consecutive school years covered by the survey) by the number of children who have moved from the first to the second grade plus the number of children who were in the first grade the previous school year, but dropped out. Both the numerator and denominator exclude children who repeated during the two school years under consideration.

Table ED6 shows the percentage of children entering first grade who eventually reach last grade of primary school. The majority of the children starting grade 1 reach grade 8 (96 percent). There are no disparities by sex and place of residence.

Table ED.6: Children reaching last grade of primary school

Percentage of children entering first grade of primary school who eventually reach the last grade of primary school (Survival rate to last grade of primary school), Turkana County MICS, 2013/14

	Percent attending grade 1 last school year who are in grade 2 this school year	Percent attending grade 2 last school year who are attending grade 3 this school year	Percent attending grade 3 last school year who are attending grade 4 this school year	Percent attending grade 4 last school year who are attending grade 5 this school year	Percent attending grade 5 last school year who are attending grade 6 this school year	Percent attending grade 6 last school year who are attending grade 7 this school year	Percent attending grade 7 last school year who are attending grade 8 this school year	Percent who reach grade 8 of those who enter grade 1 ¹
Total	100.0	99.8	99.4	100.0	99.2	99.6	98.0	96.0
Sex								
Male	100.0	99.5	98.9	100.0	100.0	100.0	100.0	98.4
Female	100.0	100.0	100.0	100.0	98.2	99.1	95.7	93.1
Area								
Urban	100.0	100.0	99.6	100.0	100.0	100.0	98.3	97.9
Rural	100.0	99.5	98.9	(100.0)	97.9	(97.8)	(96.9)	91.3
¹ 7.S3 - Children reaching last grade of primary								
() Figures that are based on 25-49 unweighted cases								

The primary school completion rate and transition rate to secondary education are presented in Table ED.7. The primary completion rate is the ratio of the total number of students, regardless of age, entering the last grade of primary school for the first time, to the number of children of the primary graduation age at the beginning of the current (or most recent) school year.

Table ED.7 shows that the primary school completion rate in Turkana is 74 percent. The completion rates differ by sex and place of residence. About 90 percent of male children compared to 60 percent of the females completed primary school. Similarly, more children in urban areas (97 percent) completed primary school compared to 39 percent for rural areas.⁹⁸

Table ED.7: Primary school completion rate		
Primary school completion rates and transition and effective transition rates to secondary school, Turkana County MICS, 2013/14		
	Primary school completion rate ¹	Number of children of primary school completion age
Total	74.0	200
Sex		
Male	90.3	92
Female	60.3	108
Area		
Urban	97.1	120
Rural	39.2	79
¹ 7.S4 - Primary completion rate		

The ratio of girls to boys attending primary and secondary education is provided in Table ED.8. These ratios are better known as the Gender Parity Index (GPI).⁹⁹ Notice that the ratios included here are obtained from net attendance ratios rather than gross attendance ratios. The latter provide an erroneous description of the GPI mainly because, in most cases, the majority of over-age children attending primary education tend to be boys.

The table shows that the primary school adjusted net attendance ratio (NAR) for girls is 65 compared to 66 for boys, giving a gender parity index of 0.97 for primary school. A similarly, secondary school adjusted net attendance ratio for boys and girls is similar (19 and 20 percent, respectively) yielding a GPI for secondary school adjusted NAR of 0.97 GPI. Both the primary school and secondary school GPI suggest that boys and girls attend primary and secondary education at the same rate.

⁹⁸ Other background characteristics variables and, both, the transition rate to secondary school and the effective transition rate to secondary school for Turkana County could not be presented in Table ED.7 due to small number of cases reported.

⁹⁹ UNESCO, 2015. EFA Monitoring Report 2015 -Education for All 2000-2015: Achievements and Challenges. Gender parity index (GPI) - Ratio of female to male values of a given indicator. A GPI between 0.97 and 1.03 indicates parity between the genders. A GPI below 0.97 indicates a disparity in favour of males. A GPI above 1.03 indicates a disparity in favour of females.

Table ED.8: Education gender parity						
Ratio of adjusted net attendance ratios of girls to boys, in primary and secondary school, Turkana County MICS, 2013/14						
	Primary school			Secondary school		
	Primary school adjusted net attendance ratio (NAR), girls	Primary school adjusted net attendance ratio (NAR), boys	Gender parity index (GPI) for primary school adjusted NAR ¹	Secondary school adjusted net attendance ratio (NAR), girls	Secondary school adjusted net attendance ratio (NAR), boys	Gender parity index (GPI) for secondary school adjusted NAR ²
Total	64.5	66.2	0.97	19.1	19.6	0.97
Area						
Urban	82.1	82.1	1.00	23.2	25.3	0.92
Rural	45.5	48.2	0.94	8.4	10.2	0.83
Mother's education						
None	56.4	59.5	0.95	4.9	9.8	0.50
Primary	86.5	85.2	1.02	(8.7)	(*)	0.40
Secondary+	89.0	89.8	0.99	(*)	(*)	1.42
Cannot be determined ^a	(*)	(*)	1.31	23.2	24.6	0.94
Wealth index quintile						
Poorest	26.0	30.3	0.86	(4.1)	5.0	0.83
Second	47.7	52.8	0.90	(6.0)	9.9	0.61
Middle	74.4	76.4	0.97	12.7	19.1	0.66
Fourth	87.8	85.9	1.02	9.4	21.6	0.44
Richest	91.5	92.0	0.99	40.0	34.3	1.17
Ethnicity of household head						
Turkana	60.4	62.8	0.96	20.8	20.7	1.01
Other ethnic group	87.0	84.3	1.03	12.8	16.6	0.78
¹ 7.S5 - Gender parity index (primary school)						
² 7.S6 - Gender parity index (secondary school)						
^a Children age 15 or higher at the time of the interview whose mothers were not living in the household						
na: not applicable						
() Figures that are based on 25-49 unweighted cases						
(*) Figures that are based on fewer than 25 unweighted cases						

The percentage of girls in the total out of school population, in both primary and secondary school, are provided in Table ED.9. The table shows that 33 percent of girls at primary level of education and 28 percent at secondary school level are out of school. Similarly, at primary level, girls accounted for 51 percent of the out-of-school population and 52 percent at secondary level.


Table ED.9: Out of school gender parity

Percentage of girls in the total out of school population, in primary and secondary school, Turkana County MICS, 2013/14

	Primary school				Secondary school			
	Percentage of out of school children	Number of children of primary school age	Percentage of girls in the total out of school population of primary school age	Number of children of primary school age out of school	Percentage of out of school children	Number of children of secondary school age	Percentage of girls in the total out of school population of secondary school age	Number of children of secondary school age out of school
Total	33.0	1,606	50.5	531	28.3	646	51.7	183
Area								
Urban	16.5	844	48.8	140	15.4	433	70.8	67
Rural	51.3	762	51.2	391	54.4	213	40.8	116
Mother's education								
None	40.9	1172	50.4	479	38.9	204	44.5	79
Primary	12.3	255	(49.0)	31	(15.3)	59	(*)	9
Secondary	9.9	154	(*)	15	(*)	29	-	0
Cannot be determined ^a	(*)	21	(*)	5	26.7	354	55.8	94
Wealth index quintile								
Poorest	71.5	338	52.0	242	73.0	88	41.9	64
Second	46.8	335	49.5	157	49.7	85	(43.3)	42
Middle	20.9	329	49.1	69	25.9	137	(63.7)	35
Fourth	13.2	327	(45.7)	43	15.0	154	(*)	23
Richest	7.3	277	(*)	20	9.7	182	(*)	18
Ethnicity of household head								
Turkana	36.7	1,352	50.8	496	32.7	504	51.3	164
Other ethnic group	13.5	249	(47.0)	33	13.2	140	(*)	18
^a Children age 15 or higher at the time of the interview whose mothers were not living in the household								
na: not applicable								
() Figures that are based on 25-49 unweighted cases								
(*) Figures that are based on fewer than 25 unweighted cases								

Figure ED.1 brings together all of the attendance and progression related education indicators covered in this chapter, by sex. Information on attendance to early childhood education is also included, which is covered in Chapter 8, in Table CD.1.

Figure ED.1: Education indicators by sex (national system), Turkana County MICS, 2013/14



All indicator values are in per cent and are calculated based on the national education system
 (*) Figures that are based on less than 25 unweighted cases

UNESCO developed the International Standard Classification of Education (ISCED) to facilitate comparisons of education statistics and indicators across countries on the basis of uniform and internationally agreed definitions.^{100, 101} The mapping of the Kenyan education system to the ISCED classification is as follows:

- (i) ISCED Level 1 is Primary Education and corresponds to Primary grades Standard 1 to 6 in the Kenyan education system.
- (ii) ISCED Level 2 is Lower Secondary Education and corresponds to Primary grades Standard 7 and 8, and Secondary grades Form 1 and 2, in the Kenyan education system.
- (iii) ISCED Level 3 is Upper Secondary Education and corresponds to Secondary grades Form 3 and 4 in the Kenyan education system.

Table ED.10 ISCED shows key education indicators in Turkana County according to the mapping of the Kenya education system to the ISCED 2011 education classification. These indicators therefore are not based on the Kenya education system but rather provide international comparison of same indicators as used in different countries education systems.

About 34 percent of children of primary school entry age enter grade 1. Approximately 62 percent of children age 6-11 years are attending primary school according to the ISCED classification (i.e. Standard 1 to 6), and 37 percent of children age 12-17 are attending secondary school (ISCED levels 2 and 3). Ninety-eight percent of the children entering primary grade 1 are expected to reach grade 6 (the last grade of the ISCED 1 level), and 96 percent transition from primary (ISCED 1 level) to secondary (ISCED 2 level).

¹⁰⁰ <http://www.uis.unesco.org/Education/Pages/international-standard-classification-of-education.aspx>
¹⁰¹ <http://www.uis.unesco.org/Education/ISCEDMappings/Pages/default.aspx>

Table ED.10: Summary of education indicators (ISCED^a)

Summary of education indicators classified according to the International Standard Classification of Education (ISCED), Turkana County MICS, 2013/14

	Primary school (ISCED 1)				Transition (ISCED 1 to 2)	Secondary school (ISCED 2+3)
	Percentage of children of primary school entry age entering grade 1 ¹	Net attendance ratio (adjusted) ²	Percent who reach grade 6 of those who enter grade 1 ³	Primary school completion rate ⁴	Transition rate to secondary school ⁵	Net attendance ratio (adjusted) ⁶
Total	33.6	62.4	98.4	84.2	96.4	36.8
Sex						
Male	36.2	63.3	98.4	95.1	94.7	36.9
Female	30.3	61.3	98.2	72.7	98.3	36.8
Gender parity index (GPI) ^{7, 8}	na	0.97	na	na	na	1.00
¹ MICS indicator 7.3 - Net intake rate in primary education						
² MICS indicator 7.4; MDG indicator 2.1 - Primary school net attendance ratio (adjusted)						
³ MICS indicator 7.6; MDG indicator 2.2 - Children reaching last grade of primary						
⁴ MICS indicator 7.7 - Primary completion rate						
⁵ MICS indicator 7.8 - Transition rate to secondary school						
⁶ MICS indicator 7.5 - Secondary school net attendance ratio (adjusted)						
⁷ MICS indicator 7.9; MDG indicator 3.1 - Gender parity index (primary school)						
⁸ MICS indicator 7.10; MDG indicator 3.1 - Gender parity index (secondary school)						
^a ISCED 1 are Standards 1-6, ISCED 2 are Standards 7-8 and Forms 1-2, and ISCED 3 are Forms 3-4.						
na: not applicable						

10. Child Protection

Kenya is committed to the survival, development and protection of children as demonstrated by its ratification of international treaties and conventions that include the 1989 United Nations Convention on the Rights of the Child (CRC), the Convention on the Elimination of all forms of Discrimination Against Women (CEDAW), the International Labour Organization (ILO) conventions on Prohibition of Child Labour and Worst Forms of Child Labour [*Chapter 182*] 1999, Palermo Protocol on Trafficking in Persons, 2000 and the Millennium Development Goals 2000 (MDGs). At regional level, Kenya ratified the 1990 African Charter on the Rights and Welfare of the Child (ACRWC).

The majority of these conventions and treaties have been domesticated into the Constitution and other enacted laws and policies that include: the Registration of Births and Deaths Act [*Chapter 149*], Rev 1990; the Children's Act, 2001; the Sexual Offences Act, 2003; the Female Genital Mutilation/Cutting Policy, 2009; the Counter Trafficking in Persons Act, 2010; the Kenya Citizenship and Immigration Act, 2011; the Labour Migration Policy, 2011; and Prohibition of Female Genital Mutilation Act, 2011; among others.

This chapter discusses birth registration, child labour, child discipline, early marriage and polygyny, female genital mutilation/cutting (FGM/C), and women's attitudes towards domestic violence.

10.1 Birth Registration

A name and nationality is every child's right, enshrined in the Convention on the Rights of the Child (CRC) and other international treaties. Yet the births of around one in four children under the age of five worldwide have never been recorded.¹⁰² This lack of formal recognition by the State usually means that a child is unable to obtain a birth certificate. As a result, he or she may be denied health care or education. Later in life, the lack of official identification documents can mean that a child may enter into marriage or the labour market, or be conscripted into the armed forces, before the legal age. In adulthood, birth certificates may be required to: obtain social assistance; acquire a job in the formal sector; prove the right to inherit property; vote; obtain a passport; etc. Registering children at birth is the first step in securing their recognition before the law, safeguarding their rights, and ensuring that any violation of these rights does not go unnoticed.¹⁰³

Birth registration requirements

The Births and Deaths Registration Act, which makes registration of all births and deaths occurring in Kenya compulsory has the following legal provisions:

- The occurrence of a birth must be registered within six months
- A registrar shall not register a birth after the expiry of six months without specific authority and payment of a late registration fee
- Registration of a birth within six months is called **current registration** and is done free of charge

¹⁰² UNICEF. 2014. *The State of the World's Children 2015*. UNICEF.

¹⁰³ UNICEF. 2013. *Every Child's Birth Right: Inequities and trends in birth registration*. UNICEF.

- Registration of a birth after six months is called **late registration** and attracts a penalty of Ksh 100. Besides, such registration is only done by the respective county registrar at their own discretion.

Births take place either within health facilities or at home. For births occurring in health facilities, the person-in charge of each facility is responsible for reporting occurrence of such births. While the primary responsibility of reporting occurrence of a birth at home is on the parents.

The midwife is responsible for completing a register of birth for every birth immediately after delivery. For every birth occurring at home, the area assistant chief is expected to complete a register of birth after receiving reports, within six months, of its occurrence within their respective areas of jurisdiction.

All completed registers of birth, from all health facilities and sub-locations are transmitted to respective county civil registries once every month. Upon receipt, they are checked for completeness and accuracy after which respective sub-county civil registrars append their signatures, thereby certifying them as legal documents. These legal documents are supposed to be maintained under safe custody within respective sub-county civil registries for purposes of issuance of certificates and other related documents.

While registration of births is compulsory, acquisition of a birth certificate is not. When in need, one makes an application for such a certificate in the county in which the event occurred. Sub-county civil registrars authorize issuance of certificates of birth from registers of birth under their custody upon application, production of supportive documentation and payment of subscribed fees. An applicant is required to pay Ksh 50 in order to acquire a birth certificate. In case of any amendment on the register of birth, before a birth certificate is issued, an extra Ksh 50 is levied.

The Births and Deaths Registration Act has provision for registering births outside the mandatory six months. Respective sub-county civil registrars have the sole discretion in approving applications for late registration of births. However, applications for late registration of births within border counties have to be vetted through the ranks of the local administration before they reach respective sub-county civil registrars. All applications for late registration must be supported by documents in relation to key characteristics pertaining to the occurrence of the birth such as date and place of occurrence, parentage, etc.

Birth Registration Status

The Turkana County MICS sought to provide an estimate of the extent of birth registration of children under-5 years of age. Mothers/caretakers of these children were asked whether children in their household had birth certificates. If they responded that a child did not have a birth certificate, additional questions were asked on whether the child's birth was registered and whether they knew how to register a birth. A child may not have been issued a birth certificate but the birth may have been registered.

Birth registration in this context includes:

- children whose birth certificates were seen by the interviewer;
- children reported to have a birth certificate that was not seen by the interviewer; and
- children who did not have a birth certificate but were reported to have been registered.


The births of 40 percent of children under-5 years in Turkana County are registered (Table CP.1). Variations are noted in birth registration for urban and rural children, 54 percent and 26 percent, respectively. Birth registration increases with mother's education and with household wealth. Only 14 percent showed a birth certificate to the interviewer. These findings are summarized in Figure CP.1.

Table CP.1: Birth registration							
Percentage of children under age 5 by whether birth is registered and percentage of children not registered whose mothers/caretakers know how to register birth, Turkana County MICS, 2013/14							
	Children under age 5 whose birth is registered with civil authorities				Number of children under age 5	Children under age 5 whose birth is not registered	
	Has birth certificate		No birth certificate	Total registered ¹		Percent of children whose mother/caretaker knows how to register birth	Number of children under age 5 without birth registration
	Seen	Not seen					
Total	13.7	8.5	18.0	40.1	1,067	23.8	639
Sex							
Male	12.9	9.2	17.2	39.3	537	25.4	326
Female	14.5	7.7	18.8	41.0	530	22.0	313
Area							
Urban	21.2	12.1	20.5	53.8	546	45.0	252
Rural	5.8	4.6	15.3	25.8	521	9.9	387
Age							
0-11 months	12.0	8.3	21.5	41.8	227	27.1	132
12-23 months	13.1	7.4	20.4	40.9	196	26.6	116
24-35 months	13.9	9.5	19.5	42.9	205	14.8	117
36-47 months	18.1	9.5	14.8	42.4	222	25.3	128
48-59 months	11.3	7.5	13.9	32.7	217	24.3	146
Mother's education							
None	9.6	6.2	15.6	31.4	758	15.9	520
Primary	19.5	13.3	26.3	59.1	207	58.2	85
Secondary	33.2	15.9	19.2	68.4	99	(63.8)	31
Wealth index quintile							
Poorest	3.9	2.6	15.3	21.7	216	9.8	169
Second	4.0	4.7	15.2	23.9	244	10.6	186
Middle	13.6	7.5	18.1	39.3	217	28.7	132
Fourth	19.9	13.6	20.4	53.8	227	49.8	105
Richest	32.8	16.0	22.1	70.9	163	54.2	47
Ethnicity of household head							
Turkana	9.6	7.2	17.4	34.2	898	23.3	591
Other ethnic group	35.4	15.3	21.1	71.7	167	(30.4)	47
¹ MICS indicator 8.1 - Birth registration							
() Figures that are based on 25-49 unweighted cases							

The lack of adequate knowledge of how to register a birth can present another major obstacle to the fulfilment of a child's right to identity. Data shows that only 24 percent of the mothers/caretakers of the children under five years of age whose births are not registered know how to register a child's birth.

Knowledge on how to register a child's birth varies by urban/rural areas, mother's/caretaker's education and household wealth.

Figure CP.1: Children under-5 whose births are registered, Turkana County MICS, 2013/14



10.2 Child Labour

Children around the world are routinely engaged in paid and unpaid forms of work that are not harmful to them. However, they are classified as child labourers when they are either too young to work or are involved in hazardous activities that may compromise their physical, mental, social or educational development. Article 32 (1) of the Convention on the Rights of the Child states: "State Parties recognize the right of the child to be protected from economic exploitation and from performing any work that is likely to be hazardous or to interfere with the child's education, or to be harmful to the child's health or physical, mental, spiritual, moral or social development".

The Employment Act [Chapter 226] 2007, and the Children Act [Chapter 141] 2007, define a child in Kenya as a person below the age of 18 years. The Employment Act, Part VII provides for protection of children including protection from the worst forms of child labour. Section 56 of the Employment Act prohibits employment of a child below age 13 years in any form of undertaking. However, it allows employment of children from age 13 to 16 years for light work, and defines those of age 16 to 18 as employable.^{104, 105}

In Turkana County, the child labour module was administered for children age 5-17 and includes questions on the type of work a child does and the number of hours he or she is engaged in it. Data were collected on both economic activities (paid or unpaid work for someone who is not a member of the household, work for a family farm or business) and domestic work (household chores such as cooking, cleaning or

¹⁰⁴ Employment Act [Chapter 226] 2007, 2012; Children Act [141] 2007, 2010.

¹⁰⁵ http://www.kenyalaw.org/kl/fileadmin/pdfdownloads/Acts/EmploymentAct_Cap226-No11of2007_01.pdf

caring for children, as well as collecting firewood or fetching water). The module also collected information on hazardous working conditions.^{106, 107}

Table CP.2 presents children's involvement in economic activities during the last week preceding the survey. The methodology of the MICS on Child Labour uses three age-specific thresholds for the number of hours a child can perform an economic activity without it being classified as in child labour. A child that performed economic activities during the last week for more than the age-specific number of hours is classified as in child labour:

- i. age 5-11: 1 hour or more
- ii. age 12-14: 14 hours or more
- iii. age 15-17: 43 hours or more

While about a third of the 5-11 year olds were involved in economic activities for at least one hour, about half of 12-14 year olds were involved in some economic activity (42 percent for less than 14 hours and four percent for more than 14 hours), and about half of the 15-17 year olds were involved in economic activities for less than 43 hours. Fourteen percent of the out of school children 12-14 years were engaged in economic activities for 14 hours or more, compared with only one percent of the children 12-14 years old attending school (Table CP.2).

¹⁰⁶ UNICEF. 2012. *How Sensitive Are Estimates of Child Labour to Definitions?* MICS Methodological Paper No. 1. UNICEF.

¹⁰⁷ The Child Labour module and the Child Discipline module were administered using random selection of a single child in all households with one or more children age 1-17 (See Appendix H: Questionnaires). The Child Labour module was administered if the selected child was age 5-17 and the Child Discipline module if the child was age 1-14 years old. To account for the random selection, the household sample weight is multiplied by the total number of children age 1-17 in each household.

Table CP.2: Children's involvement in economic activities

Percentage of children by involvement in economic activities during the last week, according to age groups, Turkana County MICS, 2013/14								
	Percentage of children age 5-11 years involved in economic activity for at least one hour	Number of children age 5-11 years	Percentage of children age 12-14 years involved in:		Number of children age 12-14 years	Percentage of children age 15-17 years involved in:		Number of children age 15-17 years
			Economic activity less than 14 hours	Economic activity for 14 hours or more		Economic activity less than 43 hours	Economic activity for 43 hours or more	
Total	32.0	1,476	41.6	4.0	645	51.5	0.0	501
Sex								
Male	37.7	760	36.5	5.5	321	46.3	0.0	295
Female	25.9	715	46.6	2.5	324	59.0	0.0	207
Area								
Urban	24.4	752	42.1	4.1	368	49.6	0.0	313
Rural	39.8	723	41.0	3.9	277	54.7	0.0	188
School attendance								
Yes	30.0	964	42.5	0.9	492	48.2	0.0	356
No	35.6	511	38.6	14.1	153	59.6	0.0	145
Mother's education								
None	33.9	1,074	44.1	5.7	457	63.5	0.0	219
Primary	28.0	259	34.2	0.0	99	(63.6)	(0.0)	46
Secondary	25.3	140	42.2	0.0	72	(*)	(*)	10
Cannot be determined ^a	(*)	1	(*)	(*)	11	36.6	0.0	226
Wealth index quintile								
Poorest	42.0	345	46.8	6.6	107	56.4	0.0	64
Second	35.9	315	42.2	0.0	128	72.5	0.0	76
Middle	27.9	248	32.4	4.8	166	42.6	0.0	80
Fourth	21.8	314	39.3	8.5	129	30.2	0.0	111
Richest	30.1	254	51.8	0.0	115	58.4	0.0	171
Ethnicity of household head								
Turkana	34.2	1,271	45.7	5.0	522	51.9	0.0	388
Other ethnic group	18.0	203	24.2	0.0	123	48.1	0.0	109
^a Children age 15 or higher at the time of the interview whose mothers were not living in the household								
na: not applicable								
() Figures that are based on 25-49 unweighted cases								
(*) Figures that are based on fewer than 25 unweighted cases								

Table CP.3 presents children's involvement in household chores. Like for economic activity above, the methodology also uses age-specific thresholds for the number of hours a child can perform household chores without it being classified as child labour. A child who performed household chores during the last week for more than the age-specific number of hours is classified as in child labour:

- i. age 5-11 and age 12-14: 28 hours or more
- ii. age 15-17: 43 hours or more

Table CP.3 shows that one percent of children age 5-11 years and three percent of children age 12-14 years were involved in household chores for 28 hours or more while one percent of children age 15-17 years were involved in household chores for 43 hours or more.

Table CP.3: Children's involvement in household chores

Percentage of children by involvement in household chores during the last week, according to age groups, Turkana County MICS											
	Percentage of children age 5-11 years involved in:			Number of children age 5-11 years	Percentage of children age 12-14 years involved in:			Number of children age 12-14 years	Percentage of children age 15-17 years involved in:		
	Household chores less than 28 hours	Household chores for 28 hours or more			Household chores less than 28 hours	Household chores for 28 hours or more			Household chores less than 43 hours	Household chores for 43 hours or more	
Total	73.1	1.0	1,476		88.6	3.1	645		89.7	0.8	501
Sex											
Male	71.1	0.8	760		87.0	0.7	321		89.0	0.4	295
Female	75.4	1.2	715		90.2	5.5	324		90.8	1.3	207
Area											
Urban	73.0	0.5	752		87.4	3.7	368		94.7	0.3	313
Rural	73.3	1.6	723		90.2	2.3	277		81.5	1.5	188
School attendance											
Yes	76.9	1.3	964		93.4	3.2	492		92.8	0.3	356
No	66.1	0.4	511		73.0	2.7	153		82.1	1.9	145
Mother's education											
None	72.2	1.2	1,074		88.5	2.6	457		93.7	1.3	219
Primary	79.0	1.0	259		95.5	1.0	99		(74.4)	(0.0)	46
Secondary	69.3	0.0	140		88.3	10.2	72		(*)	(*)	10
Cannot be determined ^a	(*)	(*)	1		(*)	(*)	11		88.5	0.5	226
Wealth index quintile											
Poorest	68.8	2.3	345		90.1	1.3	107		72.3	4.3	64
Second	71.7	0.3	315		89.5	7.2	128		93.8	1.4	76
Middle	76.8	1.4	248		81.8	1.3	166		83.9	0.0	80
Fourth	75.0	0.8	314		91.5	0.0	129		91.8	0.0	111
Richest	74.9	0.0	254		92.7	6.4	115		95.8	0.0	171
Ethnicity of household head											
Turkana	74.6	1.2	1,271		88.1	3.8	522		90.0	1.0	388
Other ethnic group	64.7	0.0	203		91.0	0.0	123		88.4	0.0	109
^a Children age 15 or higher at the time of the interview whose mothers were not living in the household											
na: not applicable											
() Figures that are based on 25-49 unweighted cases											
(*) Figures that are based on fewer than 25 unweighted cases											

Table CP.4 combines the children working and performing household chores at or above and below the age-specific thresholds as detailed in the previous tables, as well as those children reported working under hazardous conditions, into the total child labour indicator. Total child labour for Turkana County is 33 percent (35 percent for boys and 31 percent for girls). Child labour is slightly higher in rural areas (37

percent) than urban areas (30 percent). Further, child labour is higher for children less than 10 years and for those not attending school.

Table CP.4: Child labour

Percentage of children age 5-17 years by involvement in economic activities or household chores during the last week, percentage working under hazardous conditions during the last week, and percentage engaged in child labour during the last week, Turkana County MICS, 2013/14

	Children involved in economic activities for a total number of hours during last week:		Children involved in household chores for a total number of hours during last week:		Children working under hazardous conditions	Total child labour ¹	Number of children age 5-17 years
	Below the age specific threshold	At or above the age specific threshold	Below the age specific threshold	At or above the age specific threshold			
Total	22.3	19.0	80.1	1.5	22.7	32.8	2,622
Sex							
Male	21.0	22.1	78.6	0.7	23.9	34.5	1,375
Female	23.6	15.5	81.8	2.4	21.3	30.9	1,247
Area							
Urban	25.5	13.9	81.4	1.3	21.5	29.6	1,434
Rural	18.3	25.2	78.5	1.7	24.2	36.6	1,188
Age							
5-11	3.9	32.0	73.1	1.0	18.2	34.7	1,476
12-14	41.6	4.0	88.6	3.1	28.2	31.0	645
15-17	51.5	0.0	89.7	0.8	28.8	29.5	501
School attendance							
Yes	23.8	16.2	84.5	1.7	21.5	30.7	1,813
No	18.9	25.2	70.3	1.1	25.5	37.4	809
Mother's education							
None	21.3	22.3	79.1	1.5	25.0	35.5	1,750
Primary	18.8	18.0	82.5	0.9	14.1	27.1	404
Secondary+	22.0	15.9	76.9	3.3	18.0	30.3	222
Cannot be determined ^a	36.0	0.0	85.8	0.4	25.7	26.1	238
Wealth index quintile							
Poorest	16.7	29.4	73.6	2.3	28.0	43.3	516
Second	22.7	21.8	79.4	2.2	27.7	36.1	518
Middle	20.3	15.6	79.7	1.2	16.6	25.7	494
Fourth	20.2	14.3	82.2	0.5	14.3	21.3	553
Richest	31.0	14.1	85.3	1.4	27.0	37.7	540
Ethnicity of household head							
Turkana	22.2	21.1	80.6	1.8	24.7	36.1	2,180
Other ethnic group	22.0	8.4	78.1	0.0	12.0	15.3	435

¹ MICS indicator 8.2 - Child labour

^a Children age 15 or higher at the time of the interview whose mothers were not living in the household

(*) Figures that are based on fewer than 25 unweighted cases

10.3 Child Discipline

Teaching children self-control and acceptable behaviour is an integral part of child discipline in all cultures. Positive parenting practices involve providing guidance on how to handle emotions or conflicts in manners that encourage judgment and responsibility and preserve children's self-esteem, physical and psychological integrity and dignity. Too often, however, children are raised through the use of punitive methods that rely on the use of physical force or verbal intimidation to obtain desired behaviours. Studies¹⁰⁸ have found that exposing children to violent discipline have harmful consequences, which range from immediate impacts to long-term harm that children carry forward into adult life. Violence hampers children's development, learning abilities and school performance; it inhibits positive relationships, provokes low self-esteem, emotional distress and depression; and, at times, it leads to risk taking and self-harm.

Respondents to the household questionnaire in Turkana County were asked a series of questions on the methods adults in the household used to discipline a selected child during the past month. The disciplinary methods assessed ranged from non-violent approaches to psychological aggression, and moderate to severe forms of physical punishment.

Non-violent discipline: Took away privileges; explained wrong behaviour; gave the child something else to do.

Psychological aggression: Shouted, yelled, screamed; called the child 'dumb, lazy or any other name'.

Physical punishment: Shook the child; spanked, hit, slapped on bottom with bare hand; hit with belt, hairbrush, stick or other hard object; hit/slapped on the face, head or ears; hit/slapped on hand, arm or leg; beat up, hit over and over as hard as one could.

Severe punishment: hit/slapped on the face, head or ears; hit/slapped on hand, arm or leg; beat up, hit over and over as hard as one could.

Any violent discipline method: Shook the child; shouted, yelled, screamed; spanked, hit, slapped on bottom with bare hand; hit with belt, hairbrush, stick or other hard object; called the child 'dumb, lazy or any other name'; hit/slapped on the face, head or ears; hit/slapped on hand, arm or leg; beat up, hit over and over as hard as one could.

¹⁰⁸Straus, MA and Paschall MJ.2009. *Corporal Punishment by Mothers and Development of Children's Cognitive Ability: A longitudinal study of two nationally representative age cohorts.* Journal of Aggression, Maltreatment & Trauma 18(5): 459-83.
Erickson, MF and Egeland, B. 1987. *A Developmental View of the Psychological Consequences of Maltreatment.* School Psychology Review 16: 156-68.

Schneider, MW et al. 2005. *Do Allegations of Emotional Maltreatment Predict Developmental Outcomes Beyond that of Other Forms of Maltreatment?* Child Abuse & Neglect 29 (5): 513-32.

Table CP.5 shows that 64 percent of children age 1-14 years were subjected to at least one form of psychological aggression or physical punishment by household members during the past month. For the most part, households employ a combination of violent disciplinary practices, reflecting caregivers’ motivation to control children’s behaviour by any means possible. While 33 percent of children experienced psychological aggression, about 60 percent experienced some form of physical punishment. The most severe forms of physical punishment (hitting the child on the head, ears or face or hitting the child hard and repeatedly) were overall less common: 10 percent of children were subjected to severe punishment.

In rural areas, 66 percent of children age 1-14 years were subjected to at least one form of psychological or physical punishment by household members during the past month and 63 percent in urban areas. Differences are observed by education of head of household, where violent discipline of children was 67 percent for children whose head of household had no education, 62 percent for primary education, and 58 percent for those with secondary or higher education. Figure CP.2 presents a summary of the main methods of child discipline.

Figure CP.2: Child disciplining methods, children age 1-14 years, Turkana County MICS, 2013/14




Table CP.5: Child discipline						
Percentage of children age 1-14 years by child disciplining methods experienced during the last one month, Turkana County MICS, 2013/14						
	Percentage of children age 1-14 years who experienced:					Number of children age 1-14 years
	Only non-violent discipline	Psychological aggression	Physical punishment		Any violent discipline method ¹	
			Any	Severe		
Total	25.1	33.1	60.2	10.4	64.3	2,982
Sex						
Male	25.1	30.1	61.3	8.3	64.5	1,511
Female	25.1	36.1	59.1	12.5	64.2	1,471
Area						
Urban	29.1	32.5	57.7	8.9	62.5	1,552
Rural	20.8	33.7	63.0	12.1	66.3	1,430
Age						
1-2	18.9	22.6	50.4	5.1	52.5	436
3-4	14.6	35.9	73.4	11.6	74.8	425
5-9	27.4	35.2	64.3	13.1	68.5	1,077
10-14	29.6	34.0	54.8	9.4	60.8	1,044
Education of household head						
None	21.1	33.4	63.6	11.0	66.6	2,047
Primary	32.0	35.3	56.4	9.3	62.1	448
Secondary+	36.3	29.4	52.8	9.7	58.2	448
Wealth index quintile						
Poorest	20.1	34.5	64.8	11.3	68.3	635
Second	22.3	35.1	65.8	12.1	67.1	640
Middle	26.3	27.5	52.7	8.0	56.3	624
Fourth	31.0	31.7	55.8	7.9	62.0	603
Richest	26.4	37.4	62.0	13.3	68.9	480
Ethnicity of household head						
Turkana	21.4	36.1	66.0	11.5	70.0	2,528
Other ethnic group	45.6	15.9	28.3	4.2	32.8	448
¹ MICS indicator 8.3 - Violent discipline						
() Figures that are based on 25-49 unweighted cases						

Table CP.6 shows that 81 percent of respondents to the household questionnaire believe that physical punishment is a necessary part of child-rearing. There are some differentials across background variables of respondents. Overall, respondents with low educational attainment, those residing in rural areas and those from poorer households are more likely to find physical punishment as necessary in disciplining children.

Table CP.6: Attitudes toward physical punishment		
Percentage of respondents to the child discipline module who believe that physical punishment is needed to bring up, raise, or educate a child properly, Turkana County MICS, 2013/14		
	Respondent believes that a child needs to be physically punished	Number of respondents to the child discipline module
Total	80.7	913
Sex		
Male	76.4	184
Female	81.8	729
Area		
Urban	76.9	465
Rural	84.7	449
Age		
<25	71.6	164
25-39	84.7	404
40-59	80.0	243
60+	81.6	102
Respondent's relationship to selected child		
Mother	82.9	496
Father	79.7	145
Other	77.2	272
Respondent's education		
None	84.2	637
Primary	79.4	152
Secondary+	63.7	122
Wealth index quintile		
Poorest	81.0	199
Second	91.0	212
Middle	80.0	188
Fourth	75.9	164
Richest	72.0	149
Ethnicity of household head		
Turkana	84.4	781
Other ethnic group	58.7	130

10.4 Early Marriage and Polygyny

Marriage¹⁰⁹ before the age of 18 is a reality for many young girls. In many parts of the world, parents encourage the marriage of their daughters while they are still children in hopes that the marriage will benefit them both financially and socially, while also relieving financial burdens on the family. In actual fact, child marriage is a violation of human rights, compromising the development of girls and often resulting in early pregnancy and social isolation, with little education and poor vocational training reinforcing the gendered nature of poverty.¹¹⁰ The right to 'free and full' consent to a marriage is recognized in the Universal Declaration of Human Rights - with the recognition that consent cannot be 'free and full' when one of the parties involved is not sufficiently mature to make an informed decision about a life partner. Closely related to the issue of child marriage is the age at which girls become sexually active. Women who are married before the age of 18 tend to have more children than those who marry later in life. Pregnancy related deaths are known to be a leading cause of mortality for both married and unmarried girls between the ages of 15 and 19, particularly among the youngest of this cohort. There is evidence to suggest that girls who marry at young ages are more likely to marry older men which puts them at increased risk of HIV infection. The demand for such a young wife to reproduce and the power imbalance resulting from the age differential lead to very low condom use among such couples.¹¹¹

In Turkana County MICS, the percentage of women married before ages 15 and 18 years are provided in Table CP.7. Among women age 15-49 years, nine percent were married before age 15 and, among women age 20-49 years, 11 percent were married before age 15 while 32 percent were married before age 18.

Fourteen percent of young women age 15-19 years are currently married. This proportion does not vary much between urban (13 percent) and rural (17 percent). The percentage of women in a polygynous union is also provided in Table CP.7. Among all women age 15-49 years who are in union, 30 percent are in polygynous unions.

¹⁰⁹ All references to marriage in this chapter include marital union as well.

¹¹⁰ Bajracharya, A ND Amin, S. 2010. *Poverty, marriage timing, and transitions to adulthood in Nepal: A longitudinal analysis using the Nepal living standards survey*. Poverty, Gender, and Youth Working Paper No. 19. Population Council.

Godha, D et al. 2011. *The influence of child marriage on fertility, fertility-control, and maternal health care utilization*. MEASURE/Evaluation PRH Project Working paper 11-124.

¹¹¹ Clark, S et al. 2006. *Protecting young women from HIV/AIDS: the case against child and adolescent marriage*. International Family Planning Perspectives 32(2): 79-88.

Raj, A et al. 2009. *Prevalence of child marriage and its effect on fertility and fertility-control outcomes of young women in India: a cross-sectional, observational study*. The Lancet 373(9678): 1883-9.

Table CP.7: Early marriage and polygyny (women)

Percentage of women age 15-49 years who first married or entered a marital union before their 15th birthday, percentages of women age 20-49 years who first married or entered a marital union before their 15th and 18th birthdays, percentage of women age 15-19 years currently married or in union, and the percentage of women who are in a polygynous marriage or union, Turkana County MICS, 2013/14

	Women age 15-49 years		Women age 20-49 years			Women age 15-19 years		Women age 15-49 years	
	Percentage married before age 15 ¹	Number of women age 15-49 years	Percentage married before age 15	Percentage married before age 18 ²	Number of women age 20-49 years	Percentage currently married/in union ³	Number of women age 15-19 years	Percentage in polygynous marriage/union ⁴	Number of women age 15-49 years currently married/in union
Total	8.9	1,104	11.1	32.2	852	13.9	252	30.0	615
Area									
Urban	9.3	683	12.2	32.6	491	13.0	192	23.6	320
Rural	8.3	421	9.5	31.6	361	16.8	60	36.9	295
Age									
15-19	1.7	252	na	na	na	13.9	252	(24.3)	34
20-24	7.5	209	7.5	30.8	209	na	na	27.0	110
25-29	18.0	210	18.0	37.3	210	na	na	26.2	166
30-34	14.1	144	14.1	40.2	144	na	na	36.4	114
35-39	5.3	134	5.3	27.9	134	na	na	25.0	95
40-44	4.8	82	4.8	20.8	82	na	na	36.7	52
45-49	13.2	75	13.2	26.1	75	na	na	(42.8)	43
Education									
None	10.9	622	12.0	33.3	557	18.2	65	35.3	408
Primary	6.7	296	11.1	40.8	167	11.9	128	19.7	121
Secondary+	6.0	186	7.3	16.1	129	13.7	58	19.3	86
Wealth index quintile									
Poorest	8.6	169	9.8	27.8	144	18.1	25	44.8	115
Second	7.9	191	8.9	27.0	169	12.5	22	36.3	123
Middle	9.5	226	11.5	38.7	177	13.2	49	27.7	104
Fourth	11.6	249	14.3	38.3	185	26.0	64	24.8	138
Richest	6.9	270	10.4	27.8	179	5.0	91	18.6	135

Ethnicity of household head									
Turkana	8.9	883	10.8	32.1	684	15.8	198	32.2	526
Other ethnic group	9.1	218	12.0	32.4	165	6.9	53	16.3	88
¹ MICS indicator 8.4 - Marriage before age 15 ² MICS indicator 8.5 - Marriage before age 18 ³ MICS indicator 8.6 - Young women age 15-19 years currently married or in union ⁴ MICS indicator 8.7 – Polygyny									
na: not applicable									
() Figures that are based on 25-49 unweighted cases									

Table CP.8 presents the proportion of women who were first married or entered into a marital union before age 15 years and 18 years by area and age group. Examining the percentages married before age 15 years and 18 years by different age groups allows for trends to be observed in early marriage over time. Overall, nine percent of women age 15-49 years had married before age 15 while 32 percent had married before age 18. The proportion of women age 15-49 years who had married before age 15 was nine percent in urban areas, eight percent in rural areas. The proportion of all women age 15-19 years who had married before age 15 was two percent. Comparing with the same age group 10 years ago (those currently 25-29 years), 18 percent had married before age 15. Comparing with the same age group thirty years ago (those currently 45-49 years), 13 percent had married before age 15. Figure CP.3 presents a summary of the main characteristics of early marriage.

Table CP.8: Trends in early marriage (women)


Percentage of women who were first married or entered into a marital union before age 15 and 18, by area and age groups, Turkana County MICS, 2013/14

	Urban				Rural				All			
	Percentage of women married before age 15	Number of women age 15-49 years	Percentage of women married before age 18	Number of women age 20-49 years	Percentage of women married before age 15	Number of women age 15-49 years	Percentage of women married before age 18	Number of women age 20-49 years	Percentage of women married before age 15	Number of women age 15-49 years	Percentage of women married before age 18	Number of women age 20-49 years
Total	9.3	683	32.6	491	8.3	421	31.6	361	8.9	1,104	32.2	852
Age												
15-19	1.9	192	na	na	0.8	60	na	na	1.7	252	na	na
20-24	9.0	153	30.9	153	3.3	56	30.5	56	7.5	209	30.8	209
25-29	22.8	111	37.6	111	12.6	99	36.9	99	18.0	210	37.3	210
30-34	14.8	71	44.6	71	13.4	73	36.0	73	14.1	144	40.2	144
35-39	5.0	75	30.7	75	5.6	59	24.2	59	5.3	134	27.9	134
40-44	(2.1)	38	(16.5)	38	7.1	44	24.6	44	4.8	82	20.8	82
45-49	(13.4)	44	(23.2)	44	(12.8)	31	(30.4)	31	13.2	75	26.1	75

na: not applicable

() Figures that are based on 25-49 unweighted cases

Figure CP.3: Early marriage among women, Turkana County MICS, 2013/14



na: not applicable

Another important component of child marriage is the spousal age difference since the age difference between husband and wife is likely to have implications for power dynamics within the household. Table CP.9 shows that the proportion of women age 20-24 years currently married or in union with a husband or partner 10 or more years older than them is 21 percent.¹¹²

Table CP.9: Spousal age difference

Percent distribution of women currently married/in union age 20-24 years according to the age difference with their husband or partner, Turkana County MICS, 2013/14

	Percentage of currently married/in union women age 20-24 years whose husband or partner is:					Total	Number of women age 20-24 years currently married/in union
	Younger	0-4 years older	5-9 years older	10+ years older ²	Husband/Partner's age unknown		
Total	4.3	38.0	31.8	21.3	4.7	100.0	106
² MICS indicator 8.8b - Spousal age difference (among women age 20-24)							
na: not applicable							

¹¹² The cases for women age 15-19 years currently married/in union were too few to be analysed by the age of the husband/partner. As such ¹ MICS indicator 8.8a - Spousal age difference (among women age 15-19) is not shown.

10.5 Female Genital Mutilation/Cutting

Female genital mutilation/cutting (FGM/C) is the partial or total removal of the female external genitalia or other injury to the female genital organs. FGM/C is always traumatic with immediate complications including excruciating pain, shock, urine retention, ulceration of the genitals and injury to adjacent tissue. Other complications include septicaemia, infertility, obstructed labour, and even death. The procedure is generally carried out on girls between the ages of four and 14; it is also done to infants, women who are about to be married, and sometimes, to women who are pregnant with their first child or who have just given birth. It is often performed by traditional practitioners, including midwives and barbers, without anaesthesia, using scissors, razor blades, or broken glass.

FGM/C is a fundamental violation of human rights which subjects girls and women to health risks and has life-threatening consequences. Although no international human rights instruments specifically address the practice, Article 25 of the Universal Declaration of Human Rights states that “everyone has the right to a standard of living adequate for health and well-being” and has been used to argue that FGM/C violates the right to health and bodily integrity. Furthermore, it could be argued that girls, i.e. children, cannot be said to give informed consent to such a potentially damaging practice as FGM/C.

Table CP.10 presents the prevalence of FGM/C among women age 15-49 years and the type of procedure performed from the Turkana County MICS survey. Three percent of women have some form of female genital mutilation. The practice is five percent prevalent in urban areas and less than one percent in rural areas.

Table CP.10: Female genital mutilation/cutting (FGM/C) among women

Percentage of women age 15-49 years by FGM/C status and percent distribution of women who had FGM/C by type of FGM/C, Turkana County MICS, 2013/14

	Percentage of women who had any form of FGM/C ¹	Number of women age 15-49 years
Total	3.2	1,104
Area		
Urban	5.0	683
Rural	0.2	421
Age		
15-19	1.3	252
20-24	6.0	209
25-29	3.9	210
30-34	1.3	144
35-39	3.3	134
40-44	3.1	82
45-49	2.8	75
Education		
None	3.0	622
Primary	4.3	296
Secondary+	1.8	186
Wealth index quintile		
Poorest	0.0	169
Second	0.0	191
Middle	6.4	226
Fourth	5.8	249
Richest	2.3	270
Ethnicity of household head		
Turkana	0.0	883
Other ethnic group	16.0	218
¹ MICS indicator 8.10 - Prevalence of FGM/C among women		

Table CP.11 presents the prevalence and extent of FGM/C performed on all daughters, age 0-14 years, of the respondents. It is important to remember that prevalence data for girls age 0-14 years reflect their current – not final – FGM/C status, since many of them may not have reached the customary age for cutting at the time of the survey. They are reported as being uncut but are still at risk of undergoing the procedure. Overall, two percent of girls age 0-14 years have undergone FGM/C.

Table CP.11: Female genital mutilation/cutting (FGM/C) among girls

Percentage of daughters age 0-14 years by FGM/C status and percent distribution of daughters who had FGM/C by type of FGM/C, Turkana County MICS, 2013/14

	Percentage of daughters who had any form of FGM/C ¹	Number of daughters age 0-14 years
Total	1.9	381
Area		
Urban	3.0	244
Rural	0.0	137
Age		
0-4	0.0	146
5-9	2.0	126
10-14	4.4	109
Mother's Education		
None	2.1	221
Primary	1.4	102
Secondary+	(1.9)	58
Mother's FGM/C experience		
No FGM/C	0.4	355
Had FGM/C	(*)	26
Wealth index quintile		
Poorest	3.6	38
Second	0.0	68
Middle	5.3	63
Fourth	1.1	98
Richest	1.3	114
Ethnicity of household head		
Turkana	0.5	287
Other ethnic group	6.3	94
¹ MICS indicator 8.11 - Prevalence of FGM/C among girls		
() Figures that are based on 25-49 unweighted cases (*) Figures that are based on fewer than 25 unweighted cases		

Results on perceptions of women age 15-49 years towards FGM/C are presented in Table CP.12. As to whether the practice should be continued or discontinued, six percent of women think it should be continued while 87 percent believe it should be discontinued. Support for the continuation of the practice is higher among women in urban areas than rural areas.

Table CP.12: Approval of female genital mutilation/cutting (FGM/C)

Percentage of women age 15-49 years who have heard of FGM/C, and percent distribution of women according to attitudes towards whether the practice of FGM/C should be continued, Turkana County MICS, 2013/14

	Percentage of women who have heard of FGM/C	Number of women age 15-49 years	Percent distribution of women who believe the practice of FGM/C should be:				Total	Number of women age 15-49 years who have heard of FGM/C
			Continued ¹	Discontinued	Depends	DK/Missing		
Total	46.2	1,104	5.9	86.6	1.9	5.6	100.0	510
Area								
Urban	57.0	683	7.2	88.4	0.8	3.6	100.0	389
Rural	28.7	421	1.8	81.0	5.3	11.9	100.0	121
Age								
15-19	60.1	252	5.0	89.3	2.4	3.3	100.0	151
20-24	51.6	209	10.7	83.0	2.2	4.0	100.0	108
25-29	39.4	210	3.0	91.0	0.0	6.0	100.0	83
30-34	40.5	144	3.2	86.6	3.1	7.2	100.0	58
35-39	39.5	134	6.2	78.3	2.1	13.4	100.0	53
40-44	41.0	82	(6.7)	(87.5)	(2.1)	(3.7)	100.0	33
45-49	31.8	75	(*)	(*)	(*)	(*)	100.0	24
Education								
None	32.1	622	6.6	80.1	1.9	11.4	100.0	200
Primary	56.3	296	6.6	89.0	3.2	1.2	100.0	166
Secondary+	77.3	186	4.0	92.9	0.4	2.7	100.0	144
FGM/C experience								
No FGM/C	44.4	1,069	4.1	87.8	2.0	6.0	100.0	475
Had FGM/C	(100.0)	35	(29.6)	(70.4)	(0.0)	(0.0)	100.0	35
Wealth index quintile								
Poorest	24.4	169	3.3	80.4	4.4	11.9	100.0	41
Second	30.6	191	3.0	85.2	0.0	11.9	100.0	58
Middle	42.6	226	8.8	77.6	6.1	7.5	100.0	96
Fourth	55.5	249	3.5	93.0	0.5	3.0	100.0	138
Richest	65.3	270	7.8	88.5	0.7	3.0	100.0	176
Ethnicity of household head								
Turkana	41.8	883	4.4	85.8	2.6	7.2	100.0	369
Other ethnic group	63.8	218	10.0	88.6	0.0	1.4	100.0	139
¹ MICS indicator 8.9 - Approval for FGM/C								
() Figures that are based on 25-49 unweighted cases (*) Figures that are based on fewer than 25 unweighted cases								

10.6 Attitudes toward Domestic Violence

MICS assessed the attitudes of women age 15-49 years towards wife/partner beating by asking the respondents whether husbands/partners were justified to hit or beat their wives/partners in a variety of situations. The purpose of these questions was to capture the social justification of violence (in contexts where women have a lower status in society) as a disciplinary action when a woman does not comply with certain expected gender roles.

In Turkana County MICS, the responses to these questions can be found in Table CP.13. Overall, 70 percent of women in Turkana County feel that a husband/partner is justified in hitting or beating his wife in at least one of the five situations. Women who justify a husband's violence, in most cases, agree and justify violence in instances when a wife neglects the children (55 percent), or if she demonstrates her autonomy, exemplified by going out without telling her husband (51 percent) or argues with him (48 percent). Around one in three women (36 percent) believe that wife-beating is justified if the wife refuses to have sex with the husband and about one in four agree that wife-beating is justified if she burns the food (27 percent). Justification in any of the five situations is more prevalent among women less educated, and also currently or formerly married.

Table CP.13: Attitudes toward domestic violence (women)

Percentage of women age 15-49 years who believe a husband is justified in beating his wife in various circumstances, Turkana County MICS, 2013/14

	Percentage of women age 15-49 years who believe a husband is justified in beating his wife:						Number of women age 15-49 years
	If she goes out without telling him	If she neglects the children	If she argues with him	If she refuses sex with him	If she burns the food	For any of these five reasons ¹	
Total	51.4	54.6	47.5	36.3	26.6	70.1	1,104
Area							
Urban	47.4	52.0	42.6	31.6	22.9	70.2	683
Rural	57.8	58.8	55.5	43.8	32.5	70.0	421
Age							
15-19	44.5	51.1	37.6	31.6	23.0	67.8	252
20-24	48.3	50.6	42.2	27.0	21.3	66.3	209
25-29	54.6	58.7	55.3	39.4	28.9	70.3	210
30-34	55.9	59.9	51.5	41.7	27.1	75.2	144
35-39	54.5	56.2	51.2	39.9	30.4	70.3	134
40-44	56.0	53.8	52.9	46.8	31.6	71.6	82
45-49	54.4	53.8	54.1	40.9	34.0	76.3	75
Marital/Union status							
Currently married/in union	54.0	57.2	52.9	40.4	28.1	71.5	615
Formerly married/in union	52.9	55.1	46.1	39.5	34.1	70.6	171
Never married/in union	44.9	48.9	37.8	26.7	19.3	66.8	314
Education							
None	57.4	59.6	54.8	43.0	31.7	72.9	622
Primary	49.9	51.1	40.9	31.8	23.9	71.5	296
Secondary+	33.7	43.3	33.6	21.1	13.9	58.8	186
Wealth index quintile							
Poorest	58.6	59.6	58.1	48.4	33.4	74.0	169
Second	58.2	62.2	57.0	45.4	35.3	70.1	191
Middle	43.4	52.1	39.5	29.5	19.3	64.0	226
Fourth	64.0	58.9	54.2	38.6	31.3	78.7	249
Richest	37.0	44.2	34.7	25.8	17.9	64.9	270
Ethnicity of household head							
Turkana	52.0	57.1	49.8	38.0	27.4	72.0	883
Other ethnic group	48.3	45.0	37.6	29.4	23.5	62.0	218

¹ MICS indicator 8.12 - Attitudes towards domestic violence

10.7 Children's Living Arrangements

The CRC recognizes that “the child, for the full and harmonious development of his or her personality, should grow up in a family environment, in an atmosphere of happiness, love and understanding”. Millions of children around the world grow up without the care of their parents for several reasons, including due to the premature death of the parents or their migration for work. In most cases, these children are cared for by members of their extended families, while in others, children may be living in households other than their own, as live-in domestic workers for instance. Understanding the children's living arrangements, including the composition of the households where they live and the relationships with their primary caregivers, is key to design targeted interventions aimed at promoting child's care and wellbeing.

In Turkana County MICS, information on the living arrangements and orphanhood status of children under age 18 is presented in Table CP.14. About 53 percent of children age 0-17 years in Turkana County live with both their parents. Nearly 18 percent of children live with neither of their biological parents and the proportion is higher in urban areas (21 percent) than rural areas (14 percent). Older children are more likely than younger children to live with neither biological parent. Overall, 18 percent of the children age 0-17 years have lost one or both parents.

Table CP.14: Children's living arrangements and orphanhood

Percent distribution of children age 0-17 years according to living arrangements, percentage of children age 0-17 years not living with a biological parent and percentage of children who have one or both parents dead, Turkana County MICS, 2013/14

	Living with both parents	Living with neither biological parent				Living with mother only		Living with father only		Missing information on father/mother	Total	Living with neither biological parent ¹	One or both parents dead ²	Number of children age 0-17 years
		Only father alive	Only mother alive	Both alive	Both dead	Father alive	Father dead	Mother alive	Mother dead					
Total	53.0	0.0	0.0	9.7	0.0	14.2	0.0	2.4	0.0	20.7	100.0	17.7	18.4	3,699
Sex														
Male	52.9	0.0	0.0	9.0	0.0	14.3	0.0	2.5	0.0	21.3	100.0	16.6	18.7	1,876
Female	53.1	0.0	0.0	10.4	0.0	14.2	0.0	2.2	0.0	20.1	100.0	18.8	18.1	1,823
Area														
Urban	44.3	0.0	0.0	9.7	0.0	15.9	0.0	3.2	0.0	27.0	100.0	20.8	23.3	1,975
Rural	62.9	0.0	0.0	9.7	0.0	12.4	0.0	1.4	0.0	13.6	100.0	14.0	12.9	1,724
Age														
0-4	66.1	0.0	0.0	4.3	0.0	19.5	0.0	0.4	0.0	9.7	100.0	6.2	7.8	1,095
5-9	55.5	0.0	0.0	10.4	0.0	12.3	0.0	2.8	0.0	19.0	100.0	16.7	16.9	1,096
10-14	46.1	0.0	0.0	12.4	0.0	12.2	0.0	3.6	0.0	25.8	100.0	23.9	23.9	1,018
15-17	32.6	0.0	0.0	14.6	0.0	10.9	0.0	3.3	0.0	38.5	100.0	32.7	34.2	489
Wealth index quintile														
Poorest	62.0	0.0	0.0	10.9	0.0	12.0	0.0	1.4	0.0	13.6	100.0	14.5	12.9	742
Second	56.1	0.0	0.0	10.4	0.0	14.8	0.0	1.4	0.0	17.4	100.0	16.2	17.3	763
Middle	37.8	0.0	0.0	9.6	0.0	16.9	0.0	2.5	0.0	33.2	100.0	20.3	27.5	746
Fourth	55.8	0.0	0.0	7.2	0.0	13.7	0.0	2.7	0.0	20.6	100.0	17.1	17.0	762
Richest	53.2	0.0	0.0	10.4	0.0	13.7	0.0	3.9	0.0	18.8	100.0	20.4	17.3	686
Ethnicity of household head														
Turkana	55.3	0.0	0.0	9.7	0.0	14.0	0.0	2.3	0.0	18.7	100.0	17.5	17.2	3,100
Other ethnic group	40.7	0.0	0.0	9.9	0.0	15.6	0.0	2.6	0.0	31.2	100.0	18.7	24.3	588
¹ MICS indicator 8.13 - Children's living arrangements														
² MICS indicator 8.14 - Prevalence of children with one or both parents dead														

The Turkana County MICS included a simple measure of one particular aspect of migration related to what is termed children left behind, i.e. for whom one or both parents have moved abroad. While the amount of literature is growing, the long-term effects of the benefits of remittances versus the potential adverse psycho-social effects are not yet conclusive, as there is somewhat conflicting evidence available as to the effects on children.

The results of the Turkana County MICS presented in Table CP.15 will greatly help fill the data gap on this topic of migration. One percent of children age 0-17 has one or both parents living abroad. Older children are more likely to have one or both parents living abroad.

Table CP.15: Children with parents living abroad							
Percent distribution of children age 0-17 years by residence of parents in another country, Turkana County MICS, 2013/14							
	Percent distribution of children age 0-17 years:					Percentage of children age 0-17 years with at least one parent living abroad ¹	Number of children age 0-17 years
	With at least one parent living abroad			With neither parent living abroad	Total		
	Only mother abroad	Only father abroad	Both mother and father abroad				
Total	0.2	0.4	0.4	99.0	100.0	1.0	3,699
Sex							
Male	0.4	0.2	0.4	98.9	100.0	1.1	1,876
Female	0.0	0.5	0.4	99.0	100.0	1.0	1,823
Area							
Urban	0.4	0.6	0.6	98.3	100.0	1.7	1,975
Rural	0.0	0.1	0.2	99.7	100.0	0.3	1,724
Age group							
0-4	0.1	0.4	0.0	99.5	100.0	0.5	1,095
5-9	0.3	0.3	0.1	99.3	100.0	0.7	1,096
10-14	0.1	0.4	0.7	98.8	100.0	1.2	1,018
15-17	0.7	0.2	1.7	97.3	100.0	2.7	489
Wealth index quintile							
Poorest	0.4	0.0	0.0	99.6	100.0	0.4	742
Second	0.0	0.0	0.0	100.0	100.0	0.0	763
Middle	0.0	1.0	0.0	99.0	100.0	1.0	746
Fourth	0.5	0.6	2.2	96.8	100.0	3.2	762
Richest	0.4	0.2	0.0	99.4	100.0	0.6	686
Ethnicity of household head							
Turkana	0.1	0.0	0.0	99.9	100.0	0.1	3,100
Other ethnic group	1.0	2.3	2.8	93.9	100.0	6.1	588

¹ MICS indicator 8.15 - Children with at least one parent living abroad

11. HIV/AIDS and Sexual Behaviour

HIV prevalence in Kenya has declined and stabilised over the years. A trend analysis starting from 1990s how that prevalence in the general population reached a peak of 10.5 percent in 1995-96, after which it declined by about 40 percent to reach approximately 6.0 percent in 2013.¹¹³ The decline can partly be attributed to high AIDS related mortality. The prevalence has remained relatively stable since 2003 and is attributed to the rapid scale up of anti-retroviral therapy (ART) and reduction in the number of new infections that occurred during this period.

HIV and AIDS programmes in the country are guided by policies and strategies that include the Kenya National HIV/AIDS Strategic Plan; Condom Policy and Strategy, 2001; HIV and AIDS Prevention and Control ACT, 2006; HIV and AIDS policy at the workplace, 2007; Greater Involvement of People Living with HIV and AIDS (GIPA) Guidelines, 2007; Male Circumcision Policy, 2008; Reproductive Health Communication Strategy Implementation Guide for Family Planning, Adolescent and Youth Sexuality and Reproductive Health Rights, and Maternal, Neonatal, and Child Health 2010-2012; Education Sector Policy on HIV and AIDS, 2013 and many more. The current Kenya AIDS Strategic Framework - KASF 2014/15-2018/19 addresses the drivers of the HIV epidemic and builds on achievements of the previous country strategic plans to achieve its goals of contributing to the country's Vision 2030 through universal access to comprehensive HIV prevention, treatment and care.¹¹⁴

11.1 Knowledge about HIV Transmission and Misconceptions about HIV

One of the most important pre-requisites for reducing the rate of HIV infection is accurate knowledge of how HIV is transmitted and strategies for preventing transmission. Correct information is the first step towards raising awareness. Misconceptions about HIV are common and can confuse adolescents and young people and hinder prevention efforts.

The UN General Assembly Special Session on HIV/AIDS (UNGASS) called on governments to improve the knowledge and skills of young people to protect themselves from HIV. The indicators to measure this goal as well as the Millennium Development Goal (MDG) of reducing HIV infections by half include improving the level of knowledge of HIV and its prevention, and changing behaviours to prevent further spread of the disease. HIV module(s) were administered to women and men 15-49 years of age. Please note that the questions in this module often refer to "the AIDS virus". This terminology is used strictly as a method of data collection to aid respondents, preferred over the correct terminology of "HIV" that is used here in reporting the results, where appropriate.

One indicator which is both an MDG and the Global AIDS Response Progress Reporting (GARPR; formerly UNGASS) indicator is the percentage of young people who have comprehensive and correct knowledge of HIV prevention and transmission. This is defined as 1) knowing that consistent use of a condom during sexual intercourse and having just one uninfected faithful partner can reduce the chance of getting HIV, 2) knowing that a healthy-looking person can have HIV, and 3) rejecting the two most common local misconceptions about transmission/prevention of HIV. In the Turkana County MICS, all women who have heard of AIDS were asked questions on all three components and the results are detailed in Table HA.1.

¹¹³Government of Kenya 2014. Kenya AIDS Response Progress Report 2014 – Progress Towards Zero

¹¹⁴http://www.nacc.or.ke/index.php?option=com_content&view=article&id=189&Itemid=130

Almost all women age 15-49 years (98 percent) have heard of AIDS. However, the percentage of those who know the two main ways of preventing HIV transmission – having only one faithful uninfected partner and using a condom every time- is 35 percent, with about 76 percent knowing of having one faithful uninfected sex partner and 38 percent knowing of using a condom every time.

People who have comprehensive knowledge about HIV prevention include those who know of the two main ways of HIV prevention (having only one faithful uninfected partner and using a condom every time). Those who know that a healthy looking person can be HIV-positive and those who reject the two most common misconceptions. Comprehensive knowledge of HIV prevention methods and transmission is fairly low although there are differences by area, age and by woman's education. Overall, 29 percent of women have comprehensive knowledge: the level differs by place of residence (34 percent in urban and 20 percent in rural areas). Comprehensive knowledge is 35 percent for both age group 15-19 years and 20-24 years and is the highest compared to other age groups. Comprehensive knowledge is higher among women age 15-49 years who never married/in union (38 percent) than for women ever married/in union (25 percent), higher for women with secondary or higher education (39 percent) compared to those with no education (22 percent), and increases with household wealth from 16 percent in the poorest households to 39 percent in the richest.

Table HA.1: Knowledge about HIV transmission, misconceptions about HIV, and comprehensive knowledge about HIV transmission (women)

Percentage of women age 15-49 years who know the main ways of preventing HIV transmission, percentage who know that a healthy looking person can be HIV-positive, percentage who reject common misconceptions, and percentage who have comprehensive knowledge about HIV transmission, Turkana County MICS, 2013/14

	Percentage who have heard of AIDS	Percentage who know transmission can be prevented by:			Percentage who know that a healthy looking person can be HIV-positive	Percentage who know that HIV cannot be transmitted by:			Percentage who reject the two most common misconceptions and know that a healthy looking person can be HIV-positive	Percentage with comprehensive knowledge ¹	Number of women age 15-49 years
		Having only one faithful uninfected sex partner	Using a condom every time	Both		Mosquito bites	Supernatural means	Sharing food with someone with HIV			
Total	98.2	75.6	38.4	34.8	81.4	71.5	88.0	81.4	59.2	28.8	1,104
Area											
Urban	99.6	79.6	45.5	40.5	84.2	80.2	93.4	90.0	67.1	34.1	683
Rural	96.1	69.0	26.9	25.5	76.9	57.3	79.2	67.6	46.4	20.4	421
Age											
15-24 ¹	98.3	78.0	45.7	40.8	82.8	80.3	90.4	86.5	66.2	34.7	461
15-19	98.0	78.4	46.2	40.7	81.8	83.0	91.5	85.2	67.6	34.6	252
20-24	98.6	77.5	45.1	40.9	84.2	77.0	89.0	88.0	64.5	34.9	209
25-29	99.0	71.3	34.9	32.1	82.3	64.7	89.2	77.8	54.7	27.3	210
30-39	98.4	76.5	36.2	33.4	80.4	70.6	86.2	80.7	58.5	26.8	277
40-49	96.8	72.4	25.3	23.2	77.7	56.3	82.7	72.6	45.9	17.2	156
Marital status											
Ever married/in union	98.4	74.3	34.7	31.3	80.8	67.0	87.7	79.5	55.4	25.3	787
Never married/in union	97.9	78.5	47.5	43.2	82.8	82.8	88.7	86.0	68.6	37.5	314
Education											
None	97.2	72.2	29.5	26.8	75.9	59.3	82.3	72.5	46.7	21.7	622
Primary	99.7	80.5	52.1	47.1	86.1	82.9	95.3	91.9	70.3	37.5	296
Secondary+	99.3	79.0	46.2	41.8	92.4	94.0	95.7	94.5	83.2	39.0	186
Wealth index quintile											
Poorest	95.6	56.4	19.7	18.8	69.7	45.0	75.3	56.9	35.3	15.8	169
Second	95.7	74.7	30.6	28.7	77.3	55.3	79.9	68.8	43.7	21.2	191
Middle	99.6	79.8	41.7	37.0	83.8	77.1	91.0	85.8	63.1	29.1	226


Fourth	100.0	78.7	44.6	37.9	83.3	84.2	94.0	92.0	69.3	32.5	249
Richest	98.9	81.8	47.0	44.3	87.9	83.1	93.8	92.3	72.5	38.7	270
Ethnicity of household head											
Turkana	98.0	74.6	37.2	33.9	81.7	69.0	86.3	79.5	57.7	28.5	883
Other ethnic group	99.2	79.5	43.1	38.1	80.8	81.7	95.2	89.5	65.5	30.7	218

¹MICS indicator 9.1; MDG indicator 6.3 - Knowledge about HIV prevention among young women

Table HA.1 also presents the percentage of women who correctly identified misconceptions concerning HIV. The indicator is based on the two most common and relevant misconceptions in Turkana County.

Overall, 59 percent of women age 15-49 years reject the two most common misconceptions and know that a healthy-looking person can be HIV-positive. The proportion of women who know that HIV cannot be transmitted by mosquito bites, supernatural means or by sharing food with someone with HIV are 72 percent, 88 percent and 81 percent, respectively. Eighty-one percent of women know that a healthy-looking person can be HIV-positive. Some of these indicators are also presented graphically in Figure HA.1.

Figure HA.1: Women with comprehensive knowledge of HIV transmission, Turkana County MICS, 2013/14



11.2 Knowledge of mother-to-child HIV transmission (MTCT)

In Kenya, infants infected with HIV annually due to mother-to-child transmission declined from 44,000 in 2000 to 12,940 in 2013.¹¹⁵ To guide interventions on mother to child transmission of HIV, Kenya developed Guidelines for Prevention of Mother to Child Transmission (PMTCT) of HIV and AIDS, 2012 and the Kenya Strategic Framework for EMTCT, 2012. The Guidelines complement Kenya’s National Health Sector Strategic Plan II (NHSSP II) and the Kenya National AIDS Strategic Plan (KNASP III) 2009-2013 which focuses on the priority areas of prevention of new infections, improving the quality of life

¹¹⁵ Ministry of Health, 2014. Kenya HIV Estimates

of people infected and affected by HIV and AIDS, and mitigation of the social and economic impact of the infection (ibid). The strategies and guidelines are in line with the WHO PMTCT Strategic Vision 2010-2015 and the 2010 WHO Guidelines on Prevention of Mother-to-Child Transmission (PMTCT) programmes.

Knowledge of mother-to-child transmission of HIV is an important first step for women to seek HIV testing when they are pregnant to avoid infection of the baby. Women and men should know that HIV can be transmitted during pregnancy, during delivery, and through breastfeeding. The level of knowledge among women age 15-49 concerning mother-to-child transmission is presented in Tables HA.2. Overall, 91 percent of women know that HIV can be transmitted from mother-to-child by at least one of the three means. The percentage of women who know all three ways of mother-to-child transmission is 22 percent, while seven percent of women do not know of any specific way.

Table HA.2: Knowledge of mother-to-child HIV transmission (women)

Percentage of women age 15-49 years who correctly identify means of HIV transmission from mother to child, Turkana County MICS, 2013/14							
	Percentage of women age 15-49 who have heard of AIDS and:						Number of women age 15-49 years
	Know HIV can be transmitted from mother to child:					Do not know any of the specific means of HIV transmission from mother to child	
	During pregnancy	During delivery	By breastfeeding	By at least one of the three means	By all three means ¹		
Total	28.0	77.2	86.1	90.8	22.3	7.4	1,104
Area							
Urban	28.3	81.7	88.1	94.1	22.9	5.5	683
Rural	27.4	69.9	82.7	85.5	21.4	10.6	421
Age group							
15-24	25.5	76.4	86.8	90.6	20.3	7.7	461
15-19	22.7	75.2	83.8	89.3	16.1	8.7	252
20-24	28.9	77.9	90.4	92.1	25.4	6.5	209
25-29	32.5	84.5	91.2	94.4	30.1	4.6	210
30-39	28.9	75.8	85.0	91.0	21.8	7.4	277
40-49	27.5	72.0	79.1	86.5	18.7	10.4	156
Marital status							
Ever married/in union	30.2	77.8	87.4	92.2	24.1	6.2	787
Never married/in union	22.6	75.9	83.3	87.9	17.9	10.0	314
Education							
None	29.0	71.4	83.0	87.4	23.0	9.8	622
Primary	32.0	78.6	87.3	93.0	25.3	6.7	296
Secondary+	18.1	94.2	94.7	98.7	15.2	0.7	186
Wealth index quintile							
Poorest	33.4	59.0	78.2	82.2	22.9	13.5	169
Second	31.7	73.6	84.4	87.2	27.2	8.5	191
Middle	27.2	83.7	89.5	94.4	23.0	5.2	226
Fourth	26.3	79.0	86.6	92.9	21.0	7.1	249
Richest	24.1	83.9	88.8	94.0	19.2	4.9	270
Ethnicity of household head							
Turkana	28.3	76.2	85.4	90.3	22.3	7.7	883
Other ethnic group	27.1	81.0	88.7	93.1	22.7	6.2	218

¹ MICS indicator 9.2 - Knowledge of mother-to-child transmission of HIV

11.3 Accepting Attitudes toward People Living with HIV

The indicators on attitudes toward people living with HIV measure stigma and discrimination in the community. Stigma and discrimination are considered low if respondents report an accepting attitude on the following four questions: 1) would care for a family member with AIDS in own home; 2) would buy fresh vegetables from a vendor who is HIV-positive; 3) thinks that a female teacher who is HIV-positive should be allowed to teach in school; and 4) would not want to keep it a secret if a family member is HIV-positive.

Table HA.3: Accepting attitudes toward people living with HIV (women)


Percentage of women age 15-49 years who have heard of AIDS who express an accepting attitude towards people living with HIV, Turkana County MICS, 2013/14

	Percentage of women who:						Number of women age 15-49 who have heard of AIDS
	Are willing to care for a family member with AIDS in own home	Would buy fresh vegetables from a shopkeeper or vendor who is HIV-positive	Believe that a female teacher who is HIV-positive and is not sick should be allowed to continue teaching	Would not want to keep secret that a family member is HIV-positive	Agree with at least one accepting attitude	Express accepting attitudes on all four indicators ¹	
Total	84.1	61.6	74.8	33.3	96.8	16.7	1,085
Area							
Urban	91.6	68.6	81.2	31.3	98.2	19.4	680
Rural	71.4	49.6	64.1	36.7	94.5	12.2	404
Age							
15-24	87.5	62.9	77.4	31.4	97.7	15.6	453
15-19	88.9	63.2	76.0	29.9	98.1	14.4	247
20-24	85.7	62.5	79.0	33.1	97.3	17.2	206
25-29	81.5	62.3	73.5	37.6	97.2	20.2	208
30-39	83.1	59.5	73.6	31.8	96.0	15.7	273
40-49	79.2	60.2	70.9	36.0	95.0	17.0	151
Marital status							
Ever married/in union	82.4	60.4	73.7	33.2	96.5	17.0	774
Never married/in union	88.2	64.4	77.3	33.5	97.5	16.2	308
Education							
None	76.0	51.4	65.3	36.6	94.7	15.0	605
Primary	90.6	70.2	83.7	30.7	99.3	18.5	295
Secondary+	100.0	81.1	91.7	26.6	100.0	19.4	185
Wealth index quintile							
Poorest	62.9	33.7	48.5	41.0	91.5	10.1	161
Second	76.2	52.8	66.2	32.0	93.6	15.7	183
Middle	86.1	74.7	80.3	32.4	97.8	18.1	225
Fourth	90.6	64.7	82.1	29.0	98.8	17.8	249
Richest	94.4	70.4	85.1	34.4	99.5	19.2	267
Ethnicity of household head							
Turkana	82.7	59.9	73.5	34.3	96.4	16.3	866
Other ethnic group	89.4	68.1	80.1	29.6	98.5	18.5	217

¹ MICS indicator 9.3 - Accepting attitudes towards people living with HIV

Table HA.3 and Figure HA.2 present the attitudes of women age 15-49 years towards people living with HIV. Ninety-seven percent of women who have heard of AIDS agree with at least one accepting statement. The most common accepting attitude is willingness to care for a family member with AIDS in own home (84 percent). The proportion of women who express accepting attitudes towards all four indicators declines to only 17 percent.

Figure HA.2: Accepting attitudes toward people living with HIV/AIDS, Turkana County MICS, 2013/14



11.4 Knowledge of a Place for HIV Counselling and Testing during Antenatal Care

Another important indicator is the knowledge of where to be tested for HIV and use of such services. In order to protect themselves and to prevent infecting others, it is important for individuals to know their HIV status. Knowledge of own status is also a critical factor in the decision to seek treatment.

Results related to knowledge of a facility for HIV testing and whether a person had ever been tested is presented in Tables HA.4. Eighty-eight percent of women age 15-49 years know of a place where to be tested, while 78 percent have been tested. Fifty-seven percent of women know the result of their most recent test. Knowledge of a place to get tested is 94 percent in urban areas and 79 percent in rural areas. Knowledge of a place to get tested ranges from 84 percent to 97 percent by education of the woman.

The proportion of women age 15-49 years who had been tested within the last 12 months preceding the survey is 58 percent, while 53 percent were tested within the last 12 months and know the result.

Urban areas (64 percent) have a higher proportion of women who have been tested in the last 12 months preceding the survey and know their results compared to rural areas (35 percent). The proportion of women who have been tested in the last 12 months and know their results ranges from a high of 60 percent in the 20-24 years age group to a low of 43 percent for those in the 40-49 years category.

Table HA.4: Knowledge of a place for HIV testing (women)							
Percentage of women age 15-49 years who know where to get an HIV test, percentage who have ever been tested, percentage who have ever been tested and know the result of the most recent test, percentage who have been tested in the last 12 months, and percentage who have been tested in the last 12 months and know the result, Turkana County MICS, 2013/14							
	Percentage of women who:					Number of women age 15-49 years	
	Know a place to get tested ¹	Have ever been tested	Have ever been tested and know the result of the most recent test	Have been tested in the last 12 months	Have been tested in the last 12 months and know the result ^{2,3}		
Total	88.1	77.9	57.3	57.8	53.0	1,104	
Area							
Urban	93.8	83.8	67.3	67.6	64.2	683	
Rural	78.8	68.4	40.9	41.8	34.8	421	
Age							
15-24	88.5	71.8	60.3	60.3	57.0	461	
15-19	88.8	64.9	55.1	55.1	54.7	252	
20-24	88.2	80.1	66.6	66.6	59.8	209	
25-29	92.8	89.0	68.3	70.6	58.0	210	
30-39	87.6	83.0	51.4	51.6	48.2	277	
40-49	81.3	72.2	43.7	44.1	43.0	156	
Age and sexual activity in the last 12 months							
Sexually active	89.8	85.6	63.3	63.9	57.3	697	
15-24 ³	92.8	88.7	74.0	74.0	67.7	221	
15-19	94.1	89.8	77.7	77.7	76.4	81	
20-24	92.1	88.1	72.0	72.0	62.7	140	
25-49	88.4	84.1	58.3	59.2	52.5	476	
Sexually inactive	85.2	64.9	46.9	47.2	45.6	407	
Marital status							
Ever married/in union	88.0	83.3	58.4	59.0	52.4	787	
Never married/in union	88.1	64.4	54.4	54.8	54.4	314	
Education							
None	84.0	75.4	50.4	51.4	45.4	622	
Primary	91.2	76.9	63.5	63.5	59.0	296	
Secondary+	96.7	88.1	70.1	70.1	68.8	186	
Wealth index quintile							
Poorest	71.4	57.5	29.8	30.9	24.4	169	
Second	83.8	77.1	47.7	48.9	42.6	191	
Middle	93.3	82.8	62.3	63.1	56.0	226	
Fourth	92.6	84.4	67.4	67.4	63.4	249	
Richest	93.0	81.4	67.6	67.6	66.2	270	
Ethnicity of household head							
Turkana	87.4	77.9	56.2	56.8	51.7	883	
Other ethnic group	90.8	78.1	61.6	62.0	58.5	218	

¹ MICS indicator 9.4 - Women who know where to be tested for HIV

² MICS indicator 9.5 - Women who have been tested for HIV and know the results

³ MICS indicator 9.6 - Sexually active young women who have been tested for HIV and know the results

Among women who had given birth within the two years preceding the survey, the percentage who received counselling and HIV testing during antenatal care is presented in Table HA.5. Eighty-four percent of women age 15-49 with a live birth in the last two years preceding the survey received HIV counselling during ANC, 85 percent were offered an HIV test and were tested for HIV. Eighty four percent were offered an HIV test and were tested during ANC and received the results, and 78 percent received HIV counselling, offered an HIV test, accepted and received the results. A larger percentage of women in urban areas (89 percent) received HIV counselling, HIV testing, and received the results during ANC than those in rural areas (79 percent).

Table HA.5: HIV counselling and testing during antenatal care

Percentage of women age 15-49 with a live birth in the last 2 years who received antenatal care from a health professional during the last pregnancy, percentage who received HIV counselling, percentage who were offered and tested for HIV, percentage who were offered, tested and received the results of the HIV test, and percentage who received counselling and were offered, accepted and received the results of the HIV test, Turkana County MICS, 2013/14

	Percentage of women who:					Number of women age 15-49 years with a live birth in the last 2 years
	Received antenatal care from a health care professional for last pregnancy	Received HIV counselling during antenatal care ¹	Were offered an HIV test and were tested for HIV during antenatal care	Were offered an HIV test and were tested for HIV during antenatal care, and received the results ²	Received HIV counselling, were offered an HIV test, accepted and received the results	
Total	93.4	83.6	85.0	83.7	77.9	387
Area						
Urban	98.3	90.2	89.3	88.5	83.5	199
Rural	88.2	76.7	80.4	78.7	72.0	188
Age						
15-24	97.2	85.7	85.3	85.3	81.0	112
25-29	92.0	80.3	85.0	83.7	74.3	127
30-39	93.8	89.0	85.2	83.0	81.0	119
40-49	(83.7)	(68.7)	(82.8)	(80.8)	(68.7)	29
Education						
None	90.5	80.8	83.2	82.0	75.7	268
Primary	100.0	90.6	88.7	86.5	82.2	82
Secondary+	(100.0)	(88.7)	(89.9)	(89.9)	(84.2)	38
Wealth index quintile						
Poorest	84.1	72.1	71.8	71.0	66.2	69
Second	91.2	78.0	83.4	82.2	73.3	86
Middle	96.1	88.5	90.5	86.5	80.8	84
Fourth	97.6	92.9	89.9	89.9	87.3	85
Richest	97.5	85.1	87.7	87.7	80.4	63
Ethnicity of household head						
Turkana	92.5	82.8	83.3	81.8	76.3	327
Other ethnic group	98.3	88.1	94.0	94.0	86.2	59
¹ MICS indicator 9.7 - HIV counselling during antenatal care						
² MICS indicator 9.8 - HIV testing during antenatal care						
() Figures that are based on 25-49 unweighted cases						

11.5 Sexual Behaviour Related to HIV Transmission

Promoting safer sexual behaviour is critical in reducing HIV prevalence. The use of condoms during sex, especially when non-regular or multiple partners are involved, is particularly important for reducing the spread of HIV. A set of questions was administered to all women age 15-49 to assess their risk of HIV infection.

Table HA.6: Sex with multiple partners (women)

Percentage of women age 15-49 years who ever had sex, percentage who had sex in the last 12 months, percentage who had sex with more than one partner in the last 12 months, mean number of sexual partners in lifetime for women who have ever had sex, and among those who had sex with multiple partners in the last 12 months, the percentage who used a condom at last sex, Turkana County MICS, 2013/14

	Percentage of women who:			Number of women age 15-49 years	Mean number of sexual partners in lifetime	Number of women age 15-49 years who have ever had sex
	Ever had sex	Had sex in the last 12 months	Had sex with more than one partner in last 12 months ¹			
Total	82.2	63.2	1.8	1,104	1.3	907
Area						
Urban	79.7	59.0	2.3	683	1.4	545
Rural	86.1	70.0	1.2	421	1.2	363
Age						
15-24	58.7	48.0	2.7	461	1.4	270
15-19	40.5	32.1	2.8	252	1.3	102
20-24	80.5	67.2	2.7	209	1.4	168
25-29	97.8	84.9	0.9	210	1.3	205
30-39	99.7	76.9	1.9	277	1.3	277
40-49	99.2	54.3	0.4	156	1.3	155
Marital status						
Ever married/in union	100.0	78.2	1.3	787	1.3	787
Never married/in union	37.3	26.3	3.3	314	1.7	117
Education						
None	89.7	68.3	1.8	622	1.3	558
Primary	68.3	51.0	2.5	296	1.4	202
Secondary+	79.1	65.3	0.9	186	1.4	147
Wealth index quintile						
Poorest	84.8	67.5	0.9	169	1.2	143
Second	86.4	65.4	1.4	191	1.3	165
Middle	84.2	57.9	2.7	226	1.4	190
Fourth	82.5	63.2	2.4	249	1.3	205
Richest	75.6	63.3	1.5	270	1.4	204
Ethnicity of household head						
Turkana	84.4	66.6	1.5	883	1.3	745
Other ethnic group	73.0	49.3	3.4	218	1.4	159

¹ MICS indicator 9.12 - Multiple sexual partnerships

² MICS indicator 9.13 - Condom use at last sex among people with multiple sexual partnerships (this indicator could not be produced due to insufficient sample size)

As shown in Table HA.6, two percent of women 15-49 years of age reported that they had had sex with more than one partner in the last 12 months. Overall, the mean number of lifetime sexual partners is 1.3.¹¹⁶

11.6 HIV Indicators for Young Women

In many countries, over half of new adult HIV infections are among young people of age 15-24 years thus a change in behaviour among members of this age group is especially important to reduce new infections.

Table HA.7 summarizes information on key HIV indicators for young women in Turkana. Thirty-five percent of young women have comprehensive knowledge. Young women who know of three means of HIV transmission from mother-to-child are 20 percent and 89 percent have knowledge of a place to get tested. Young urban women are more knowledgeable (36 percent) than their rural counterparts (30 percent). Comprehensive knowledge increases with household wealth from 18 percent among the poorest to 44 percent in the richest.

Overall, 68 percent of young women in this age group, who were sexually active, had been tested for HIV in the last 12 months and know the result. There are disparities by place of residence, and marital status. The percentage of sexually active women who had been tested for HIV in the past 12 months and know the result is 78 percent in urban areas and 34 percent in rural areas. The proportion is high among never married/in union (75 percent) compared with ever married/in union (64 percent).

¹¹⁶ The percentage of women who had more than one sexual partner in the last 12 months reporting that a condom was used the last time they had sex could not be included in the table due to small number of cases reported.

Table HA.7: Key HIV and AIDS indicators (young women)

Percentage of women age 15-24 years by key HIV and AIDS indicators, Turkana County MICS, 2013/14

	Percentage of women age 15-24 years who:										Number of women age 15-24 years who have heard of AIDS
	Have comprehensive knowledge ¹	Know all three means of HIV transmission from mother to child	Know a place to get tested for HIV	Have ever been tested and know the result of the most recent test	Have been tested for HIV in the last 12 months and know the result	Had sex in the last 12 months	Number of women age 15-24 years	Percentage of sexually active young women who have been tested for HIV in the last 12 months and know the result ²	Number of women age 15-24 years who had sex in the last 12 months	Percentage who express accepting attitudes towards people living with HIV on all four indicators ^a	
Total	34.7	20.3	88.5	60.3	57.0	48.0	461	67.7	221	15.6	453
Area											
Urban	36.4	21.5	94.3	68.2	65.7	49.7	345	77.5	171	18.1	345
Rural	29.7	16.7	71.3	36.7	31.3	43.1	116	33.9	50	7.9	108
Age											
15-19	34.6	16.1	88.8	55.1	54.7	32.1	252	76.4	81	14.4	247
15-17	33.9	16.6	86.7	48.8	48.8	25.2	170	(78.1)	43	19.0	166
18-19	36.0	15.0	93.2	68.4	67.1	46.6	82	(74.5)	38	4.8	81
20-24	34.9	25.4	88.2	66.6	59.8	67.2	209	62.7	140	17.2	206
20-22	34.2	27.9	93.2	69.5	65.3	69.5	139	67.3	97	14.6	139
23-24	36.2	20.4	78.3	60.7	49.0	62.4	70	(52.4)	43	22.4	67
Marital status											
Ever married/in union	33.7	24.4	89.6	70.8	62.2	86.7	176	64.4	153	18.1	174
Never married/in union	35.3	17.8	87.9	53.8	53.8	24.1	285	75.1	69	14.1	278
Education											
None	25.5	18.9	77.7	50.9	45.0	48.3	152	56.3	74	8.6	145
Primary	39.1	25.1	92.4	63.9	61.2	42.2	196	74.8	83	20.3	195
Secondary+	39.7	13.8	96.5	66.8	66.2	57.9	112	71.6	65	16.6	112
Wealth index quintile											
Poorest	17.5	18.1	55.9	22.2	20.5	38.0	44	(*)	17	6.3	40
Second	29.4	24.4	80.7	50.6	44.3	30.9	45	(*)	14	(12.0)	42
Middle	31.4	21.7	93.0	66.4	61.0	50.4	100	64.7	50	16.1	99
Fourth	34.4	20.7	92.2	70.0	65.8	57.8	127	77.0	73	20.0	127

Richest	44.2	18.4	94.7	62.3	61.8	46.2	145	77.0	67	15.2	145
Ethnicity of household head											
Turkana	35.0	20.6	88.7	61.4	57.4	51.3	350	66.6	180	14.9	342
Other ethnic group	33.8	19.3	87.9	56.8	55.8	37.4	110	(72.3)	41	18.0	110
¹ MICS indicator 9.1; MDG indicator 6.3 - Knowledge about HIV prevention among young women											
² MICS indicator 9.6 - Sexually active young women who have been tested for HIV and know the results											
^a Refer to Table HA.3 for the four indicators.											
() Figures that are based on 25-49 unweighted cases											
(*) Figures that are based on fewer than 25 unweighted cases											

Certain behaviour may create, increase, or perpetuate risk of exposure to HIV. For this young age group, such behaviour includes sex at an early age and women having sex with older men.

Table HA.8 shows results on sexual behaviour of young women age 15-24 years. Overall, eight percent of young women reported ever having sex before age 15. Further, three percent of young women had sex with more than one partner in the last 12 months. On the other hand, 17 percent of the young women who had sex in the last 12 months reported that it involved a non-marital, non-cohabiting partner; of those only 31 percent of women used a condom the last time. About 18 percent of women age 15-24 years had sex with a man 10 or more years older in the last 12 months. Analysis on the percentage of women age 15-24 years reporting that a condom was used the last time they had sex could not be undertaken due to insufficient sample size.

Table HA.8: Key sexual behaviour indicators (young women)


Percentage of women age 15-24 years by key sexual behaviour indicators, Turkana County MICS, 2013/14

	Percentage of women age 15-24 years who:			Number of women age 15-24 years	Percentage of women who never had sex ²	Number of never-married women age 15-24 years	Percentage of women age 15-24 years who in the last 12 months had sex with:		Number of women age 15-24 years who had sex in the last 12 months	Percentage reporting the use of a condom during the last sexual intercourse with a non-marital, non-cohabiting partner in the last 12 months ⁵	Number of women age 15-24 years who had sex with a non-marital, non-cohabiting partner in last 12 months
	Had sex before age 15 ¹	Ever had sex	Had sex with more than one partner in last 12 months				A man 10 or more years older ³	A non-marital, non-cohabiting partner ⁴			
Total	8.2	58.7	2.7	461	66.9	285	17.6	17.0	221	31.2	78
Area											
Urban	8.9	61.0	3.0	345	62.6	215	17.3	19.3	171	33.3	67
Rural	6.2	51.7	1.8	116	80.1	70	18.7	10.0	50	(*)	12
Age											
15-19	10.0	40.5	2.8	252	69.1	217	13.3	19.7	81	(36.0)	50
15-17	11.8	31.3	4.1	170	71.4	163	(14.5)	(21.4)	43	(43.0)	36
18-19	6.2	59.8	0.0	82	62.0	53	(12.1)	(16.1)	38	(*)	13
20-24	6.0	80.5	2.7	209	59.8	68	20.0	13.7	140	(23.0)	29
20-22	7.2	78.1	1.9	139	60.0	51	19.7	15.8	97	(*)	22
23-24	3.7	85.4	4.2	70	(*)	17	(20.9)	(9.5)	43	(*)	7
Marital status											
Ever married/in union	9.9	100.0	2.9	176	na	na	23.1	8.6	153	(*)	15
Never married/in union	7.2	33.1	2.7	285	66.9	285	5.3	22.1	69	35.9	63
Education											
None	8.5	60.1	3.4	152	76.2	80	26.3	12.6	74	(*)	19
Primary	9.1	53.2	3.0	196	71.8	128	18.3	14.2	83	(*)	28
Secondary+	6.2	66.4	1.5	112	49.0	77	6.9	27.8	65	(*)	31
Wealth index quintile											
Poorest	7.8	46.9	2.3	44	(82.8)	28	(*)	(*)	17	(*)	4
Second	4.9	45.2	1.2	45	(82.6)	30	(*)	(*)	14	(*)	5
Middle	7.1	64.2	3.4	100	55.7	64	15.5	18.8	50	(*)	19

Fourth	9.4	67.5	3.5	127	69.5	59	14.2	16.8	73	(*)	21
Richest	9.0	54.9	2.2	145	63.4	103	19.6	20.7	67	(*)	30
Ethnicity of household head											
Turkana	10.5	62.2	2.4	350	62.5	212	18.7	17.4	180	32.3	61
Other ethnic group	0.9	47.4	3.7	110	79.5	73	(12.8)	(15.6)	41	(*)	17
¹ MICS indicator 9.10 - Sex before age 15 among young women											
² MICS indicator 9.9 - Young women who have never had sex											
³ MICS indicator 9.11 - Age-mixing among sexual partners											
⁴ MICS indicator 9.14 - Sex with non-regular partners											
⁵ MICS indicator 9.15; MDG indicator 6.2 - Condom use with non-regular partners											
na: not applicable											
() Figures that are based on 25-49 unweighted cases											
(*) Figures that are based on fewer than 25 unweighted cases											

Figure HA.3 brings together two critical behaviours that are known to increase the risk of HIV infection, sex before age 15, and sex with multiple partners, from tables HA.8 and HA.6.

Figure HA.3: Sexual behaviour that increases the risk of HIV infection, young people age 15-24, Turkana County MICS, 2013/14



11.7 Orphans

While the number of children orphaned due to AIDS has stabilized globally since 2009, efforts to mitigate the impact of AIDS on households, communities, and children continue to be intensified by national programmes and global partners. Children who are orphaned may be at increased risk of neglect or exploitation when the parents are not available to assist them. Monitoring the variations in different outcomes for orphans and comparing them to their peers gives us a measure of how well communities and governments are responding to their needs. Please refer to Table CP.14 on page 155 for detailed information on living conditions of children and overall prevalence of orphanhood.

Table HA.9 presents information on the orphanhood status of children age 10-14 years, and their school attendance. About seven percent of children age 10-14 years in Turkana County are orphans. Eighty-six percent of orphans are attending school, compared with a 72 percent attendance amongst non-orphan children of the same age group who are living with at least one parent. The orphans to non-orphans school attendance ratio is 1.19 which suggests that orphans are not disadvantaged in relation to non-orphans.

Table HA.9: School attendance of orphans and non-orphans

School attendance of children age 10-14 years by orphanhood, Turkana County MICS, 2013/14

	Percentage of children whose mother and father have died (orphans)	Percentage of children whose parents are still alive and who are living with at least one parent (non-orphans)	Number of children age 10-14 years	Percentage of children whose mother and father have died (orphans) and are attending school	Total number of orphan children age 10-14 years	Percentage of children whose parents are still alive, who are living with at least one parent (non-orphans), and who are attending school	Total number of non-orphan children age 10-14 years	Orphans to non-orphans school attendance ratio ¹
Total	6.9	61.8	1,018	85.9	70	72.2	630	1.19
Sex								
Male	7.5	63.4	508	(85.9)	38	73.9	322	(1.16)
Female	6.3	60.3	511	(86.0)	32	70.4	308	(1.22)
Area								
Urban	10.3	53.6	550	(93.1)	56	95.2	295	(0.98)
Rural	3.0	71.5	468	(*)	14	51.8	335	(*)
¹ MICS indicator 9.16; MDG indicator 6.4 - Ratio of school attendance of orphans to school attendance of non-orphans								
See Table CP.14 for further overall results related to children's living arrangements and orphanhood								
() Figures that are based on 25-49 unweighted cases								
(*) Figures that are based on fewer than 25 unweighted cases								

12. Access to Mass Media and Use of Information/Communication Technology

The Government of Kenya recognizes the role of Information and Communication Technology (ICT) in the social and economic development of the nation and has developed a national ICT Policy based on the Economic Recovery Strategy for Wealth and Employment Creation (2003-2007). In the National ICT Policy (2006), the Government's vision is to make Kenya 'a prosperous ICT-driven society'.^{117, 118}

The Turkana County MICS collected information on exposure to mass media and the use of computers and the internet. Information was collected on exposure to newspapers/magazines, radio and television among women age 15-49 years, while the questions on the use of computers and the use of the internet were asked to young women age 15-24 years. This chapter, therefore, discusses access to mass media and use of ICT.

12.1 Access to Mass Media

The proportion of women who read a newspaper or magazine, listen to the radio and watch television at least once a week is shown in Table MT.1. About nine percent of women in Turkana County read a newspaper or magazine, 19 percent listen to the radio, and 16 percent watch television at least once a week. Overall, 72 percent do not have regular exposure to any of the three media, while 27 percent are exposed to at least one and three percent to all the three types of media on a weekly basis.

Women under age 25 years are more likely than older women to report exposure to all three types of mass media. Strong differentials by area, education and household wealth are observed for exposure to all types of media. Women with secondary or higher education are more likely to have been exposed to all three types of media (16 percent) than women with only primary education (2 percent). Similarly, women from the richest households are more likely to have been exposed to all three types of media (12 percent) than women from the poorest households (0 percent).

¹¹⁷ <http://www1.american.edu/initeb/en6343a/ICT-policy.htm>

¹¹⁸ Ministry of Information and Communications. 2006. National Information and Communications Technology (ICT) Policy.

Table MT.1: Exposure to mass media (women)

Percentage of women age 15-49 years who are exposed to specific mass media on a weekly basis, Turkana County MICS, 2013/14

	Percentage of women age 15-49 years who:						Number of women age 15-49 years
	Read a newspaper at least once a week	Listen to the radio at least once a week	Watch television at least once a week	All three media at least once a week ¹	Any media at least once a week	None of the media at least once a week	
Total	8.5	18.9	15.9	3.2	27.4	72.4	1,104
Age							
15-19	18.6	28.0	25.3	7.2	42.6	57.4	252
20-24	7.9	17.9	19.2	1.8	33.7	65.8	209
25-29	8.0	18.8	14.7	3.5	24.6	75.4	210
30-34	3.1	14.5	8.2	1.9	15.7	83.7	144
35-39	5.0	16.7	7.4	1.9	17.9	82.1	134
40-44	3.1	10.0	14.1	1.4	18.7	81.3	82
45-49	0.0	13.3	9.7	0.0	15.5	84.5	75
Area							
Urban	12.5	28.5	24.1	4.6	41.1	58.8	683
Rural	2.0	3.2	2.5	1.0	5.3	94.5	421
Education							
None	0.0	4.8	3.0	0.0	7.2	92.8	622
Primary	11.3	29.5	21.4	2.3	42.6	57.1	296
Secondary+	32.4	48.9	49.8	15.5	70.9	28.6	186
Wealth index quintile							
Poorest	0.0	0.0	0.0	0.0	0.0	100.0	169
Second	0.0	0.7	0.0	0.0	0.7	99.3	191
Middle	4.0	7.2	5.5	0.8	12.8	87.2	226
Fourth	6.0	16.2	11.7	1.2	26.4	73.2	249
Richest	25.9	55.8	49.5	11.5	76.7	23.0	270
Ethnicity of household head							
Turkana	7.8	19.3	12.4	3.0	25.0	74.9	883
Other ethnic group	11.5	16.9	29.5	4.1	36.8	62.8	218

¹ MICS indicator 10.1 - Exposure to mass media

12.2 Use of Information/Communication Technology

The questions on computer and internet use were asked only to young women age 15-24 years. As shown in Table MT.2, 14 percent of young women age 15-24 years ever used a computer, 11 percent had used a computer during the last 12 months and seven percent used a computer at least once a week during the last month.

Overall, 12 percent of young women age 15-24 years ever used the internet, while a similar proportion had used the internet during the last 12 months. The proportion of young women who use the internet more frequently, at least once a week during the last month, is lower, at 10 percent.

Both computer and internet use during the last 12 months are more widespread among the 20-24 year-old women. Use of a computer and the internet is also strongly associated with place of residence, education and household wealth. None of the women with primary education had used

computer during the last 12 months, while 39 percent of women with secondary or higher education used a computer during the same period. Similarly, higher utilisation of a computer and the internet is observed among young women in the richest households. For example, 17 percent of women living in the wealthiest households used a computer at least once during the month before the survey, and 21 percent of them used the internet at least once a week during the same period and none from the poorest households for both indicators.

Table MT.2: Use of computers and internet (women)

Percentage of young women age 15-24 years who have ever used a computer and the internet, percentage who have used during the last 12 months, and percentage who have used at least once weekly during the last one month, Turkana County MICS, 2013/14

	Percentage of women age 15-24 years who have:						Number of women age 15-24 years
	Ever used a computer	Used a computer during the last 12 months ¹	Used a computer at least once a week during the last one month	Ever used the internet	Used the internet during the last 12 months ²	Used the internet at least once a week during the last one month	
Total	13.7	11.2	6.7	12.4	11.8	10.0	461
Age							
15-19	10.9	9.4	5.1	10.4	10.4	8.4	252
20-24	17.2	13.2	8.6	14.8	13.4	12.0	209
Area							
Urban	17.3	14.1	8.7	14.4	13.9	12.3	345
Rural	3.2	2.3	0.9	6.4	5.5	3.2	116
Education							
None	0.0	0.0	0.0	0.0	0.0	0.0	152
Primary	4.1	3.7	1.1	5.2	4.6	4.0	196
Secondary+	49.2	39.3	25.7	41.9	40.3	34.2	112
Wealth index quintile							
Poorest	0.0	0.0	0.0	0.0	0.0	0.0	44
Second	0.0	0.0	0.0	2.5	2.5	2.5	45
Middle	8.9	6.0	2.8	7.5	6.4	5.2	100
Fourth	10.2	8.0	2.9	9.7	9.1	7.1	127
Richest	28.7	24.3	16.9	25.1	24.4	21.4	145
Ethnicity of household head							
Turkana	10.5	8.6	5.1	10.3	9.8	8.3	350
Other ethnic group	23.8	19.2	11.8	19.2	18.1	15.5	110
				¹ MICS indicator 10.2 - Use of computers			
				² MICS indicator 10.3 - Use of internet			

13. Subjective well-being

Subjective perceptions of individuals of their incomes, health, living environments and the like, play a significant role in their lives and can impact their perception of well-being, irrespective of objective conditions such as actual income and physical health status.¹¹⁹ In the MICS, a set of questions were asked to women age 15-24 years to understand how satisfied this group of young people is in different areas of their lives, such as their family life, friendships, school, current job, health, where they live, how they are treated by others, how they look, and their current income.

Life satisfaction is a measure of an individual's perceived level of well-being. Understanding young women's satisfaction in different areas of their lives can help to gain a comprehensive picture of young people's life situations. A distinction can also be made between life satisfaction and happiness. Happiness is a fleeting emotion that can be affected by numerous factors, including day-to-day factors such as the weather, or a recent death in the family. It is possible for a person to be satisfied with job, income, family life, friends, and other aspects of life, but still be unhappy, or vice versa. In addition to the set of questions on life satisfaction, the survey also asked questions about happiness and the respondents' perceptions of a better life.

To assist respondents in answering the set of questions on happiness and life satisfaction, they were shown a card with smiling faces (and not so smiling faces) that corresponded to the response categories (see the Questionnaires in Appendix H) 'very satisfied', 'somewhat satisfied', 'neither satisfied nor unsatisfied', 'somewhat unsatisfied' and 'very unsatisfied'. For the question on happiness, the same scale was used, this time ranging from 'very happy' to 'very unhappy', in the same fashion.

Table SW.1 shows the proportion of young women age 15-24 years, who are very or somewhat satisfied in selected domains. Note that for three domains, satisfaction with school, job and income, the denominators are confined to those who are currently attending school, have a job, and have an income. Of the different domains, young women are the most satisfied with the way they look (98 percent), followed by their health (96 percent), and their family life (90 percent). The percentage of women age 15-24 years who are very or somewhat satisfied with school is 91 percent.¹²⁰

¹¹⁹ OECD. 2013. *OECD Guidelines on Measuring Subjective Well Being*. OECD. <http://dx.doi.org/10.1787/9789264191655-en>

¹²⁰ Four columns assessing the proportion of women age 15-24 years who were very or somewhat satisfied with their job and those who were very or somewhat satisfied with their income, were removed from the table due to the few number of cases reported.

Table SW.1: Domains of life satisfaction (women)

Percentage of women age 15-24 years who are very or somewhat satisfied in selected domains of satisfaction, Turkana County MICS, 2013/14

	Percentage of women age 15-24 years who are very or somewhat satisfied in selected domains:						Percentage of women age 15-24 years who:			Number of women age 15-24 years	Percentage of women age 15-24 years who are very or somewhat satisfied with school	Number of women age 15-24 years attending school
	Family life	Friendships	Health	Living environment	Treatment by others	The way they look	Are attending school	Have a job	Have an income			
Total	89.5	88.6	96.2	87.1	86.0	98.3	40.8	4.1	4.4	461	91.2	186
Age												
15-19	89.2	89.0	95.8	90.0	87.9	98.5	61.2	0.9	2.0	252	91.9	153
20-24	90.0	88.2	96.7	83.6	83.7	98.1	16.0	8.1	7.3	209	(88.3)	33
Area												
Urban	89.2	88.3	96.0	89.6	85.3	98.0	47.3	3.9	4.1	345	91.0	162
Rural	90.6	89.7	97.1	79.4	88.3	99.2	21.0	4.9	5.4	116	(92.6)	24
Marital Status												
Ever married/in union	90.2	89.8	96.5	85.6	85.5	97.2	9.9	8.2	8.8	176	(*)	17
Never married/in union	89.1	87.9	96.0	88.0	86.3	99.0	59.7	1.6	1.8	285	90.9	169
Education												
None	92.2	82.8	98.5	84.1	86.8	98.7	1.8	5.1	4.9	152	(*)	3
Primary	89.2	89.2	94.5	87.9	86.7	97.9	57.1	2.3	2.9	196	94.1	111
Secondary+	86.6	95.3	96.2	89.8	83.7	98.5	64.7	6.0	6.5	112	86.5	73
Wealth index quintile												
Poorest	92.4	84.7	100.0	80.6	85.1	100.0	5.3	4.6	4.6	44	(*)	2
Second	87.9	80.6	92.8	80.4	88.2	98.0	13.4	9.0	7.7	45	(*)	6
Middle	88.2	91.6	95.4	84.3	85.6	97.1	44.4	0.7	1.8	100	(90.9)	44
Fourth	84.2	90.2	96.2	87.0	85.2	99.1	40.5	3.3	2.0	127	(92.3)	50
Richest	94.7	88.8	96.7	93.0	86.6	98.1	57.5	5.6	7.3	145	91.9	83
Ethnicity of household head												
Turkana	91.3	88.0	96.7	88.6	87.6	98.1	37.4	4.2	5.0	350	92.8	130
Other ethnic group	83.9	90.7	94.7	82.2	81.0	99.0	51.6	3.8	2.7	110	(87.7)	56

() Figures that are based on 25-49 unweighted cases
(*) Figures that are based on fewer than 25 unweighted cases

In Table SW.2, proportions of women age 15-24 years with overall life satisfaction are shown. “Life satisfaction” is defined as those who are very or somewhat satisfied with their life overall, and is based on a single question which was asked after the life satisfaction questions on all of the above-mentioned domains, with the exception of the question on satisfaction with income, which was asked later.

In Turkana County, 90 percent of women age 15-24 years are satisfied with their life. The proportions do not vary significantly by age and marital status.

As a summary measure, the average life satisfaction score is also calculated and presented in Table SW.2. The score is simply calculated by averaging the responses to the question on overall life satisfaction, ranging from very satisfied (1) to very unsatisfied (5) (see Questionnaires in Appendix H). Therefore, the lower the average score, the higher the life satisfaction levels. Average life satisfaction score for women age 15-24 years is 1.6. The table also shows that 92 percent of women age 15-24 years are very or somewhat happy.

Table SW.2: Overall life satisfaction and happiness (women)				
Percentage of women age 15-24 years who are very or somewhat satisfied with their life overall, the average overall life satisfaction score, and percentage of women age 15-24 years who are very or somewhat happy, Turkana County MICS, 2013/14				
	Percentage of women with overall life satisfaction ¹	Average life satisfaction score	Percentage of women who are very or somewhat happy ²	Number of women age 15-24 years
Total	89.7	1.6	91.5	461
Age				
15-19	90.8	1.5	90.6	252
20-24	88.4	1.6	92.6	209
Area				
Urban	91.3	1.5	92.0	345
Rural	85.1	1.7	89.9	116
Marital Status				
Ever married/in union	89.0	1.6	94.1	176
Never married/in union	90.2	1.6	89.9	285
Education				
None	87.4	1.7	93.8	152
Primary	88.9	1.6	89.4	196
Secondary+	94.4	1.4	92.1	112
Wealth index quintile				
Poorest	87.7	1.8	91.2	44
Second	79.7	1.8	91.7	45
Middle	87.2	1.6	90.9	100
Fourth	89.8	1.7	88.5	127
Richest	95.0	1.3	94.6	145
Ethnicity of household head				
Turkana	90.2	1.5	93.0	350
Other ethnic group	88.3	1.7	86.8	110
¹ MICS Indicator 11.1 - Life satisfaction				
² MICS indicator 11.2 - Happiness				

In addition to the series of questions on life satisfaction and happiness, respondents were also asked two simple questions on whether they think their life improved during the last one year, and whether they think their life will be better in one year's time. Such information may contribute to our understanding of desperation that may exist among young people, as well as hopelessness and hopes for the future. Specific combinations of the perceptions during the last one year and expectations for the next one year may be valuable information to understand the general sense of well-being among young people.

In Table SW.3, women's perceptions of a better life are shown. The proportion of women age 15-24 years who believe that their lives improved during the last one year and who expect that their lives would get better after one year, was 63 percent. Differences in the perception of a better life can be observed by age: 69 percent of young women age 15-24 years believe that their lives had improved during the last one year and expect that it would get better after one year, while the corresponding proportion for young women age 20-24 years is 56 percent; and by urban/rural areas, 68 percent and 48 percent, respectively.

Table SW.3: Perception of a better life (women)				
Percentage of women age 15-24 years who think that their lives improved during the last one year and those who expect that their lives will get better after one year, Turkana County MICS, 2013/14				
	Percentage of women who think that their life			Number of women age 15-24 years
	Improved during the last one year	Will get better after one year	Both ¹	
Total	65.3	86.8	62.9	461
Age				
15-19	71.4	90.0	69.0	252
20-24	57.9	83.0	55.6	209
Area				
Urban	69.8	89.7	67.8	345
Rural	51.8	78.0	48.2	116
Marital Status				
Ever married/in union	65.7	87.2	63.2	176
Never married/in union	65.1	86.6	62.7	285
Education				
None	56.6	81.7	53.9	152
Primary	70.0	90.3	69.2	196
Secondary+	69.0	87.6	64.2	112
Wealth index quintile				
Poorest	48.7	75.8	45.2	44
Second	51.2	77.4	48.7	45
Middle	59.3	85.0	57.4	100
Fourth	66.7	90.8	65.5	127
Richest	77.6	90.7	74.4	145
Ethnicity of household head				
Turkana	64.7	84.9	61.7	350
Other ethnic group	67.5	92.9	66.8	110

¹ MICS indicator 11.3 - Perception of a better life

14. Tobacco and Alcohol Use

Tobacco products are products made entirely or partly of leaf tobacco as raw material, which are intended to be smoked, sucked, chewed, or snuffed. All contain the highly addictive psychoactive ingredient, nicotine. Tobacco use is one of the main risk factors for a number of chronic diseases, including cancer, lung diseases, and cardiovascular diseases.¹²¹

The consumption of alcohol carries a risk of adverse health and social consequences related to its intoxicating, toxic and dependence-producing properties. In addition to the chronic diseases that may develop in those who drink large amounts of alcohol over a number of years, alcohol use is also associated with an increased risk of acute health conditions, such as injuries, including from traffic accidents.¹²² Alcohol use also causes harm far beyond the physical and psychological health of the drinker. It harms the well-being and health of people around the drinker. An intoxicated person can harm others or put them at risk of traffic accidents or violent behaviour, or negatively affect co-workers, relatives, friends or strangers. Thus, the impact of the harmful use of alcohol reaches deep into society.¹²³

Tobacco control campaigns were initiated in Kenya in 1992 as part of the World No Tobacco Day celebration. In 2001, the Ministry of Health (MOH) established the National Tobacco Free Initiative Committee (NTFIC) to coordinate tobacco control activities, and a tobacco control focal point was designated.¹²⁴ The Government of Kenya participated in formulation of the 2003 WHO Framework Convention on Tobacco Control (FCTC) which contains articles aimed at reducing the supply of and demand for tobacco; protection from exposure to smoke; and a provision that addresses liability¹²⁵. Kenya ratified the convention in 2004. Tobacco Control Act [*Chapter 245A*] was enacted in 2007 to control the production, manufacture, sale, labelling, advertising, promotion and sponsorship of tobacco products, and the National Tobacco Control Action Plan was launched in 2010. Liquor control in the country is through the Liquor Licensing Act [*Chapter 121*].

The MICS collected information on ever and current use of tobacco and alcohol and intensity of use among women age 15-49 years in Turkana County. This section presents the main results.

14.1 Tobacco Use

Table TA.1 presents the current and ever use of tobacco products by women age 15-49 years. In Turkana County MICS, ever use of any tobacco product among women is 22 percent, while 21 percent smoke cigarettes, or used smoked or smokeless tobacco products on one or more days during the last one month prior to the survey. Smoking was more prevalent in rural than urban areas, and among women with no education. The results further indicate that majority of the smokers are from the poorest and the second poorest wealth quintiles.

¹²¹WHO. <http://www.who.int/topics/tobacco/en/>

¹²²WHO. http://www.who.int/topics/alcohol_drinking/en/

¹²³ WHO. <http://www.who.int/mediacentre/factsheets/fs349/en/>

¹²⁴ WHO. 2012. Joint national capacity assessment on the implementation of effective tobacco control policies in Kenya.

¹²⁵ WHO. 2005. Framework Convention on Tobacco Control


Table TA.1: Current and ever use of tobacco (women)

Percentage of women age 15-49 years by pattern of use of tobacco, Turkana County MICS, 2013/14

	Never smoked cigarettes or used other tobacco products	Ever users				Users of tobacco products at any time during the last one month				Number of women age 15-49 years
		Only cigarettes	Cigarettes and other tobacco products	Only other tobacco products	Any tobacco product	Only cigarettes	Cigarettes and other tobacco products	Only other tobacco products	Any tobacco product ¹	
Total	77.5	0.4	0.2	21.7	22.3	0.1	0.0	20.6	20.7	1104
Age										
15-19	93.0	0.0	0.0	6.7	6.7	0.0	0.0	5.7	5.7	252
20-24	83.6	1.6	0.5	13.6	15.7	0.0	0.0	13.6	13.6	209
25-29	74.2	0.5	0.0	25.2	25.6	0.5	0.0	24.0	24.4	210
30-34	69.2	0.0	0.0	30.8	30.8	0.0	0.0	28.5	28.5	144
35-39	74.1	0.0	0.5	25.4	25.9	0.0	0.0	24.9	24.9	134
40-44	56.7	0.0	0.6	42.8	43.3	0.0	0.0	41.6	41.6	82
45-49	62.1	0.0	0.0	37.9	37.9	0.0	0.0	35.5	35.5	75
Area										
Urban	88.3	0.6	0.2	10.8	11.7	0.1	0.0	9.5	9.6	683
Rural	59.9	0.0	0.1	39.4	39.5	0.0	0.0	38.7	38.7	421
Education										
None	64.1	0.0	0.2	35.3	35.5	0.0	0.0	34.1	34.1	622
Primary	92.8	0.9	0.3	5.9	7.2	0.3	0.0	5.0	5.4	296
Secondary+	98.0	0.9	0.0	1.2	2.0	0.0	0.0	0.3	0.3	186
Under-5s in the same household										
At least one	77.9	0.5	0.2	21.3	21.9	0.0	0.0	20.4	20.4	757
None	76.6	0.3	0.3	22.5	23.0	0.3	0.0	21.1	21.4	347
Wealth index quintile										
Poorest	47.4	0.0	0.3	51.9	52.1	0.0	0.0	50.7	50.7	169
Second	57.2	0.0	0.4	41.4	41.8	0.0	0.0	40.8	40.8	191
Middle	81.8	0.8	0.0	17.4	18.2	0.0	0.0	17.0	17.0	226
Fourth	89.2	1.1	0.4	9.4	10.8	0.4	0.0	7.9	8.2	249
Richest	96.2	0.0	0.0	3.8	3.8	0.0	0.0	2.3	2.3	270
Ethnicity of household head										
Turkana	72.4	0.2	0.2	26.9	27.4	0.0	0.0	25.6	25.6	883
Other ethnic group	98.1	1.2	0.0	0.7	1.9	0.5	0.0	0.7	1.2	218

¹ MICS indicator 12.1 - Tobacco use

Figure TA.1: Ever and current smokers, Turkana County MICS, 2013/14



None among the 1,104 women age 15-49 years sampled in the survey stated having smoked a whole cigarette before age 15, which indicates that the actual percentage in the population must be extremely low. Further, only one woman age 15-49 years in our sample stated to be a current cigarette smoker. For this reason, corresponding statistics are not presented in this report.

14.2 Alcohol Use

Table TA.2 shows the use of alcohol among women. About 10 percent of women age 15-49 years, had at least one drink of alcohol on one or more days during the last one month preceding the survey while four percent have had at least one alcoholic drink before the age of 15 years. The proportion who had an alcoholic drink in the last month preceding the survey ranged between three percent and 18 percent by age. Women age 15-49 years in urban areas are twice (5 percent) as likely to have had at least one alcoholic drink before age 16 than their rural counterparts (2 percent). The results further indicate that drinking of at least one alcoholic drink before age 15 increases with increase in wealth quintiles. About two percent of the women in the poorest wealth quintile have had at least one alcoholic drink before age 15 compared to five percent of those from the richest wealth quintile.

Table TA.2: Use of alcohol (women)				
Percentage of women age 15-49 years who have never had an alcoholic drink, percentage who first had an alcoholic drink before age 15, and percentage of women who have had at least one alcoholic drink at any time during the last one month, Turkana County MICS, 2013/14				
	Percentage of women who:			Number of women age 15-49 years
	Never had an alcoholic drink	Had at least one alcoholic drink before age 15 ¹	Had at least one alcoholic drink at any time during the last one month ²	
Total	83.0	4.2	9.8	1.104
Age				
15-19	85.6	8.5	3.2	252
20-24	82.4	2.6	8.5	209
25-29	86.4	3.1	8.7	210
30-34	85.1	1.8	12.5	144
35-39	78.8	4.1	15.6	134
40-44	76.2	5.4	14.1	82
45-49	76.4	0.8	17.7	75
Area				
Urban	81.9	5.3	9.3	683
Rural	84.7	2.4	10.5	421
Education				
None	83.0	3.3	12.5	622
Primary	82.7	7.2	5.5	296
Secondary+	83.3	2.4	7.5	186
Wealth index quintile				
Poorest	86.9	2.4	7.7	169
Second	81.9	2.6	12.8	191
Middle	79.0	4.7	15.4	226
Fourth	87.4	4.9	4.1	249
Richest	80.4	5.5	9.4	270
Ethnicity of household head				
Turkana	80.8	4.9	11.3	883
Other ethnic group	91.9	1.5	3.6	218
¹ MICS indicator 12.4 - Use of alcohol before age 15				
² MICS indicator 12.3 - Use of alcohol				

Appendix A. Documents Reviewed

African Development Bank Group, 2014. Kenya Country Strategy Paper 2024-2018.

Bajracharya, A ND Amin, S. 2010. *Poverty, marriage timing, and transitions to adulthood in Nepal: A longitudinal analysis using the Nepal living standards survey*. Poverty, Gender, and Youth Working Paper No. 19. Population Council.

Belsky, J et al. 2006. *Socioeconomic Risk, Parenting During the Preschool Years and Child Health Age 6 Years*. European Journal of Public Health 17(5): 511–2.

Bhuta, Z. et al. 2013. *Evidence-based interventions for improvement of maternal and child nutrition: what can be done and at what cost? The Lancet June 6, 2013*.

Cairncross, S and Cliff, JL. 1987. *Water use and Health in Mueda, Mozambique*. Transactions of the Royal Society of Tropical Medicine and Hygiene 81: 51-4.

Cairncross, S and Valdmanis, V. 2006. *Water supply, sanitation and hygiene promotion* Chapter 41 in *Disease Control Priorities in Developing Countries*. 2nd Edition, Edt. Jameson et al. The World Bank.

Cairncross, S et al. 2010. *Water, sanitation and hygiene for the prevention of diarrhoea*. International Journal of Epidemiology 39: i193-i205

Campbell, H. et al. 2013. *Measuring Coverage in MNCH: Challenges in Monitoring the Proportion of Young Children with Pneumonia Who Receive Antibiotic Treatment*. PLoS Med 10(5): e1001421.
doi:10.1371/journal.pmed.1001421

Clark, S et al. 2006. *Protecting young women from HIV/AIDS: the case against child and adolescent marriage*. International Family Planning Perspectives 32(2): 79-88.

Constitution of Kenya of 2010 [Promulgated on 25th August 2010]

D'Acremont, V et al. 2010. *Reduction in the proportion of fevers associated with Plasmodium falciparum parasitaemia in Africa: a systematic review*. Malaria Journal 9(240).

DfID. 2013. *Water, Sanitation and Hygiene: Evidence Paper*.

<http://r4d.dfid.gov.uk/pdf/outputs/sanitation/WASH-evidence-paper-april2013.pdf>

Division of Malaria Control [Ministry of Public Health and Sanitation], Kenya National Bureau of Statistics, and ICF Macro. 2011. 2010 Kenya Malaria Indicator Survey. Nairobi, Kenya: DOMC, KNBS and ICF Macro.

Erickson, MF and Egeland, B. 1987. *A Developmental View of the Psychological Consequences of Maltreatment*. School Psychology Review 16: 156-68.

Framework of Actions for the follow - up to the Programme of Action of the International Conference on Population and Development Beyond 2014

Gaisie, S. Fertility Trend in Ghana. African Population Studies Vol. 20 N°2/Etude de la population africaine vol. 20 n° 2.

Godha, D et al. 2011. *The influence of child marriage on fertility, fertility-control, and maternal health care utilization*. MEASURE/Evaluation PRH Project Working paper 11-124.

Government of Kenya 2014. Kenya AIDS Response Progress Report 2014 – Progress Towards Zero.

Government of Kenya, 2011. National Food and Nutrition Security Policy.

Government of Kenya. 2010. The Constitution of Kenya 2010

Government of Kenya. National School Health Policy. Ministry of Public Health and Sanitation and Ministry of Education.

Nairobi: Republic of Kenya; 2009.

Government of Kenya. Children Act [141] 2007, 2010.

Government of Kenya. Employment Act [Chapter 226] 2007, 2012

Government of Kenya. 2006. Early Childhood Development Service Standard Guidelines for Kenya 2006.

Government of Kenya. 2006. National Early Childhood Development Policy Framework 2006.

Grantham-McGregor, S et al. 2007. *Developmental Potential in the First 5 Years for Children in Developing Countries*. The Lancet 369: 60–70

Grossman, DC. 2000. The History of Injury Control and the Epidemiology of Child and Adolescent Injuries. The Future of Children, 10(1): 23-52.

HMN, UNICEF, and WHO. 2008. *Countdown to 2015: Tracking Progress in Maternal, Newborn & Child Survival, The 2008 Report*.

Kenta Health Sector Strategic and Investment Plan (KHSSPI) July 2013-June 2017.

Kenya Demographic and Health Survey 2014

Kenya Human Rights Commission. 2010. *Towards Equality and Anti-Discrimination: An Overview of International and Domestic Law an Anti-discrimination in Kenya*.

Kenya National Bureau of Statistics, 2013. *Statistical Abstract 2013*.

Kenya National Bureau of Statistics, 2014. *Kenya Facts and Figures 2014*.

Kenya National Bureau of Statistics. 2015. *Kenya Demographic and Health Survey, 2014*.

Kenya National Commission for Human Rights. 2012. *Realising Sexual and Reproductive Health Rights in Kenya: A myth or reality? A Report of the Public Inquiry into Violations of Sexual and Reproductive Health Rights in Kenya April 2012*.

Lawn, JE et al. 2005. *4 million neonatal deaths: When? Where? Why?* *Lancet* 2005; 365:891–900.

Liquor Licensing Act [*Chapter 121*]

Ministry of Education, Science and Technology. 2015. *Education for All 2015 National Review*.

Ministry of Education Science and Technology, 2014. *Basic Education Statistical Booklet*.

Ministry of Education, Science and Technology, 2005. *Kenya Education Sector Support Programme 2005-2010*.

Ministry of Health, 2014. *Kenya HIV Estimates*.

Ministry of Health, 2013. *Mother and Child Health Booklet*. Republic of Kenya.

Ministry of Health, 2013. *Transforming Health: Accelerating Attainment of Universal Health Coverage*.

Ministry of Health. 2006. *Division of Vaccines and Immunization (DVI) Multi Year Plan 2006-2010*.

Ministry of Health. 2012. *Guidelines for the Prevention of Mother-To-Child Transmission (PMTCT) of HIV/AIDS in Kenya*.

- Ministry of Information and Communications. 2006. National Information and Communications Technology (ICT) Policy.
- Ministry of Public Health and Sanitation Division of Malaria Control. National Malaria Communication Strategy 2010 – 2013
- Ministry of Public Health and Sanitation Division of Malaria Control. 2010. National Malaria Policy April 2010.
- Ministry of Public Health and Sanitation Division of Malaria Control. 2009. National Malaria Strategy 2009-2017 – Towards a Malaria-free Kenya
- Ministry of Public Health and Sanitation, and Ministry of Medical Services. 2010. National Guidelines for the Diagnosis, Treatment and Prevention of Malaria in Kenya.
- Ministry of Public Health and Sanitation. 2010. National Tobacco Control Programme.
- Ministry of Public Health and Sanitation. 2012. National Nutrition Action Plan 2012-2017
- Ministry of Water and Irrigation. 2012. The National Water Policy 2012
- OECD. 2013. *OECD Guidelines on Measuring Subjective Well Being*. OECD. <http://dx.doi.org/10.1787/9789264191655-en>
- PAHO. 2003. *Guiding principles for complementary feeding of the breastfed child*.
- President's Malaria Initiative - Kenya Malaria Operational Plan FY 2014
- Raj, A et al. 2009. *Prevalence of child marriage and its effect on fertility and fertility-control outcomes of young women in India: a cross-sectional, observational study*. The Lancet 373(9678): 1883–9.
- Ram, P et al. editors. 2008. *Use of a novel method to detect reactivity to structured observation for measurement of handwashing behavior*. American Society of Tropical Medicine and Hygiene.
- Save the Children with support from Oxfam, 2013. A Cost of the Diet analysis in Turkana district of Kenya. Kerio Agropastoral livelihood zone, Turkana. Funded by ECHO.
- Say, L et al. 2014. *Global causes of maternal death: a WHO systematic analysis*. The Lancet Global Health 2(6): e323-33. DOI: 10.1016/S2214-109X(14)70227-X
- Schneider, MW et al. 2005. *Do Allegations of Emotional Maltreatment Predict Developmental Outcomes Beyond that of Other Forms of Maltreatment?* Child Abuse & Neglect 29(5): 513–32.

Shonkoff, J and Phillips, D (eds). 2000. *From neurons to neighborhoods: the science of early childhood development*. Committee on Integrating the Science of Early Childhood Development, National Research Council, 2000.

Shulman, CE and Dorman, EK. 2003. *Importance and prevention of malaria in pregnancy*. *Trans R Soc Trop Med Hyg* 97(1): 30–55.

Straus, MA and Paschall MJ. 2009. *Corporal Punishment by Mothers and Development of Children's Cognitive Ability: A longitudinal study of two nationally representative age cohorts*. *Journal of Aggression, Maltreatment & Trauma* 18(5): 459-83.

The Kenyan Section of the International Commission of Jurists. 2004. *International Human Rights Standards: Reporting Obligations – The Convention of the Rights of the Child*.

The Partnership for Maternal, Newborn and Child Health, 2012. *Maternal and Child Health: Kenya*

The United Nations General Assembly. 1989. *The Convention on the Rights of the Child*.

Tobacco Control Act [*Chapter 245A*] 2012.

UNESCO, 2015. *EFA Monitoring Report 2015 -Education for All 2000-2015: Achievements and Challenges*

UN Interagency Group for Child Mortality Estimation. 2013. *Levels and Trends in Child Mortality: Report 2013*

UNICEF Fact sheet http://www.unicef.org/media/media_81674.html

UNICEF. 2014. *The State of the World's Children 2015*. UNICEF.

United Nations Children's Fund (UNICEF), September 2014. *Committing to Child Survival: A Promise Renewed - Progress Report 2014*.

UNICEF. 2013. *Every Child's Birth Right: Inequities and trends in birth registration*. UNICEF.

UNICEF. 2012. *How Sensitive Are Estimates of Child Labour to Definitions?* MICS Methodological Paper No. 1. UNICEF.

UNICEF. 2002. *A World Fit For Children* adopted by the UN General Assembly at the 27th Special Session, 10 May 2002: 2.

United Nations General Assembly Special Session on HIV/AIDS 2001. Summary of the Declaration of Commitment on HIV/AIDS 25-27 June 2001, New York

United Nations. 2002. A World Fit for Children. Resolution adopted by the United Nations General Assembly 10 May 2002.

WHO, UNICEF, UNFPA, The World Bank. 2012. *Trends in Maternal Mortality: 1990-2010*. World Health Organization.

WHO. 2014. Implementing the Commission on Information and Accountability Recommendations 2014: Progress Report Accountability for Women's and Children's Health.

WHO. 2005. Framework Convention on Tobacco Control

WHO. 2005. *Guiding principles for feeding non-breastfed children 6-24 months of age*.

WHO. 2003. *Global Strategy for Infant and Young Child Feeding*.

WHO. 2003. *Implementing the Global Strategy for Infant and Young Child Feeding*. Meeting Report Geneva, 3-5 February, 2003.

WHO. 2012. Joint national capacity assessment on the implementation of effective tobacco control policies in Kenya.

WHO. 2010. PMTCT Strategic Vision 2010-15: Preventing mother-to-child transmission of HIV to reach the UNGASS and Millennium Development Goals - Moving towards the elimination of paediatric HIV

WHO. 2010. WHO Guidelines on Prevention of Mother-to-Child Transmission (PMTCT) programmes

WHO. 2008. *Indicators for assessing infant and young child feeding practices. Part 1: Definitions*.

WHO. 2002. [Clinical Guidelines for Diagnosis and Treatment of Common Conditions in Kenya](#) (WHO; 2002; 344 pages)

WHO. <http://www.who.int/mediacentre/factsheets/fs349/en/>

WHO. http://www.who.int/topics/alcohol_drinking/en/

WHO. <http://www.who.int/topics/tobacco/en/>

WHO/UNICEF JMP. 2008. *MDG assessment report*.

http://www.wssinfo.org/fileadmin/user_upload/resources/1251794333-JMP_08_en.pdf

WHO/UNICEF. 2012. *Progress on Drinking water and Sanitation: 2012 update*.

Wolf, J et al. 2014. *Systematic review: Assessing the impact of drinking water and sanitation on diarrhoeal disease in low- and middle-income settings: systematic review and meta-regression*.

Tropical Medicine and International Health 2014.

Appendix B. Education ISCED Tables

Table ED.4: Primary school attendance and out of school children (ISCED)

Percentage of children of primary school age attending primary or secondary school (adjusted net attendance ratio), percentage attending preschool, and percentage out of school, Turkana County MICS, 2013/14

	Male					Female					Total				
	Percentage of children:					Percentage of children:					Percentage of children:				
	Net attendance ratio (adjusted)	Not attending school or preschool	Attending preschool	Out of school ^a	Number of children	Net attendance ratio (adjusted)	Not attending school or preschool	Attending preschool	Out of school ^a	Number of children	Net attendance ratio (adjusted) ¹	Not attending school or preschool	Attending preschool	Out of school ^a	Number of children
Total	63.3	25.9	9.1	35.0	613	61.3	27.3	9.6	36.9	563	62.4	26.6	9.3	35.9	1,176
Area															
Urban	78.4	8.0	11.8	19.9	319	77.9	6.6	13.9	20.4	280	78.2	7.3	12.8	20.1	599
Rural	47.0	45.3	6.0	51.3	294	44.9	47.8	5.4	53.2	283	46.0	46.5	5.7	52.2	577
Age at beginning of school year															
6	46.5	26.5	24.9	51.4	112	38.7	28.2	29.6	57.8	90	43.1	27.3	27.0	54.3	202
7	46.7	32.4	15.1	47.4	115	60.6	22.3	16.0	38.4	117	53.7	27.3	15.6	42.8	232
8	65.5	28.0	5.0	33.0	104	62.3	29.8	4.0	33.9	76	64.1	28.8	4.6	33.4	180
9	70.6	26.1	3.4	29.4	110	69.4	29.8	0.8	30.6	109	70.0	27.9	2.1	30.0	219
10	78.4	21.0	0.6	21.6	83	71.3	26.5	2.2	28.7	85	74.8	23.8	1.4	25.2	169
11	80.2	18.7	1.1	19.8	89	65.3	28.4	3.1	31.5	84	73.0	23.4	2.1	25.5	173
Mother's education															
None	56.1	33.0	9.5	42.5	456	54.1	35.9	9.1	44.9	419	55.2	34.4	9.3	43.7	875
Primary	84.2	5.5	7.1	12.6	98	85.5	1.1	12.2	13.3	86	84.8	3.5	9.5	12.9	184
Secondary+	86.2	4.8	8.9	13.8	58	81.9	3.7	12.1	15.9	46	84.3	4.3	10.4	14.7	104
Cannot be determined ^b		-	-	-	0	(*)	(*)	(*)	(*)	9	(*)	(*)	(*)	(*)	9
Wealth index quintile															
Poorest	31.5	58.3	10.1	68.5	127	27.4	63.5	8.6	72.1	131	29.4	60.9	9.4	70.3	258
Second	46.1	41.8	9.3	51.0	139	47.2	42.3	7.7	50.1	116	46.6	42.0	8.6	50.6	255
Middle	74.6	12.7	8.8	21.5	127	70.2	14.8	10.5	25.2	111	72.6	13.7	9.6	23.3	238

Fourth	84.1	7.5	8.4	15.9	128	85.4	2.9	11.7	14.6	109	84.7	5.4	9.9	15.3	237
Richest	88.6	1.1	8.5	9.6	93	87.1	1.8	10.0	11.8	96	87.9	1.5	9.2	10.7	189
Ethnicity of household head															
Turkana	60.6	28.3	9.5	37.8	523	57.2	31.2	9.5	40.7	480	59.0	29.7	9.5	39.2	1,004
Other ethnic group	79.3	11.4	6.6	18.0	88	85.4	4.1	10.5	14.6	82	82.3	7.9	8.5	16.3	170

¹MICS indicator 7.4; MDG indicator 2.1 - Primary school net attendance ratio (adjusted)

^aThe percentage of children of primary school age out of school are those not attending school and those attending preschool

^bChildren age 15 or higher at the time of the interview whose mothers were not living in the household

(*) Figures that are based on fewer than 25 unweighted cases

Table ED.5: Secondary school attendance and out of school children (ISCED)

Percentage of children of secondary school age attending secondary school or higher (adjusted net attendance ratio), percentage attending primary school, and percentage out of school, Turkana County MICS, 2013/14

	Male				Female				Total			
	Percentage of children:				Percentage of children:				Percentage of children:			
	Net attendance ratio (adjusted)	Attending primary school	Out of school ^a	Number of children	Net attendance ratio (adjusted)	Attending primary school	Out of school ^a	Number of children	Net attendance ratio (adjusted) ¹	Attending primary school	Out of school ^a	Number of children
Total	36.9	36.6	24.6	554	36.8	32.0	29.7	522	36.8	34.4	27.1	1,076
Area												
Urban	46.5	43.4	8.3	335	49.1	31.9	16.9	343	47.8	37.6	12.7	678
Rural	22.2	26.2	49.7	219	13.2	32.0	54.1	179	18.2	28.8	51.7	398
Age at beginning of school year												
12	10.9	63.0	23.1	115	11.3	61.1	26.6	116	11.1	62.1	24.8	230
13	19.7	54.8	23.7	92	29.7	42.8	27.5	108	25.1	48.3	25.8	200
14	28.3	47.4	20.6	99	38.3	28.4	31.7	74	32.6	39.3	25.3	174
15	48.9	19.0	32.0	77	41.4	29.0	29.7	75	45.2	23.9	30.9	153
16	64.4	14.5	21.2	81	64.8	3.5	31.8	77	64.6	9.1	26.3	157
17	62.1	7.3	28.8	90	52.1	6.4	33.6	73	57.6	6.9	30.9	162

Mother's education												
None	20.3	45.8	32.0	274	15.9	44.8	39.1	227	18.3	45.3	35.2	501
Primary	(39.8)	(49.3)	(10.9)	56	37.7	47.3	14.2	73	38.6	48.2	12.8	129
Secondary	(*)	(*)	(*)	30	(81.0)	(19.0)	(0.0)	49	65.7	34.3	0.0	79
Cannot be determined ^b	59.3	16.6	22.2	192	51.3	12.4	32.3	173	55.5	14.6	27.0	366
Wealth index quintile												
Poorest	14.9	14.9	68.8	97	4.1	14.9	81.1	72	10.3	14.9	74.0	169
Second	25.5	38.3	32.8	98	12.2	30.4	56.5	67	20.1	35.1	42.4	165
Middle	31.7	48.1	17.5	118	25.5	45.8	25.8	110	28.7	47.0	21.5	228
Fourth	44.7	45.0	10.3	116	39.8	44.2	13.9	129	42.1	44.6	12.2	245
Richest	60.5	33.5	4.1	125	70.3	19.8	8.7	145	65.8	26.1	6.5	270
Ethnicity of household head												
Turkana	36.7	32.4	29.0	436	36.6	27.9	33.8	416	36.6	30.2	31.4	852
Other ethnic group	36.6	52.9	8.8	113	37.8	47.7	13.4	105	37.1	50.4	11.0	219
¹ MICS indicator 7.5 - Secondary school net attendance ratio (adjusted)												
^a The percentage of children of secondary school age out of school are those who are not attending primary, secondary, or higher education												
^b Children age 15 or higher at the time of the interview whose mothers were not living in the household												
() Figures that are based on 25-49 unweighted cases												
(*) Figures that are based on fewer than 25 unweighted cases												

Table ED.7: Primary school completion and transition to secondary school (ISCED)

Primary school completion rates and transition and effective transition rates to secondary school, Turkana County MICS, 2013/14						
	Primary school completion rate ¹	Number of children of primary school completion age	Transition rate to secondary school ²	Number of children who were in the last grade of primary school the previous year	Effective transition rate to secondary school	Number of children who were in the last grade of primary school the previous year and are not repeating that grade in the current school year
Total	84.2	173	96.4	190	98.6	186
Sex						
Male	95.1	89	94.7	99	98.9	95
Female	72.7	84	98.3	91	98.3	91
Area						
Urban	93.2	102	98.0	149	99.3	147
Rural	71.3	71	(90.7)	41	(95.9)	39
¹ MICS indicator 7.7 - Primary completion rate						
² MICS indicator 7.8 - Transition rate to secondary school						
() Figures that are based on 25-49 unweighted cases						

Table ED.8: Education gender parity (ISCED)

Ratio of adjusted net attendance ratios of girls to boys, in primary and secondary school, Turkana County MICS, 2013/14						
	Primary school			Secondary school		
	Primary school adjusted net attendance ratio (NAR), girls	Primary school adjusted net attendance ratio (NAR), boys	Gender parity index (GPI) for primary school adjusted NAR ¹	Secondary school adjusted net attendance ratio (NAR), girls	Secondary school adjusted net attendance ratio (NAR), boys	Gender parity index (GPI) for secondary school adjusted NAR ²
Total	61.3	63.3	0.97	36.8	36.9	1.00
Area						
Urban	77.9	78.4	0.99	49.1	46.5	1.06
Rural	44.9	47.0	0.96	13.2	22.2	0.60
Mother's education						
None	54.1	56.1	0.96	15.9	20.3	0.78
Primary	85.5	84.2	1.01	(37.7)	(*)	0.94
Secondary+	81.9	86.2	0.95	(*)	(*)	1.98
Cannot be determined ^a	(*)	-	-	51.3	59.3	0.87
Wealth index quintile						
Poorest	27.4	31.5	0.87	(4.1)	14.9	0.27
Second	47.2	46.1	1.02	(12.2)	25.5	0.48
Middle	70.2	74.6	0.94	25.5	31.7	0.81
Fourth	85.4	84.1	1.02	39.8	44.7	0.89
Richest	87.1	88.6	0.98	70.3	60.5	1.16
Ethnicity of household head						
Turkana	57.2	60.6	0.94	36.6	36.7	1.00
Other ethnic group	85.4	79.3	1.08	37.8	36.6	1.03
¹ MICS indicator 7.9; MDG indicator 3.1 - Gender parity index (primary school)						

² MICS indicator 7.10; MDG indicator 3.1 - Gender parity index (secondary school)

^a Children age 15 or higher at the time of the interview whose mothers were not living in the household

na: not applicable

() Figures that are based on 25-49 unweighted cases

(*) Figures that are based on fewer than 25 unweighted cases

Table ED.9: Out of school gender parity (ISCED)

Percentage of girls in the total out of school population, in primary and secondary school, Turkana County MICS, 2013/14

	Primary school				Secondary school			
	Percentage of out of school children	Number of children of primary school age	Percentage of girls in the total out of school population of primary school age	Number of children of primary school age out of school	Percentage of out of school children	Number of children of secondary school age	Percentage of girls in the total out of school population of secondary school age	Number of children of secondary school age out of school
Total	35.9	1,176	49.2	422	27.1	1,076	53.2	291
Area								
Urban	20.1	599	47.5	121	12.7	678	67.7	86
Rural	52.2	577	49.9	301	51.7	398	47.2	206
Mother's education								
None	43.7	875	49.2	382	35.2	501	50.3	176
Primary	12.9	184	(*)	24	12.8	129	(*)	16
Secondary+	14.7	104	(*)	15	0.0	79	-	0
Cannot be determined ^a	na	na	na	na	27.0	366	56.7	99
Wealth index quintile								
Poorest	70.3	258	52.2	181	74.0	169	46.6	125
Second	50.6	255	44.9	129	42.4	165	54.2	70
Middle	23.3	238	50.6	55	21.5	228	58.0	49
Fourth	15.3	237	(43.9)	36	12.2	245	(60.0)	30
Richest	10.7	189	(*)	20	6.5	270	(*)	18
Ethnicity of household head								
Turkana	39.2	1,004	49.7	393	31.4	852	52.7	267
Other ethnic group	16.3	170	(42.9)	28	11.0	219	(*)	24
^a Children age 15 or higher at the time of the interview whose mothers were not living in the household								
na: not applicable								
() Figures that are based on 25-49 unweighted cases								
(*) Figures that are based on fewer than 25 unweighted cases								

Appendix C. Sample Design

The major features of the sample design are described in this appendix. Sample design features include sample size, sample allocation, sampling frame and listing, choice of domains, sampling stages, stratification, and the calculation of sample weights.

The primary objective of the sample design for the Turkana County MICS was to produce statistically reliable estimates of indicators, at county level. The urban and rural areas in Turkana County were the sampling strata. A multi-stage, stratified cluster sampling approach was used for the selection of the survey sample.

Sample Size and Sample Allocation

The sample size for the Turkana County MICS was calculated as 1,740 households. For the calculation of the sample size, the key indicator used was the basic immunization for children aged 12-23 months. The following formula was used to estimate the required sample size for this indicator:

$$n = \frac{[4(r)(1-r)(deff)]}{[(0.12r)^2 (pb)(AveSize)(RR)]}$$

where

n is the required sample size, expressed as number of households

4 is a factor to achieve the 95 percent level of confidence

r is the predicted or anticipated value of the indicator, expressed in the form of a proportion

$deff$ is the design effect for the indicator, estimated from a previous survey or using a default value of 1.5

$0.12r$ is the margin of error to be tolerated at the 95 percent level of confidence, defined as 12 per cent of r (relative margin of error of r)

pb is the proportion of the total population upon which the indicator, r , is based

$AveSize$ is the average household size (number of persons per household)

RR is the predicted response rate

For the calculation, r (basic immunization for children aged 12-23 months) was assumed to be 57.3 percent based on the results for MICS3 in the neighbouring Marsabit district. The value of $deff$ (design effect) was taken as 1.5 based on estimates from previous surveys, pb (percentage of children aged 12-23 months in Turkana County) was taken as 1.7 percent, $AveSize$ (average household size in Turkana County) was taken as 6.9. Both pb and $AveSize$ were based on the results from the 2009 Kenya Population and Housing Census. The margin of error to be tolerated at the 95 per cent level of confidence was fixed at $0.15r$ and the response rate was assumed to be 90 per cent based on experience from previous surveys.

The resulting number of households from this exercise was 1,740 households which is the sample size for Turkana County. The number of households selected per cluster was 30 households, and was based on a number of considerations, including design effect, the budget available, and the time that would be needed per team to complete one cluster. By dividing the total number of households by the number of sample households per cluster, it was determined that 58 clusters be sampled in the county.

Power allocation method was used to allocate the sample to the urban and rural strata of Turkana County. The table below shows the distribution of sampled households and clusters in the sampling strata.

	Number of households			Number of Clusters		
	Total	Urban	Rural	Total	Urban	Rural
Total	1,740	660	1,080	58	22	36

Sampling Frame and Selection of Clusters

MICS5 utilized the recently created fifth National Sample Survey and Evaluation Programme (NASSEP V) frame which is a household based master sampling frame developed and maintained by KNBS. The frame was implemented using a multi-tiered structure, in which a set of 4 sub-samples (C1, C2, C3, C4) were developed. It is based on the list of enumeration areas (EAs) from the 2009 Kenya Population and Housing Census. The frame is stratified according to County and further into rural and urban. Each of the sub-samples is representative at county level and at national (i.e. Urban/rural) level and contains 1,340 clusters.

The Primary Sampling Units (PSUs) for the survey were clusters drawn from the NASSEP V sampling frame, so the first component of the probabilities and weights are based on that master sample. Within each stratum the PSUs for the MICS were selected independently from one of the subsamples of the master sample using Equal Probability Selection Method (EPSEM). A total of 58 clusters were selected from the master sample in this way.

Cluster Updating Activities

Out of the 58 clusters selected for Turkana County, it was established that 30 had been listed more than six months prior to the start of the survey. These listing for these clusters was updated prior to selection of households. For this purpose, listing teams visited each cluster, and listed all occupied households. For the remaining 28 sample clusters a more recent listing was available, so it was used for selecting the sample households.

Selection of Households

A uniform sample of 30 households per cluster was selected using equal probability systematic sampling method. Non responding households were not replaced. Systematic sampling is a probability sample selection method in which the sample is obtained by selecting every k th element of the population where k is an integer greater than 1. The first number of the sample is selected randomly from within the first k elements.

Calculation of Sample Weights

The MICS5 sample was not self-weighting and thus a weighting process was required to provide estimates representative of the target population. Two main sampling weights were calculated: household weights and individual (women and children) weights. The base weights incorporated the probabilities of selection of the clusters from the census EAs database into the NASSEP V sample frame, the probabilities of selection of the MICS clusters from NASSEP V frame and the probabilities of selection of the households from each of the NASSEP V frame clusters. Base weights were then adjusted for cluster and household non-response by multiplying them by the inverse of the clusters and household response rates. The individual weight of a woman or child case was calculated as the household weight multiplied by the inverse of the individual response rate. Given that the MICS5 sample was a two-stage stratified cluster sample, sampling probabilities were calculated separately for each sampling stage. We will use the following notations:

P_{0hi} : sampling probability of the i^{th} EA in stratum h in the selection of the master sample from the 2009 census frame

P_{1hi} : first stage sampling probability of the i^{th} cluster in stratum h

P_{2hi} : second-stage sampling probability within the i^{th} cluster (households)

P_{hi} : overall sampling probability of any households of the i^{th} cluster in stratum h

For the NASSEP V master sample, EAs in each stratum were selected using a systematic probability proportional to size (PPS) sampling procedure. Let a_h be the number of EAs selected in stratum h , M_{hi} the measure of size (number of households) according to the 2009 census frame in the i^{th} EA, and $\sum M_{hi}$ the total measure of size (total number of households) in the stratum h . The probability of selecting the i^{th} EA in the NASSEP V master sample is calculated as follows:

$$P_{0hi} = \frac{a_h M_{hi}}{\sum M_{hi}}$$

Let b_h be the total number of clusters in stratum h of the NASSEP V master sample for the MICS5 and s_i the total number of segments created during listing of the i^{th} cluster. The probability of selecting

the i^{th} cluster in stratum h from the NASSEP V frame is calculated as follows: $P_{1hi} = \frac{a_h}{b_h} \times \frac{1}{s_h}$

Let L_{hi} be the number of households listed in the household listing operation in cluster i in stratum h ,

let g_{hi} be the number of households selected in the cluster. The second stage selection probability for each household in the cluster is calculated as follows:

$$P_{2hi} = \frac{g_{hi}}{L_{hi}}$$

The overall selection probability of each household in cluster i of stratum h is the product of the selection probabilities:

$$P_{hi} = D_{hi} \times P_{1hi} \times P_{2hi}$$

The sampling weight for each household in cluster i of stratum h is the inverse of its selection probability:

$$W_{hi} = \frac{1}{P_{hi}}$$

The individual weight of children or Women (W_{hi}) in cluster i is the household weight multiplied by the inverse of the individual response rate;

$$W_{hi} = W_{hi} \times \frac{E_{hi}}{I_{hi}},$$

Where, E_{hi} is the total eligible individuals (women or children) found in the i^{th} cluster of stratum h and I_{hi} is the total number of Individuals (women or children) with a successful interview.

After the completion of fieldwork, response rates were calculated for each cluster. These were used to adjust the sample weights calculated for each cluster. Response rates in the Turkana County MICS are shown in Table HH.1 in this report.

The non-response adjustment factors for the individual women and under-5 questionnaires were applied to the adjusted household weights. Numbers of eligible women and under-5 children were obtained from the roster of household members in the Household Questionnaire for households where interviews were completed.

The design weights for the households were calculated by multiplying the inverse of the probabilities of selection by the non-response adjustment factor for each cluster. These weights were then standardized (or normalized), one purpose of which is to make the weighted sum of the interviewed sample units equal to the unweighted total number of observations at the national level. Normalization is achieved by dividing the full sample weights (adjusted for nonresponse) by the average of these weights across all households at the national level. This is performed by multiplying the sample weights by a constant factor equal to the unweighted number of households at the national level divided by the weighted total number of households (using the full sample weights adjusted for nonresponse). A similar standardization procedure was followed in obtaining standardized weights for the individual women and under-5 questionnaires.

Sample weights were appended to all data sets and analyses were performed by weighting households, women or under-5s with these normalized sample weights.

Appendix D. Estimates of Sampling Errors

The sample of respondents selected in the Turkana County MICS is only one of the samples that could have been selected from the same population, using the same design and size. Each of these samples would yield results that differ somewhat from the results of the actual sample selected. Sampling errors are a measure of the variability between the estimates from all possible samples. The extent of variability is not known exactly, but can be estimated statistically from the survey data.

The following sampling error measures are presented in this appendix for each of the selected indicators:

Standard error (se): Standard error is the square root of the variance of the estimate. For survey indicators that are means, proportions or ratios, the Taylor series linearization method is used for the estimation of standard errors. For more complex statistics, such as fertility and mortality rates, the Jackknife repeated replication method is used for standard error estimation.

Coefficient of variation (se/r) is the ratio of the standard error to the value (r) of the indicator, and is a measure of the relative sampling error.

Design effect (deff) is the ratio of the actual variance of an indicator, under the sampling method used in the survey, to the variance calculated under the assumption of simple random sampling based on the same sample size. The *square root of the design effect (deft)* is used to show the efficiency of the sample design in relation to the precision. A *deft* value of 1.0 indicates that the sample design of the survey is as efficient as a simple random sample for a particular indicator, while a *deft* value above 1.0 indicates an increase in the standard error due to the use of a more complex sample design.

Confidence limits are calculated to show the interval within which the true value for the population can be reasonably assumed to fall, with a specified level of confidence. For any given statistic calculated from the survey, the value of that statistic will fall within a range of plus or minus two times the standard error ($r + 2.se$ or $r - 2.se$) of the statistic in 95 percent of all possible samples of identical size and design.

For the calculation of sampling errors from the MICS data, programs developed in CPro Version 5.0, SPSS Version 21 Complex Samples module and CMRJack¹²⁶ have been used.

The results are shown in the tables that follow. In addition to the sampling error measures described above, the tables also include weighted and unweighted counts of denominators for each indicator. Given the use of normalized weights, by comparing the weighted and unweighted counts it is possible to determine whether a particular domain has been under-sampled or over-sampled compared to the average sampling rate. If the weighted count is smaller than the unweighted count, this means that the particular domain had been over-sampled. As explained later in the footnote of Table SE.1, there is an exception in the case of indicators 4.1 and 4.3, for which the unweighted count represents the number of sample households, and the weighted counts reflect the total population.

¹²⁶ CMRJack is a software developed by FAFO, an independent and multidisciplinary research foundation. CMRJack produces mortality estimates and standard errors for surveys with complete birth histories or summary birth histories. See http://www.fafono.org/ais/child_mortality/index.html

Sampling errors are calculated for indicators of primary interest, for the county level, and for urban and rural areas within Turkana County. Three of the selected indicators are based on household members, eight are based on women, and two are based on children under-5 years. Table SE.1 shows the list of indicators for which sampling errors are calculated, including the base population (denominator) for each indicator. Tables SE.2 to SE.4 show the calculated sampling errors for selected domains.

Table SE.1: Indicators selected for sampling error calculations

List of indicators selected for sampling error calculations, and base populations (denominators) for each indicator, Turkana County MICS, 2013/14

MICS5 Indicator	Base Population
Household members	
4.1 Use of improved drinking water sources	All household members ^a
4.3 Use of improved sanitation	All household members ^a
7.4 Primary school net attendance ratio (adjusted)	Children of primary school age
Women	
5.3 Contraceptive prevalence rate	Women age 15-49 years who are currently married or in union
5.4 Unmet need	Women age 15-49 years who are currently married or in union
5.5a Antenatal care coverage (1+ times, skilled provider)	Women age 15-49 years with a live birth in the last 2 years
5.5b Antenatal care coverage (4+ times, any provider)	Women age 15-49 years with a live birth in the last 2 years
5.7 Skilled attendant at delivery	Women age 15-49 years with a live birth in the last 2 years
7.1 Literacy rate (young women)	Women age 15-24 years
9.1 Knowledge about HIV prevention (young women)	Women age 15-24 years
9.15 Condom use with non-regular partners	Women age 15-24 years who had a non-marital, non-cohabiting partner in the last 12 months
Under-5s	
3.18 Children under age 5 who slept under an ITN	Children under age 5 years who spent the previous night in the household
3.22 Anti-malarial treatment of children under age 5	Children under age 5 years with fever in the last 2 weeks
^a To calculate the weighted results of MICS Indicators 4.1 and 4.3, the household weight is multiplied by the number of household members in each household. Therefore the unweighted base population presented in the SE tables reflect the unweighted number of households, whereas the weighted numbers reflect the household population.	

Table SE.2: Sampling errors: Total sampleStandard errors, coefficients of variation, design effects (*deff*), square root of design effects (*deff*), and confidence intervals for selected indicators, Turkana County MICS,2013/14

	MICS Indicator	MDG Indicator	Value (<i>r</i>)	Standard error (<i>se</i>)	Coefficient of variation (<i>se/r</i>)	Design effect (<i>deff</i>)	Square root of design effect (<i>deff</i>)	Weighted count	Unweighted count	Confidence limits	
										Lower bound <i>r</i> - 2 <i>se</i>	Upper bound <i>r</i> + 2 <i>se</i>
Household members											
Use of improved drinking water sources	4.1	7.8	0.719	0.0334	0.046	7.028	2.651	6,594	1,277	0.652	0.785
Use of improved sanitation	4.3	7.9	0.123	0.0243	0.198	7.002	2.646	6,594	1,277	0.074	0.171
Primary school net attendance ratio (adjusted)	7.4	2.1	0.654	0.0241	0.037	4.202	2.050	1,606	1,641	0.606	0.702
Women											
Contraceptive prevalence rate	5.3	5.3	0.141	0.0186	0.132	1.848	1.360	615	649	0.103	0.178
Unmet need	5.4	5.6	0.340	0.0142	0.042	0.585	0.765	615	649	0.312	0.369
Antenatal care coverage (1+ times, skilled provider)	5.5a	5.5	0.9360	0.0099	0.011	0.670	0.819	387	412	0.916	0.956
Antenatal care coverage (4+ times, any provider)	5.5b	5.5	0.440	0.0312	0.071	1.623	1.274	387	412	0.378	0.502
Skilled attendant at delivery	5.7	5.2	0.351	0.0442	0.126	3.515	1.875	387	412	0.263	0.440
Literacy rate (young women)	7.1	2.3	0.547	0.0369	0.067	2.324	1.524	461	425	0.473	0.620
Knowledge about HIV prevention (young women)	9.1	6.3	0.347	0.0216	0.062	0.874	0.935	461	425	0.304	0.390
Condom use with non-regular partners	9.15	6.2	0.312	0.0486	0.156	0.714	0.845	78	66	0.215	0.409
Under-5s											
Children under age 5 who slept under an ITN	3.18	6.7	0.2470	0.0278	0.113	4.351	2.086	1,051	1,049	0.191	0.302
Anti-malarial treatment of children under age 5	3.22	6.8	0.175	0.0257	0.147	0.877	0.937	198	192	0.123	0.226
na: not applicable											

Table SE.3: Sampling errors: UrbanStandard errors, coefficients of variation, design effects (*deff*), square root of design effects (*deff*), and confidence intervals for selected indicators, Turkana County MICS, 2013/14

	MICS Indicator	MDG Indicator	Value (<i>r</i>)	Standard error (<i>se</i>)	Coefficient of variation (<i>se/r</i>)	Design effect (<i>deff</i>)	Square root of design effect (<i>deff</i>)	Weighted count	Unweighted count	Confidence limits	
										Lower bound <i>r</i> - 2 <i>se</i>	Upper bound <i>r</i> + 2 <i>se</i>
Household members											
Use of improved drinking water sources	4.1	7.8	0.855	0.0385	0.045	6.377	2.525	3,598	533	0.778	0.932
Use of improved sanitation	4.3	7.9	0.201	0.0392	0.196	5.110	2.260	3,598	533	0.122	0.279
Primary school net attendance ratio (adjusted)	7.4	2.1	0.821	0.0246	0.030	2.785	1.669	844	677	0.772	0.870
Women											
Contraceptive prevalence rate	5.3	5.3	0.227	0.0331	0.146	1.651	1.285	320	265	0.160	0.293
Unmet need	5.4	5.6	0.355	0.0193	0.054	0.430	0.656	320	265	0.317	0.394
Antenatal care coverage (1+ times, skilled provider)	5.5a	5.5	0.983	0.0052	0.005	0.274	0.524	199	169	0.973	0.994
Antenatal care coverage (4+ times, any provider)	5.5b	5.5	0.531	0.0462	0.087	1.437	1.199	199	169	0.439	0.624
Skilled attendant at delivery	5.7	5.2	0.565	0.0705	0.125	3.397	1.843	199	169	0.424	0.706
Literacy rate (young women)	7.1	2.3	0.648	0.0438	0.068	2.300	1.517	345	274	0.561	0.736
Knowledge about HIV prevention (young women)	9.1	6.3	0.364	0.0225	0.062	0.598	0.774	345	274	0.319	0.409
Condom use with non-regular partners	9.15	6.2	0.333	0.0535	0.161	0.645	0.803	67	51	0.226	0.440
Under-5s											
Children under age 5 who slept under an ITN	3.18	6.7	0.371	0.0502	0.135	4.497	2.120	538	418	0.271	0.472
Anti-malarial treatment of children under age 5	3.22	6.8	0.225	0.0442	0.197	0.965	0.982	114	87	0.136	0.313
na: not applicable											

Table SE.4: Sampling errors: RuralStandard errors, coefficients of variation, design effects (*deff*), square root of design effects (*deff*), and confidence intervals for selected indicators, Turkana County MICS, 2013/14

	MICS Indicator	MDG Indicator	Value (<i>r</i>)	Standard error (<i>se</i>)	Coefficient of variation (<i>se/r</i>)	Design effect (<i>deff</i>)	Square root of design effect (<i>deff</i>)	Weighted count	Unweighted count	Confidence limits	
										Lower bound <i>r</i> - 2 <i>se</i>	Upper bound <i>r</i> + 2 <i>se</i>
Household members											
Use of improved drinking water sources	4.1	7.8	0.554	0.0566	0.102	9.625	3.102	2,996	744	0.441	0.667
Use of improved sanitation	4.3	7.9	0.029	0.0154	0.531	6.264	2.503	2,996	744	0.000	0.060
Primary school net attendance ratio (adjusted)	7.4	2.1	0.469	0.0406	0.087	6.365	2.523	762	964	0.387	0.550
Women											
Contraceptive prevalence rate	5.3	5.3	0.047	0.0120	0.256	1.240	1.113	295	384	0.023	0.071
Unmet need	5.4	5.6	0.323	0.0212	0.065	0.783	0.885	295	384	0.281	0.366
Antenatal care coverage (1+ times, skilled provider)	5.5a	5.5	0.886	0.0206	0.023	1.021	1.010	188	243	0.845	0.927
Antenatal care coverage (4+ times, any provider)	5.5b	5.5	0.343	0.0457	0.133	2.241	1.497	188	243	0.252	0.435
Skilled attendant at delivery	5.7	5.2	0.126	0.0392	0.312	3.383	1.839	188	243	0.047	0.204
Literacy rate (young women)	7.1	2.3	0.244	0.0574	0.235	2.679	1.637	116	151	0.129	0.359
Knowledge about HIV prevention (young women)	9.1	6.3	0.297	0.0543	0.183	2.116	1.455	116	151	0.189	0.406
Condom use with non-regular partners	9.15	6.2	(*)	(*)	(*)	(*)	(*)	12	15	(*)	(*)
Under-5s											
Children under age 5 who slept under an ITN	3.18	6.7	0.117	0.0221	0.190	3.002	1.733	513	631	0.072	0.161
Anti-malarial treatment of children under age 5	3.22	6.8	0.107	0.0138	0.129	0.208	0.456	84	105	0.080	0.135
na: not applicable											

Appendix E. List of Personnel Involved in the Survey

Survey Management Team

PSRI

Murungaru Kimani, Director
Lawrence Ikamari, Director

KNBS

Zachary Mwangi, Director General
Macdonald Obudho, Director

UNICEF

Pirkko Heinonen, Representative a.i.
Kanyankore Marcel Rundasigwa, former Representative (RIP)
Madhavi Ashok, Deputy Representative
Joanne Bosworth, Chief of Social Policy
Paul Mpuga, Chief of PME

Technical Co-ordinators

PSRI

Alfred Agwanda
Samuel Wakibi
Anne Khasakhala
Ben Jarabi
Wanjiru Gichuhi
Andrew Mutuku
George Odipo

KNBS

Macdonald Obudho
James Ng'ang'a
James Munguti
William O. Nyongo, CSO

UNICEF

UNICEF

Paul Mpuga
Monica Chizororo
Nicholas Oloo
Robert Peter Ndugwa
John Ndegwa Wagai

Survey Support Team

UNICEF

Susan Govedi
Linda Claire
Moses Mwangi

UNICEF HQ/Regional Technical Backstopping Team

Bo Pedersen
Yadigar Coskun
Eva Quintana
Pierre Martel

Report Author (Consultant)

Nyasha Madzingira

Report Review Team

KNBS

Macdonald Obudho
Robert Buluma
Godfrey Otieno
James Ng'ang'a
Dickson A Makuba

UNICEF

Monica Chizororo
Nyasha Madzingira

PSRI

Lawrence Ikamari
Samuel Wakibi
Ben Jarabi

Ministry of Health

Samuel Murage
Charles Mabakha
Lydia Wanjiru Karimurio
John Wanyungu

National Registration Bureau

Immaculate K. Ndetei

NACC

Mercy Khasiani

Turkana County

Nancy Kinyonge
Wycliffe Machani
Joseph Orata

Kakamega County

Enoch Obuolo
Paul Manyasi
Ernest O. Odwori

Bungoma County

Thomas Shiundu
Hedwick Wasike
Alice Barasa

Data Collection Team**County Coordinator**

Wanjiru Gichuhi

Supervisors

Ezra Erupe
Susan Akuyo

Field Interviewers

Pauline Lokidor
Monica Ewoi
Irene Eno
Maureen Ekidor
Anne Kooil
Teresa Alimlim
Patrick Eyepan
Gladys Lokwawi
Deborah Namii
Alice Loukworo
Dorryshella Eyen
Loice Ekitela

Cluster updating field work**Coordinator**

Thomas W. Alubokho

Cartographer

Franco Mwendwa

Supervisors

Thomas Sitienei
Paul Tanui
David Kigen

Enumerators

Corby Tata Emoit
Muya Fredrick Lokol
Jeniffer N. Ekidor

Tioko S. Ekalimon
Paulo B. Chemjor
Joseph Kipkirui Rono

Data Processing**Data Entry Personnel**

Yvonne Chebet Ronoh
Melissa Ayuma Muyale
Alex Mutuku Judah
Eunice W. Maina
Redempta K. Muyuma
Esther K. Kinyanjui
Lilian Cheronu
Milcah W. Mwangangi
Domitilla M. Kivuvo
Hezbon Nango
Margaret Nyamuok
Daniel Mutembei Marete
Catherine Wakanyi Wangaruro
Habil Joash Onyango
Benedict Rono
Daniel Otieno Ochola

Data Cleaning and Validation

Bernard Obasi
Samuel Wakibi
John Ndegwa Wagai

Data Weighting

James Ng'ang'a

Data Analysts


Lawrence Ikamari
Alfred Agwanda
Murungaru Kimani
James Ng'ang'a
Samuel Wakibi
Anne Khasakhala
Ben Jarabi
Wanjiru Gichuhi
Andrew Mutuku
George Odipo
Bernard Obasi

Appendix F. Data Quality Tables

Table DQ.1: Age distribution of household population									
Single-year age distribution of household population by sex, Turkana County MICS, 2013/14									
	Males		Females		Age	Males		Females	
	Number	Percent	Number	Percent		Number	Percent	Number	Percent
Age					Age				
0	120	3.7	119	3.6	45	28	0.9	22	0.6
1	102	3.1	94	2.8	46	10	0.3	21	0.6
2	104	3.2	104	3.1	47	9	0.3	16	0.5
3	112	3.4	118	3.6	48	17	0.5	15	0.4
4	113	3.5	109	3.3	49	12	0.4	11	0.3
5	124	3.8	122	3.7	50	29	0.9	28	0.9
6	112	3.4	128	3.9	51	11	0.3	19	0.6
7	113	3.5	97	2.9	52	24	0.7	18	0.5
8	114	3.5	115	3.5	53	11	0.3	19	0.6
9	97	3.0	72	2.2	54	10	0.3	24	0.7
10	119	3.6	120	3.6	55	14	0.4	33	1.0
11	83	2.5	75	2.3	56	10	0.3	23	0.7
12	91	2.8	93	2.8	57	7	0.2	16	0.5
13	102	3.1	109	3.3	58	5	0.1	13	0.4
14	113	3.4	113	3.4	59	5	0.2	11	0.3
15	90	2.7	72	2.2	60	19	0.6	15	0.5
16	84	2.6	78	2.4	61	11	0.3	5	0.2
17	82	2.5	83	2.5	62	7	0.2	8	0.2
18	80	2.4	69	2.1	63	13	0.4	13	0.4
19	78	2.4	49	1.5	64	5	0.1	12	0.4
20	58	1.8	75	2.3	65	17	0.5	21	0.6
21	55	1.7	53	1.6	66	5	0.1	8	0.3
22	54	1.6	42	1.3	67	10	0.3	7	0.2
23	49	1.5	45	1.4	68	5	0.2	12	0.4
24	49	1.5	47	1.4	69	5	0.2	6	0.2
25	43	1.3	47	1.4	70	10	0.3	5	0.2
26	44	1.4	57	1.7	71	5	0.1	3	0.1
27	63	1.9	40	1.2	72	1	0.0	4	0.1
28	46	1.4	63	1.9	73	5	0.2	4	0.1
29	31	1.0	43	1.3	74	9	0.3	12	0.4
30	39	1.2	52	1.6	75	2	0.1	10	0.3
31	27	0.8	23	0.7	76	0	0.0	0	0.0
32	20	0.6	29	0.9	77	0	0.0	4	0.1
33	24	0.7	30	0.9	78	1	0.0	0	0.0
34	31	0.9	28	0.8	79	1	0.0	2	0.1
35	70	2.1	55	1.7	80	2	0.1	2	0.1
36	13	0.4	22	0.7	81	1	0.0	0	0.0
37	21	0.6	22	0.7	82	1	0.0	0	0.0
38	23	0.7	32	1.0	83	2	0.1	4	0.1
39	25	0.8	26	0.8	84	3	0.1	0	0.0
40	52	1.6	34	1.0	85+	3	0.1	2	0.0
41	13	0.4	14	0.4					
42	8	0.2	20	0.6	DK/Missing	2	0.1	0	0.0
43	13	0.4	13	0.4					

44	24	0.7	19	0.6	Total	3,274	100.0	3,321	100.0
----	----	-----	----	-----	-------	-------	-------	-------	-------

Figure DQ.1: Household population by single ages, Turkana County MICS, 2013/14



Note: The figure excludes 2 household members with unknown age and/or sex

Table DQ.2: Age distribution of eligible and interviewed women

Household population of women age 10-54 years, interviewed women age 15-49 years, and percentage of eligible women who were interviewed, by five-year age groups, Turkana County MICS, 2013/14

Age	Household population of women age 10-54 years	Interviewed women age 15-49 years		Percentage of eligible women interviewed (Completion rate)
	Number	Number	Percent	
10-14	511	na	na	na
15-19	351	266	22.9	75.8
20-24	262	221	19.0	84.5
25-29	249	221	19.0	88.8
30-34	161	151	13.0	93.6
35-39	157	141	12.1	89.8
40-44	100	86	7.4	85.9
45-49	85	79	6.7	92.8
50-54	109	na	na	na
Total (15-49)	1,365	1,165	100.0	85.4
Ratio of 50-54 to 45-49	1.29	na	na	na

na: not applicable

Table DQ.4: Age distribution of children in household and under-5 questionnaires

Household population of children age 0-7 years, children age 0-4 years whose mothers/caretakers were interviewed, and percentage of under-5 children whose mothers/caretakers were interviewed, by single years of age, Turkana County MICS, 2013/14

	<u>Household population of children 0-7 years</u>	<u>Under-5s with completed interviews</u>		Percentage of eligible under-5s with completed interviews (Completion rate)
	Number	Number	Percent	
Age				
0	239	230	22.0	96.2
1	196	191	18.3	97.5
2	208	198	19.0	95.3
3	231	219	20.9	94.7
4	222	207	19.8	93.3
5	246	na	na	na
6	240	na	na	na
7	210	na	na	na
Total (0-4)	1,095	1,045	100.0	95.4
Ratio of 5 to 4	1.11	na	na	na
na: not applicable				

Table DQ.5: Birth date reporting: Household population

Percent distribution of household population by completeness of date of birth information, Turkana County MICS, 2013/14

	<u>Completeness of reporting of month and year of birth</u>				Total	Number of household members
	Year and month of birth	Year of birth only	Month of birth only	Both missing		
Total	41.9	58.0	0.0	0.1	100.0	6,594
Age						
0-4	82.9	17.1	0.0	0.0	100.0	1,095
5-14	50.2	49.7	0.0	0.1	100.0	2,114
15-24	38.6	61.4	0.0	0.0	100.0	1,292
25-49	17.2	82.7	0.0	0.1	100.0	1,459
50-64	9.0	90.9	0.0	0.1	100.0	440
65-84	0.8	99.2	0.0	0.0	100.0	188
85+	16.8	83.2	0.0	0.0	100.0	4
DK/Missing	0.0	0.0	0.0	100.0	100.0	2
Area						
Urban	50.8	49.1	0.0	0.1	100.0	3,598
Rural	31.2	68.8	0.0	0.0	100.0	2,996
na: not applicable						

Table DQ.6: Birth date and age reporting: Women

Percent distribution of women age 15-49 years by completeness of date of birth/age information, Turkana County MICS, 2013/14

	Completeness of reporting of date of birth and age					Total	Number of women age 15-49 years
	Year and month of birth	Year of birth and age	Year of birth only	Age only	Other/DK/Missing		
Total	27.9	71.9	0.0	0.0	0.2	100.0	1,104
Area							
Urban	40.0	59.9	0.0	0.0	0.2	100.0	683
Rural	8.3	91.4	0.0	0.0	0.3	100.0	421

Table DQ.8: Birth date and age reporting: Under-5s

Percent distribution children under 5 years by completeness of date of birth/age information, Turkana County MICS, 2013/14

	Completeness of reporting of date of birth and age					Total	Number of men age 15-49 years
	Year and month of birth	Year of birth and age	Year of birth only	Age only	Other/DK/Missing		
Total	86.8	13.2	0.0	0.0	0.0	100.0	1,067
Area							
Urban	92.5	7.5	0.0	0.0	0.0	100.0	546
Rural	80.8	19.2	0.0	0.0	0.0	100.0	521

Table DQ.9: Birth date reporting: Children, adolescents and young people

Percent distribution of children, adolescents and young people age 5-24 years by completeness of date of birth information, Turkana County MICS, 2013/14

	Completeness of reporting of month and year of birth				Total	Number of children, adolescents and young people age 5-24 years
	Year and month of birth	Year of birth only	Month of birth only	Both missing		
Total	45.8	54.1	0.0	0.1	100.0	3,406
Area						
Urban	54.1	45.8	0.0	0.1	100.0	1,973
Rural	34.4	65.6	0.0	0.0	100.0	1,433

Table DQ.10: Birth date reporting: First and last births

Percent distribution of first and last births to women age 15-49 years by completeness of date of birth, Turkana County MICS, 2013/14

	Completeness of reporting of date of birth											
	Date of first birth					Number of first births	Date of last birth					Number of last births
	Year and month of birth	Year of birth only	Completed since first birth only	Other/DK/Missing	Total		Year and month of birth	Year of birth only	Other/DK/Missing	Total		
Total	64.3	35.1	0.2	0.4	100.0	800	80.5	19.2	0.2	100.0	683	
Area												
Urban	75.2	24.3	0.0	0.4	100.0	455	86.1	13.9	0.0	100.0	369	
Rural	49.8	49.3	0.5	0.4	100.0	345	74.0	25.5	0.5	100.0	314	

Table DQ.11: Completeness of reporting

Percentage of observations that are missing information for selected questions and indicators, Turkana County MICS, 2013/14

Questionnaire and type of missing information	Reference group	Percent with missing/incomplete information ^a	Number of cases
Household			
Salt test result	All households interviewed that have salt	1.1	1,277
Starting time of interview	All households interviewed	0.2	1,277
Ending time of interview	All households interviewed	0.1	1,277
Women			
Date of first marriage/union	All ever married women age 15-49		
Only month		58.0	790
Both month and year		26.0	790
Age at first marriage/union	All ever married women age 15-49 with year of first marriage not known	2.1	790
Age at first intercourse	All women age 15-24 who have ever had sex	1.6	270
Time since last intercourse	All women age 15-24 who have ever had sex	1.1	270
Starting time of interview	All women interviewed	0.0	1,104
Ending time of interview	All women interviewed	0.0	1,104
Under-5			
Starting time of interview	All under-5 children	0.0	1,067
Ending time of interview	All under-5 children	0.3	1,067

^a Includes "Don't know" responses

Table DQ.12: Completeness of information for anthropometric indicators: Underweight

Percent distribution of children under 5 by completeness of information on date of birth and weight, Turkana County MICS, 2013/14

	Valid weight and date of birth	Reason for exclusion from analysis				Total	Percent of children excluded from analysis	Number of children under 5
		Weight not measured	Incomplete date of birth	Weight not measured and incomplete date of birth	Flagged cases (outliers)			
Total	83.9	2.9	12.6	0.6	0.1	100.0	16.1	1,067
Age								
<6 months	92.0	5.4	2.6	0.0	0.0	100.0	8.0	107
6-11 months	96.2	0.9	2.9	0.0	0.0	100.0	3.8	120
12-23 months	91.1	4.1	4.6	0.0	0.2	100.0	8.9	196
24-35 months	82.6	2.4	14.7	0.0	0.2	100.0	17.4	205
36-47 months	85.3	1.1	12.9	0.8	0.0	100.0	14.7	222
48-59 months	66.4	3.8	27.8	2.0	0.0	100.0	33.6	217

Table DQ.13: Completeness of information for anthropometric indicators: Stunting

Percent distribution of children under 5 by completeness of information on date of birth and length or height, Turkana County MICS, 2013/14

	Valid length/height and date of birth	Reason for exclusion from analysis				Total	Percent of children excluded from analysis	Number of children under 5
		Length/Height not measured	Incomplete date of birth	Length/Height not measured, incomplete date of birth	Flagged cases (outliers)			
Total	82.1	2.9	12.7	0.5	1.9	100.0	17.9	1,067
Age								
<6 months	89.2	6.3	2.6	0.0	1.9	100.0	10.8	107
6-11 months	92.6	0.9	2.9	0.0	3.6	100.0	7.4	120
12-23 months	89.2	4.1	4.6	0.0	2.1	100.0	10.8	196
24-35 months	80.0	2.1	14.7	0.0	3.2	100.0	20.0	205
36-47 months	83.9	1.1	13.2	0.5	1.3	100.0	16.1	222
48-59 months	66.4	3.8	27.8	2.0	0.0	100.0	33.6	217

Table DQ.14: Completeness of information for anthropometric indicators: Wasting

Percent distribution of children under 5 by completeness of information on weight and length or height, Turkana County MICS, 2013/14

	Reason for exclusion from analysis					Total	Percent of children excluded from analysis	Number of children under 5
	Valid weight and length/height	Weight not measured	Length/Height not measured	Weight and length/height not measured	Flagged cases (outliers)			
Total	93.8	0.1	0.1	3.3	2.7	100.0	6.2	1,067
Age								
<6 months	89.2	0.0	0.9	5.4	4.5	100.0	10.8	107
6-11 months	91.5	0.0	0.0	0.9	7.6	100.0	8.5	120
12-23 months	94.2	0.0	0.0	4.1	1.6	100.0	5.8	196
24-35 months	96.8	0.3	0.0	2.1	0.8	100.0	3.2	205
36-47 months	97.3	0.3	0.0	1.5	0.8	100.0	2.7	222
48-59 months	90.4	0.0	0.0	5.8	3.8	100.0	9.6	217

Table DQ.15: Heaping in anthropometric measurements

Distribution of weight and height/length measurements by digits reported for the decimal points, Turkana County MICS, 2013/14

	Weight		Height or length	
	Number	Percent	Number	Percent
Total	1,030	100.0	1,032	100.0
Digits				
0	122	11.8	124	12.0
1	110	10.6	78	7.6
2	117	11.4	128	12.5
3	108	10.5	112	10.9
4	97	9.4	98	9.5
5	96	9.3	177	17.2
6	91	8.9	88	8.6
7	113	10.9	103	10.0
8	94	9.2	62	6.0
9	82	8.0	61	5.9
0 or 5	218	21.2	301	29.2

Figure DQ.2: Weight and height/length measurements by digits reported for the decimal points, Turkana County MICS, 2013/14




Table DQ.16: Observation of birth certificates

Percent distribution of children under 5 by presence of birth certificates, and percentage of birth certificates seen, Turkana County MICS, 2013/14

	Child has birth certificate		Child does not have birth certificate	DK/Missing	Total	Percentage of birth certificates seen by the interviewer (1)/(1+2)*100	Number of children under age 5
	Seen by the interviewer (1)	Not seen by the interviewer (2)					
Total	13.7	8.5	77.3	0.5	100.0	61.8	1,067
Area							
Urban	21.2	12.1	66.2	0.5	100.0	63.7	546
Rural	5.8	4.6	89.0	0.6	100.0	55.5	521
Child's age							
0-5 months	8.9	6.6	84.5	0.0	100.0	57.3	107
6-11 months	14.7	9.9	75.4	0.0	100.0	59.9	120
12-23 months	13.1	7.4	79.5	0.0	100.0	63.8	196
24-35 months	13.9	9.5	76.1	0.4	100.0	59.4	205
36-47 months	18.1	9.5	71.5	0.9	100.0	65.6	222
48-59 months	11.3	7.5	79.9	1.3	100.0	60.2	217

Table DQ.17: Observation of vaccination cards

Percent distribution of children age 0-35 months by presence of a vaccination card, and the percentage of vaccination cards seen by the interviewers, Turkana County MICS, 2013/14

	Child does not have vaccination card		Child has vaccination card		DK/Missing	Total	Percentage of vaccination cards seen by the interviewer (1)/(1+2)*100	Number of children age 0-35 months
	Had vaccination card previously	Never had vaccination card	Seen by the interviewer (1)	Not seen by the interviewer (2)				
Total	5.5	4.5	59.6	30.3	0.1	100.0	66.3	627
Area								
Urban	5.8	1.2	62.4	30.4	0.3	100.0	67.3	321
Rural	5.2	8.0	56.5	30.2	0.0	100.0	65.1	306
Child's age								
0-5 months	2.3	12.3	67.5	17.1	0.8	100.0	79.8	107
6-11 months	1.3	4.1	74.4	20.2	0.0	100.0	78.7	120
12-23 months	6.1	2.0	64.2	27.8	0.0	100.0	69.8	196
24-35 months	8.9	3.2	42.4	45.5	0.0	100.0	48.2	205

Table DQ.18: Observation of women's health cards

Percent distribution of women with a live birth in the last 2 years by presence of a health card, and the percentage of health cards seen by the interviewers, Turkana County MICS, 2013/14

	Woman has health card			DK/Missing	Total	Percent of health cards seen by the interviewer (1)/(1+2)*100	Number of women with a live birth in the last two years
	Woman does not have health card	Seen by the interviewer (1)	Not seen by the interviewer (2)				
Total	11.0	60.6	27.6	0.8	100.0	68.7	387
Area							
Urban	4.2	73.0	22.0	0.7	100.0	76.8	199
Rural	18.2	47.4	33.4	1.0	100.0	58.7	188
Age							
15-24	3.5	69.6	26.9	0.0	100.0	72.1	112
25-34	12.5	59.1	27.5	0.9	100.0	68.2	199
35-49	17.9	51.5	28.8	1.9	100.0	64.1	77

Table DQ.19: Observation of bednets and places for handwashing

Percentage of bednets in all households observed by the interviewers, and percent distribution of places for handwashing observed by the interviewers in all interviewed households, Turkana County MICS, 2013/14

	Percentage of bednets observed by interviewer	Total number of bednets	Place for handwashing				Total	Number of households interviewed
			Observed	Not observed				
				Not in the dwelling, plot or yard	No permission to see	Other reason		
Total	77.8	953	4.3	87.0	3.9	4.7	100.0	1,277
Area								
Urban	76.0	722	5.9	85.0	2.6	6.5	100.0	684
Rural	83.5	231	2.6	89.3	5.3	2.7	100.0	593
Wealth index quintile								
Poorest	82.4	62	1.8	85.3	7.8	5.1	100.0	256
Second	80.7	92	4.1	88.6	6.0	1.0	100.0	282
Middle	80.8	121	2.8	90.0	2.8	4.4	100.0	255
Fourth	85.4	209	5.5	87.2	2.2	5.1	100.0	239
Richest	72.5	469	7.7	83.7	0.0	8.7	100.0	245

Table DQ.20: Respondent to the under-5 questionnaire

Distribution of children under five by respondent to the under-5 questionnaire, Turkana County MICS, 2013/14

	Mother in the household	Mother not in the household and primary caretaker identified:				Total	Number of children under 5
		Father	Other adult female	Other adult male			
Total	92.7	0.2	6.8	0.3	100.0	1,095	
Age							
0	98.5	0.2	1.3	0.0	100.0	239	
1	97.2	0.0	2.8	0.0	100.0	196	
2	90.7	0.2	9.1	0.0	100.0	208	
3	91.8	0.7	7.5	0.0	100.0	231	
4	85.2	0.0	13.5	1.3	100.0	222	

Table DQ.21: Selection of children age 1-17 years for the child labour and child discipline modules

Percent distribution of households by the number of children age 1-17 years, and the percentage of households with at least two children age 1-17 years where correct selection of one child for the child labour and child discipline modules was performed, Turkana County MICS, 2013/14

	Number of children age 1-17 years			Total	Number of households	Percentage of households where correct selection was performed	Number of households with 2 or more children age 1- 17 years
	None	One	Two or more				
Total	17.2	14.3	68.5	100.0	1,277	95.8	875
Area							
Urban	18.8	14.1	67.1	100.0	684	94.9	459
Rural	15.3	14.6	70.1	100.0	593	96.8	416
Wealth index quintile							
Poorest	15.5	13.5	71.0	100.0	256	97.8	182
Second	15.8	15.9	68.3	100.0	282	96.0	193
Middle	15.9	13.9	70.2	100.0	255	95.7	179
Fourth	18.0	12.9	69.1	100.0	239	93.6	165
Richest	20.9	15.2	63.8	100.0	245	95.9	157

Table DQ.22: School attendance by single age

Distribution of household population age 5-24 years by educational level and grade attended in the current (or most recent) school year, Turkana County MICS, 2013/14

Age at beginning of school year	Not attending school	Currently attending																	Number of household members		
		Preschool	Primary school Grade										Secondary school Grade				Higher than secondary	DK/Missing		Total	
			1	2	3	4	5	6	7	8	DK/Missing	1	2	3	4						
5	34.3	40.6	13.1	8.3	1.5	0.2	0.4	0.0	0.8	0.0	0.8	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	100.0	243
6	29.4	27.0	23.5	10.1	5.2	2.6	0.0	0.0	0.0	0.0	1.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	100.0	202
7	30.4	15.6	19.5	20.3	8.7	2.8	1.6	0.0	0.0	0.0	0.8	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3	100.0	232
8	31.3	4.6	12.8	20.2	17.8	9.1	3.5	0.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	100.0	180
9	27.9	2.1	5.3	20.3	24.7	12.2	2.7	2.9	1.0	0.0	0.4	0.0	0.5	0.0	0.0	0.0	0.0	0.0	0.0	100.0	219
10	23.8	1.4	2.9	15.4	17.5	18.2	12.9	4.3	3.0	0.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	100.0	169
11	24.9	2.1	3.9	13.1	14.1	19.4	8.6	8.5	5.0	0.0	0.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	100.0	173
12	26.6	0.0	1.5	4.7	11.7	19.3	13.9	11.0	6.5	3.6	0.3	0.9	0.0	0.0	0.0	0.0	0.0	0.0	0.0	100.0	230
13	26.1	0.0	0.7	4.5	6.3	6.1	14.9	15.8	14.7	10.4	0.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	100.0	200
14	28.1	0.0	0.0	2.9	1.5	9.5	11.8	13.6	14.4	13.3	0.0	1.5	3.4	0.0	0.0	0.0	0.0	0.0	0.0	100.0	174
15	30.9	0.0	0.0	0.0	0.5	4.2	8.2	11.1	22.1	10.3	0.0	4.6	6.9	1.3	0.0	0.0	0.0	0.0	0.0	100.0	153
16	26.3	0.0	0.0	0.0	0.4	1.1	1.3	6.3	18.4	13.8	0.0	4.9	13.1	7.2	7.2	0.0	0.0	0.0	0.0	100.0	157
17	36.8	0.0	1.1	0.7	0.0	1.5	0.7	3.0	9.3	20.0	0.0	3.5	8.1	6.7	6.0	1.8	1.1	100.0	162	162	
18	43.6	0.0	0.0	0.0	0.0	2.7	0.9	3.9	8.4	8.5	0.0	4.5	11.4	11.9	2.8	1.4	0.0	100.0	129	129	
19	58.0	0.0	0.0	0.6	0.0	0.0	0.0	2.1	6.9	7.1	0.0	3.8	5.4	3.2	9.4	3.5	0.0	100.0	126	126	
20	61.3	0.0	0.0	0.0	0.0	1.6	0.0	0.0	3.5	3.2	0.0	4.3	3.3	7.0	10.0	5.7	0.0	100.0	111	111	
21	50.9	0.0	0.0	0.0	0.0	0.0	2.0	0.6	3.1	4.0	0.0	3.2	8.0	9.7	8.3	8.7	1.5	100.0	87	87	
22	76.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2.0	2.5	1.1	1.3	2.2	3.8	4.2	6.9	0.0	100.0	112	112	
23	81.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.2	1.5	1.0	15.2	0.0	100.0	81	81	
24 ^a	100.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	100.0	2	2	

^a Those age 25 at the time of interview who were age 24 at beginning of school year are excluded as current attendance was only collected for those age 5-24 at the time of interview

Table DQ.23: Sex ratio at birth among children ever born and living

Sex ratio (number of males per 100 females) among children ever born (at birth), children living, and deceased children, by age of women, Turkana County MICS, 2013/14

	Children Ever Born			Children Living			Children Deceased			Number of women
	Sons	Daughters	Sex ratio at birth	Sons	Daughters	Sex ratio	Sons	Daughters	Sex ratio	
Total	1,579	1,469	1.07	1,458	1,369	1.07	121	101	1.21	1,104
Age										
15-19	27	15	1.75	25	15	1.63	2	0	-	252
20-24	120	118	1.02	115	111	1.04	5	8	0.67	209
25-29	328	295	1.11	316	284	1.11	12	11	1.10	210
30-34	308	305	1.01	291	287	1.01	17	18	0.93	144
35-39	330	312	1.06	304	293	1.04	26	19	1.36	134
40-44	239	208	1.15	215	189	1.13	24	19	1.31	82
45-49	227	215	1.06	192	189	1.02	35	26	1.34	75

Table DQ.24: Births by periods preceding the survey

Number of births, sex ratio at birth, and period ratio by periods preceding the survey, according to living, deceased, and total children (imputed), as reported in the birth histories, Turkana County MICS, 2013/14

	Number of births			Percent with complete birth date ^a			Sex ratio at birth ^b			Period ratio ^c		
	Living	Deceased	Total	Living	Deceased	Total	Living	Deceased	Total	Living	Deceased	Total
Total	2,827	222	3,048	65.5	36.5	63.4	106.5	120.7	107.5	na	na	na
Years												
0	207	4	211	94.4	86.5	94.2	111.7	22.8	108.9	na	na	na
1	179	7	186	92.8	89.4	92.7	107.5	172.2	109.3	97.0	79.5	96.3
2	163	13	176	84.6	77.5	84.1	102.8	65.1	99.6	88.6	161.1	91.6
3	189	9	198	83.6	47.5	81.9	110.2	297.2	114.9	112.8	76.8	110.4
4	172	11	183	67.0	47.1	65.8	113.0	75.9	110.3	93.3	116.0	94.4
5	180	10	190	70.8	45.2	69.5	84.6	96.0	85.2	101.2	112.1	101.7
6	183	7	190	66.4	19.9	64.7	92.1	na	99.3	115.6	79.2	113.7
7	137	7	145	65.4	55.8	64.9	122.3	74.6	119.3	77.7	95.9	78.5
8	170	8	178	64.0	0.0	61.1	90.7	272.3	95.1	138.9	113.2	137.4
9	108	7	115	60.0	44.0	59.0	134.9	55.8	127.6	16.4	9.8	15.8
10+	1,137	139	1,276	49.7	28.2	47.4	109.3	122.1	110.6	na	na	na
Five-year periods												
0-4	911	43	954	84.9	65.9	84.0	109.2	98.4	108.6	na	na	na
5-9	778	40	818	65.8	33.2	64.3	99.7	145.5	101.5	na	na	na
10-14	592	42	634	54.4	52.2	54.3	100.6	95.0	100.3	na	na	na
15-19	304	43	347	52.2	24.2	48.7	112.1	104.5	111.1	na	na	na
20+	241	54	295	35.1	12.7	31.0	129.8	169.8	136.2	na	na	na

na: not applicable

^a Both month and year of birth given. The inverse of the percent reported is the percent with incomplete and therefore imputed date of birth^b $(B_m/B_f) \times 100$, where B_m and B_f are the numbers of male and female births, respectively^c $(2 \times B_t / (B_{t-1} + B_{t+1})) \times 100$, where B_t is the number of births in year t preceding the survey

Table DQ.25: Reporting of age at death in days

Distribution of reported deaths under one month of age by age at death in days and the percentage of neonatal deaths reported to occur at ages 0–6 days, by 5-year periods preceding the survey (imputed), Turkana County MICS, 2013/14

	Number of years preceding the survey				Total (0–19)
	0–4	5–9	10–14	15–19	
Age at death (days)					
0	0	0	0	1	1
1	7	2	5	6	20
2	0	1	1	0	1
3	0	3	0	0	3
4	3	0	0	0	3
5	0	0	0	1	1
7	1	0	2	0	2
16	0	1	0	0	1
Total 0–30 days	11	6	7	8	33
Percent early neonatal ^a	93.9	81.1	77.1	100.0	88.9

^a Deaths during the first 7 days (0-6), divided by deaths during the first month (0-30 days)

Table DQ.26: Reporting of age at death in months

Distribution of reported deaths under two years of age by age at death in months and the percentage of infant deaths reported to occur at age under one month, for the 5-year periods of birth preceding the survey (imputed), Turkana County MICS, 2013/14

	Number of years preceding the survey				Total (0-19)
	0–4	5–9	10–14	15–19	
Age at death (months)					
0 ^a	11	6	7	8	33
1	4	1	2	0	7
2	2	2	1	2	7
3	1	2	3	1	7
4	2	3	0	0	5
5	1	0	2	0	2
6	1	2	0	0	3
7	0	1	0	0	1
8	0	1	0	0	1
9	3	2	1	2	8
10	1	0	0	0	1
14	0	0	0	0	0
Reported as 1 year	5	2	8	4	19
Total 0–11 months	25	20	17	13	75
Percent neonatal ^b	44.3	32.7	43.9	58.4	43.6

^a Includes deaths under one month reported in days
^b Deaths under one month, divided by deaths under one year

Appendix G. Turkana County MICS5 Indicators: Numerators and Denominators

MICS INDICATOR		Module ¹²⁷	Numerator	Denominator	MDG Indicator Reference ¹²⁸
MORTALITY¹²⁹					
1.1	Neonatal mortality rate	BH	Probability of dying within the first month of life		
1.2	Infant mortality rate	CM - BH	Probability of dying between birth and the first birthday		MDG 4.2
1.3	Post-neonatal mortality rate	BH	Difference between infant and neonatal mortality rates		
1.4	Child mortality rate	BH	Probability of dying between the first and the fifth birthdays		
1.5	Under-five mortality rate	CM - BH	Probability of dying between birth and the fifth birthday		MDG 4.1

NUTRITION					
2.1a 2.1b	Underweight prevalence	AN	Number of children under age 5 who (a) fall below minus two standard deviations (moderate and severe) (b) fall below minus three standard deviations (severe) of the median weight for age of the WHO standard	Total number of children under age 5	MDG 1.8
2.2a 2.2b	Stunting prevalence	AN	Number of children under age 5 who (a) fall below minus two standard deviations (moderate and severe) (b) fall below minus three standard deviations (severe) of the median height for age of the WHO standard	Total number of children under age 5	
2.3a 2.3b	Wasting prevalence	AN	Number of children under age 5 who (a) fall below minus two standard deviations (moderate and severe) (b) fall below minus three standard deviations (severe) of the median weight for height of the WHO standard	Total number of children under age 5	
2.4	Overweight prevalence	AN	Number of children under age 5 who are above two standard deviations of	Total number of children under age 5	

¹²⁷Some indicators are constructed by using questions in several modules in the MICS questionnaires. In such cases, only the module(s) which contains most of the necessary information is indicated.

¹²⁸Millennium Development Goals (MDG) indicators, effective 15 January 2008 - <http://mdgs.un.org/unsd/mdg/Host.aspx?Content=Indicators/OfficialList.htm>, accessed 10 June 2013.

¹²⁹When the Birth History module is used, mortality indicators are calculated for the last 5-year period. When the indicators are estimated indirectly (using the Fertility module only), the rates refer to dates as estimated by the indirect technique.

			the median weight for height of the WHO standard		
2.5	Children ever breastfed	MN	Number of women with a live birth in the last 2 years who breastfed their last live-born child at any time	Total number of women with a live birth in the last 2 years	
2.6	Early initiation of breastfeeding	MN	Number of women with a live birth in the last 2 years who put their last newborn to the breast within one hour of birth	Total number of women with a live birth in the last 2 years	
2.7	Exclusive breastfeeding under 6 months	BD	Number of infants under 6 months of age who are exclusively breastfed ¹³⁰	Total number of infants under 6 months of age	
2.8	Predominant breastfeeding under 6 months	BD	Number of infants under 6 months of age who received breast milk as the predominant source of nourishment ¹³¹ during the previous day	Total number of infants under 6 months of age	
2.9	Continued breastfeeding at 1 year	BD	Number of children age 12-15 months who received breast milk during the previous day	Total number of children age 12-15 months	
2.10	Continued breastfeeding at 2 years	BD	Number of children age 20-23 months who received breast milk during the previous day	Total number of children age 20-23 months	
2.11	Duration of breastfeeding	BD	The age in months when 50 percent of children age 0-35 months did not receive breast milk during the previous day		
2.12	Age-appropriate breastfeeding	BD	Number of children age 0-23 months appropriately fed ¹³² during the previous day	Total number of children age 0-23 months	
2.13	Introduction of solid, semi-solid or soft foods	BD	Number of infants age 6-8 months who received solid, semi-solid or soft foods during the previous day	Total number of infants age 6-8 months	
2.14	Milk feeding frequency for non-breastfed children	BD	Number of non-breastfed children age 6-23 months who received at least 2 milk feedings during the previous day	Total number of non-breastfed children age 6-23 months	
2.15	Minimum meal frequency	BD	Number of children age 6-23 months who received solid, semi-solid and soft foods (plus milk feeds for non-breastfed children) the minimum number of times ¹³³ or more during the previous day	Total number of children age 6-23 months	

¹³⁰Infants receiving breast milk, and not receiving any other fluids or foods, with the exception of oral rehydration solution, vitamins, mineral supplements and medicines

¹³¹Infants who receive breast milk and certain fluids (water and water-based drinks, fruit juice, ritual fluids, oral rehydration solution, drops, vitamins, minerals, and medicines), but do not receive anything else (in particular, non-human milk and food-based fluids)

¹³²Infants age 0-5 months who are exclusively breastfed, and children age 6-23 months who are breastfed and ate solid, semi-solid or soft foods

¹³³Breastfeeding children: Solid, semi-solid, or soft foods, two times for infants age 6-8 months, and three times for children 9-23 months; Non-breastfeeding children: Solid, semi-solid, or soft foods, or milk feeds, four times for children age 6-23 months

2.16	Minimum dietary diversity	BD	Number of children age 6–23 months who received foods from 4 or more food groups ¹³⁴ during the previous day	Total number of children age 6–23 months	
2.17a 2.17b	Minimum acceptable diet	BD	(a) Number of breastfed children age 6–23 months who had at least the minimum dietary diversity and the minimum meal frequency during the previous day (b) Number of non-breastfed children age 6–23 months who received at least 2 milk feedings and had at least the minimum dietary diversity not including milk feeds and the minimum meal frequency during the previous day	(a) Number of breastfed children age 6–23 months (b) Number of non-breastfed children age 6–23 months	
2.18	Bottle feeding	BD	Number of children age 0-23 months who were fed with a bottle during the previous day	Total number of children age 0-23 months	
2.19	Iodized salt consumption	SI	Number of households with salt testing 15 parts per million or more of iodide/iodate	Total number of households in which salt was tested or where there was no salt	
2.20	Low-birthweight infants	MN	Number of most recent live births in the last 2 years weighing below 2,500 grams at birth	Total number of most recent live births in the last 2 years	
2.21	Infants weighed at birth	MN	Number of most recent live births in the last 2 years who were weighed at birth	Total number of most recent live births in the last 2 years	

CHILD HEALTH

3.1	Tuberculosis immunization coverage	IM	Number of children age 12-23 months who received BCG vaccine before their first birthday	Total number of children age 12-23 months	
3.2	Polio immunization coverage	IM	Number of children age 12-23 months who received the third dose of OPV vaccine (OPV3) before their first birthday	Total number of children age 12-23 months	
3.3	Diphtheria, pertussis and tetanus (DPT) immunization coverage	IM	Number of children age 12-23 months who received the third dose of DPT vaccine (DPT3) before their first birthday	Total number of children age 12-23 months	
3.4	Measles immunization coverage ¹³⁵	IM	Number of children age 12-23 months who received measles vaccine before their first birthday	Total number of children age 12-23 months	MDG 4.3
3.5	Hepatitis B immunization coverage	IM	Number of children age 12-23 months who received the third dose of Hepatitis B vaccine (HepB3) by their first birthday	Total number of children age 12-23 months	
3.6	Haemophilus influenzae type b (Hib) immunization coverage	IM	Number of children age 12-23 months who received the third dose of Hib vaccine (Hib3) by their first birthday	Total number of children age 12-23 months	

¹³⁴The indicator is based on consumption of any amount of food from at least 4 out of the 7 following food groups: 1) grains, roots and tubers, 2) legumes and nuts, 3) dairy products (milk, yogurt, cheese), 4) flesh foods (meat, fish, poultry and liver/organ meats), 5) eggs, 6) vitamin-A rich fruits and vegetables, and 7) other fruits and vegetables

3.7	Yellow fever immunization coverage	IM	Number of children age 12-23 months who received yellow fever vaccine by their first birthday	Total number of children age 12-23 months	
3.8	Full immunization coverage	IM	Number of children age 12-23 months who received all vaccinations recommended in the national immunization schedule before their first birthday	Total number of children age 12-23 months	
3.9	Neonatal tetanus protection	MN	Number of women age 15-49 years with a live birth in the 2 years preceding the survey who were given at least two doses of tetanus toxoid vaccine within the appropriate interval ¹³⁶ prior to the most recent birth	Total number of women age 15-49 years with a live birth in the 2 years preceding the survey	
3.10	Care-seeking for diarrhoea	CA	Number of children under age 5 with diarrhoea in the last 2 weeks for whom advice or treatment was sought from a health facility or provider	Total number of children under age 5 with diarrhoea in the last 2 weeks	
3.11	Diarrhoea treatment with oral rehydration salts (ORS) and zinc	CA	Number of children under age 5 with diarrhoea in the last 2 weeks who received ORS and zinc	Total number of children under age 5 with diarrhoea in the last 2 weeks	
3.12	Diarrhoea treatment with oral rehydration therapy (ORT) and continued feeding	CA	Number of children under age 5 with diarrhoea in the last 2 weeks who received ORT (ORS packet, pre-packaged ORS fluid, recommended homemade fluid or increased fluids) and continued feeding during the episode of diarrhoea	Total number of children under age 5 with diarrhoea in the last 2 weeks	
3.13	Care-seeking for children with acute respiratory infection (ARI) symptoms	CA	Number of children under age 5 with ARI symptoms in the last 2 weeks for whom advice or treatment was sought from a health facility or provider	Total number of children under age 5 with ARI symptoms in the last 2 weeks	
3.14	Antibiotic treatment for children with ARI symptoms	CA	Number of children under age 5 with ARI symptoms in the last 2 weeks who received antibiotics	Total number of children under age 5 with ARI symptoms in the last 2 weeks	
3.15	Use of solid fuels for cooking	HC	Number of household members in households that use solid fuels as the primary source of domestic energy to cook	Total number of household members	
3.16a 3.16b	Household availability of insecticide-treated nets (ITNs) ¹³⁷	TN	Number of households with (a) at least one ITN (b) at least one ITN for every two people	Total number of households	

¹³⁶See the MICS tabulation plan for a detailed description

¹³⁷An ITN is (a) a conventionally treated net which has been soaked with an insecticide within the past 12 months, (b) factory treated net which does not require any treatment (LLIN), (b) a pretreated net obtained within the past 12 months, or (c) a net that has been soaked with or dipped in insecticide within the past 12 months

3.17a 3.17b	Household vector control ¹³⁸	TN - IR	Number of households (a) with at least one ITN or that have been sprayed by IRS ¹³⁹ in the last 12 months (b) with at least one ITN for every two people or that have been sprayed by IRS in the last 12 months	Total number of households	
3.18	Children under age 5 who slept under an ITN	TN	Number of children under age 5 who slept under an ITN the previous night	Total number of children under age 5	MDG 6.7
3.19	Population that slept under an ITN	TN	Number of household members who slept under an ITN the previous night	Total number of household members who spent the previous night in the interviewed households	
3.20	Care-seeking for fever	CA	Number of children under age 5 with fever in the last 2 weeks for whom advice or treatment was sought from a health facility or provider	Total number of children under age 5 with fever in the last 2 weeks	
3.21	Malaria diagnostics usage	CA	Number of children under age 5 with fever in the last 2 weeks who had a finger or heel prick for malaria testing	Total number of children under age 5 with fever in the last 2 weeks	
3.22	Anti-malarial treatment of children under age 5	CA	Number of children under age 5 who tested positive for malaria in the last 2 weeks who received any antimalarial treatment	Total number of children under age 5 who tested positive for malaria in the last 2 weeks	MDG 6.8
3.23	Treatment with Artemisinin-based Combination Therapy (ACT) among children who received malarial treatment	CA	Number of children under age 5 with fever in the last 2 weeks who received ACT or Quinine (or other first-line treatment according to national policy)	Total number of children under age 5 with fever in the last 2 weeks who received any anti-malarial drugs	
3.24	Pregnant women who slept under an ITN	TN - CP	Number of pregnant women who slept under an ITN the previous night	Total number of pregnant women	
3.25	Intermittent preventive treatment for malaria during pregnancy	MN	Number of women age 15-49 years who received two or more doses of SP/Fansidar, at least one of which was received during an ANC visit, to prevent malaria during their last pregnancy that led to a live birth in the last 2 years	Total number of women age 15-49 years who have had a live birth in the last 2 years	

¹³⁸(a) Households covered by vector control, (b) Universal coverage of vector control

¹³⁹Indoor Residual Spraying

WATER AND SANITATION					
4.1	Use of improved drinking water sources	WS	Number of household members using improved sources of drinking water	Total number of household members	MDG 7.8
4.2	Water treatment	WS	Number of household members in households using unimproved drinking water sources who use an appropriate treatment method	Total number of household members in households using unimproved drinking water sources	
4.3	Use of improved sanitation	WS	Number of household members using improved sanitation facilities which are not shared	Total number of household members	MDG 7.9
4.4	Safe disposal of child's faeces	CA	Number of children age 0-2 years whose last stools were disposed of safely	Total number of children age 0-2 years	
4.5	Place for handwashing	HW	Number of households with a specific place for hand washing where water and soap or other cleansing agent are present	Total number of households	
4.6	Availability of soap or other cleansing agent	HW	Number of households with soap or other cleansing agent	Total number of households	

REPRODUCTIVE HEALTH					
5.1	Adolescent birth rate ¹⁴⁰	CM - BH	Age-specific fertility rate for women age 15-19 years		MDG 5.4
5.2	Early childbearing	CM - BH	Number of women age 20-24 years who had at least one live birth before age 18	Total number of women age 20-24 years	
5.3	Contraceptive prevalence rate	CP	Number of women age 15-49 years currently married or in union who are using (or whose partner is using) a (modern or traditional) contraceptive method	Total number of women age 15-49 years who are currently married or in union	MDG 5.3
5.4	Unmet need ¹⁴¹	UN	Number of women age 15-49 years who are currently married or in union who are fecund and want to space their births or limit the number of children they have and who are not currently using contraception	Total number of women age 15-49 years who are currently married or in union	MDG 5.6
5.5a 5.5b	Antenatal care coverage	MN	Number of women age 15-49 years with a live birth in the last 2 years who were attended (a) at least once by skilled personnel (b) at least four times by skilled personnel during their last pregnancy that led to a live birth	Total number of women age 15-49 years with a live birth in the last 2 years	MDG 5.5

¹⁴⁰The indicator is calculated for the last 3-year period.

¹⁴¹See the MICS tabulation plan for a detailed description

5.6	Content of antenatal care	MN	Number of women age 15-49 years with a live birth in the last 2 years who had their blood pressure measured and gave urine and blood samples during the last pregnancy that led to a live birth	Total number of women age 15-49 years with a live birth in the last 2 years	
5.7	Skilled attendant at delivery	MN	Number of women age 15-49 years with a live birth in the last 2 years who were attended by skilled health personnel during their most recent live birth	Total number of women age 15-49 years with a live birth in the last 2 years	MDG 5.2
5.8	Institutional deliveries	MN	Number of women age 15-49 years with a live birth in the last 2 years whose most recent live birth was delivered in a health facility	Total number of women age 15-49 years with a live birth in the last 2 years	
5.9	Caesarean section	MN	Number of women age 15-49 years whose most recent live birth in the last 2 years was delivered by caesarean section	Total number of women age 15-49 years with a live birth in the last 2 years	
5.10	Post-partum stay in health facility	PN	Number of women age 15-49 years who stayed in the health facility for 24 hours or more after the delivery of their most recent live birth in the last 2 years	Total number of women age 15-49 years with a live birth in the last 2 years	
5.11	Post-natal health check for the newborn	PN	Number of last live births in the last 2 years who received a health check while in facility or at home following delivery, or a post-natal care visit within 2 days after delivery	Total number of last live births in the last 2 years	
5.12	Post-natal health check for the mother	PN	Number of women age 15-49 years who received a health check while in facility or at home following delivery, or a post-natal care visit within 2 days after delivery of their most recent live birth in the last 2 years	Total number of women age 15-49 years with a live birth in the last 2 years	
5.13	Maternal mortality ratio	MM	Deaths during pregnancy, childbirth, or within two months after delivery or termination of pregnancy, per 100,000 births within the 5-year period preceding the survey		MDG 5.1

CHILD DEVELOPMENT

6.1	Net Attendance to early childhood education	EC	Number of children age 36-59 months who are attending an early childhood education programme	Total number of children age 36-59 months	
6.2	Support for learning	EC	Number of children age 36-59 months with whom an adult has engaged in four or more activities to promote learning and school readiness in the last 3 days	Total number of children age 36-59 months	
6.3	Father's support for learning	EC	Number of children age 36-59 months whose father has engaged in four or more activities to promote learning and school readiness in the last 3 days	Total number of children age 36-59 months	
6.4	Mother's support for learning	EC	Number of children age 36-59 months whose mother has engaged in four or more activities to promote learning and school readiness in the last 3 days	Total number of children age 36-59 months	

6.5	Availability of children's books	EC	Number of children under age 5 who have three or more children's books	Total number of children under age 5	
6.6	Availability of playthings	EC	Number of children under age 5 with two or more types of playthings	Total number of children under age 5	
6.7	Inadequate care	EC	Number of children under age 5 left alone or in the care of another child younger than 10 years of age for more than one hour at least once in the last week	Total number of children under age 5	
6.8	Early child development index	EC	Number of children age 36-59 months who are developmentally on track in literacy-numeracy, physical, social-emotional, and learning domains	Total number of children age 36-59 months	

LITERACY AND EDUCATION					
7.1	Literacy rate among young women	WB	Number of women age 15-24 years who are able to read a short simple statement about everyday life or who attended secondary or higher education	Total number of women age 15-24 years	MDG 2.3
7.2	School readiness	ED	Number of children in first grade of primary school who attended pre-school during the previous school year	Total number of children attending the first grade of primary school	
7.3	Net intake rate in primary education	ED	Number of children of school-entry age who enter the first grade of primary school	Total number of children of school-entry age	
7.4	Primary school net attendance ratio (adjusted) ¹⁴²	ED	Number of children of primary school age currently attending primary (primary 1-6; ISCED 1) or secondary school	Total number of children of primary school age (ISCED)	MDG 2.1
7.S1	Primary school net attendance ratio (adjusted)	ED	Number of children of primary school age currently attending primary (primary 1-8; national) or secondary school	Total number of children of primary school age (national)	
7.5	Secondary school net attendance ratio (adjusted)	ED	Number children of secondary school age currently attending secondary (primary 7-8 included; ISCED) school or higher	Total number of children of secondary school age (ISCED)	
7.S2	Secondary school net attendance ratio (adjusted)	ED	Number of children of secondary school age currently attending secondary school (national) or higher	Total number of children of secondary school age	
7.6	Children reaching last grade of primary	ED	Proportion of children entering the first grade of primary school who eventually reach last grade (primary 6; ISCED)		MDG 2.2
7.S3	Children reaching last grade of primary	ED	Proportion of children entering the first grade of primary school who eventually reach last grade (primary 8; national)		
7.7	Primary completion rate	ED	Number of children attending the last grade of primary school (excluding repeaters) (ISCED)	Total number of children of primary school completion age (age appropriate to final grade of primary school) (ISCED)	

¹⁴²For Kenya, the International Standard Classification of Education (ISCED) 1997 classifies Primary 7 and 8 as Lower Secondary education. The indicators labelled ISCED calculates Primary School indicators based on Primary 1-6 only, whereas Primary 7 and 8 are included in Secondary School indicators. Those indicators labelled national and marked with S are based on the national education system, which includes Primary 7-8 in Primary School indicators.

7.S4	Primary completion rate	ED	Number of children attending the last grade of primary school (excluding repeaters) (national)	Total number of children of primary school completion age (age appropriate to final grade of primary school) (national)	
7.7a	Secondary completion rate	ED	Number of children attending the last grade of secondary school (form four), excluding repeaters	Total number of children of secondary school (form four) completion age (age appropriate to final grade of secondary school)	
7.8	Transition rate to secondary school	ED	Number of children attending the last grade of primary school during the previous school year who are in the first grade of secondary school during the current school year (ISCED)	Total number of children attending the last grade of primary school during the previous school year (ISCED)	
7.S5	Transition rate to secondary school	ED	Number of children attending the last grade of primary school during the previous school year who are in the first grade of secondary school during the current school year (national)	Total number of children attending the last grade of primary school during the previous school year (national)	
7.9	Gender parity index (primary school)	ED	Primary school net attendance ratio (adjusted) for girls (ISCED)	Primary school net attendance ratio (adjusted) for boys (ISCED)	MDG 3.1
7.S6	Gender parity index (primary school)	ED	Primary school net attendance ratio (adjusted) for girls (national)	Primary school net attendance ratio (adjusted) for boys (national)	
7.10	Gender parity index (secondary school)	ED	Secondary school net attendance ratio (adjusted) for girls (ISCED)	Secondary school net attendance ratio (adjusted) for boys (ISCED)	MDG 3.1
7.S7	Gender parity index (secondary school)	ED	Secondary school net attendance ratio (adjusted) for girls (national)	Secondary school net attendance ratio (adjusted) for boys (national)	

CHILD PROTECTION					
8.1	Birth registration	BR	Number of children under age 5 whose births are reported registered	Total number of children under age 5	
8.2	Child labour	CL	Number of children age 5-17 years who are involved in child labour	Total number of children age 5-17 years	
8.3	Violent discipline	CD	Number of children age 1-14 years who experienced psychological aggression or physical punishment during the last one month	Total number of children age 1-14 years	
8.4	Marriage before age 15	MA	Number of women age 15-49 years who were first married or in union before age 15	Total number of women age 15-49 years	
8.5	Marriage before age 18	MA	Number of women age 20-49 years who were first married or in union before age 18	Total number of women age 20-49 years	
8.6	Young women age 15-19 years currently married or in union	MA	Number of women age 15-19 years who are married or in union	Total number of women age 15-19 years	
8.7	Polygyny	MA	Number of women age 15-49 years who are in a polygynous union	Total number of women age 15-49 years who are married or in union	
8.8a 8.8b	Spousal age difference	MA	Number of women who are married or in union and whose spouse is 10 or more years older, (a) among women age 15-19 years, (b) among women age 20-24 years	Total number of women who are married or in union (a) age 15-19 years, (b) age 20-24 years	
8.9	Approval for female genital mutilation/cutting (FGM/C)	FGM/C	Number of women age 15-49 years who state that FGM/C should be continued	Total number of women age 15-49 years	
8.10	Prevalence of FGM/C among women	FGM/C	Number of women age 15-49 years who report to have undergone any form of FGM/C	Total number of women age 15-49 years	
8.11	Prevalence of FGM/C among girls	FGM/C	Number of daughters age 0-14 years who have undergone any form of FGM/C, as reported by mothers age 15-49 years	Total number of daughters age 0-14 years	
8.12	Attitudes towards domestic violence	DV	Number of women who state that a husband/partner is justified in hitting or beating his wife in at least one of the following circumstances: (1) she goes out without telling him, (2) she neglects the children, (3) she argues with him, (4) she refuses sex with him, (5) she burns the food	Total number of women age 15-49 years	

8.13	Children's living arrangements	HL	Number of children age 0-17 years living with neither biological parent	Total number of children age 0-17 years	
8.14	Prevalence of children with one or both parents dead	HL	Number of children age 0-17 years with one or both parents dead	Total number of children age 0-17 years	
8.15	Children with at least one parent living abroad	HL	Number of children 0-17 years with at least one parent living abroad	Number of children 0-17 years	

HIV/AIDS AND SEXUAL BEHAVIOUR

9.1	Knowledge about HIV prevention among young women	HA	Number of women age 15-24 years who correctly identify ways of preventing the sexual transmission of HIV ¹⁴³ , and who reject major misconceptions about HIV transmission	Total number of women age 15-24 years	MDG 6.3
9.2	Knowledge of mother-to-child transmission of HIV	HA	Number of women age 15-49 years who correctly identify all three means ¹⁴⁴ of mother-to-child transmission of HIV	Total number of women age 15-49 years	
9.3	Accepting attitudes towards people living with HIV	HA	Number of women age 15-49 years expressing accepting attitudes on all four questions ¹⁴⁵ toward people living with HIV	Total number of women age 15-49 years who have heard of HIV	
9.4	Women who know where to be tested for HIV	HA	Number of women age 15-49 years who state knowledge of a place to be tested for HIV	Total number of women age 15-49 years	
9.5	Women who have been tested for HIV and know the results	HA	Number of women age 15-49 years who have been tested for HIV in the last 12 months and who know their results	Total number of women age 15-49 years	
9.6	Sexually active young women who have been tested for HIV and know the results	HA	Number of women age 15-24 years who have had sex in the last 12 months, who have been tested for HIV in the last 12 months and who know their results	Total number of women age 15-24 years who have had sex in the last 12 months	
9.7	HIV counselling during antenatal care	HA	Number of women age 15-49 years who had a live birth in the last 2 years and received antenatal care during the pregnancy of their most recent birth, reporting that they received counselling on HIV during antenatal care	Total number of women age 15-49 years who had a live birth in the last 2 years	

¹⁴³Using condoms and limiting sex to one faithful, uninfected partner

¹⁴⁴Transmission during pregnancy, during delivery, and by breastfeeding

¹⁴⁵Women (1) who think that a female teacher with the AIDS virus should be allowed to teach in school, (2) who would buy fresh vegetables from a shopkeeper or vendor who has the AIDS virus, (3) who would not want to keep it as a secret if a family member became infected with the AIDS virus, and (4) who would be willing to care for a family member who became sick with the AIDS virus

9.8	HIV testing during antenatal care	HA	Number of women age 15-49 years who had a live birth in the last 2 years and received antenatal care during the pregnancy of their most recent birth, reporting that they were offered and accepted an HIV test during antenatal care and received their results	Total number of women age 15-49 years who had a live birth in the last 2 years	
9.9	Young women who have never had sex	SB	Number of never married women age 15-24 years who have never had sex	Total number of never married women age 15-24 years	
9.10	Sex before age 15 among young women	SB	Number of women age 15-24 years who had sexual intercourse before age 15	Total number of women age 15-24 years	
9.11	Age-mixing among sexual partners	SB	Number of women age 15-24 years who had sex in the last 12 months with a partner who was 10 or more years older	Total number of women age 15-24 years who had sex in the last 12 months	
9.12	Multiple sexual partnerships	SB	Number of women age 15-49 years who had sexual intercourse with more than one partner in the last 12 months	Total number of women age 15-49 years	
9.13	Condom use at last sex among people with multiple sexual partnerships	SB	Number of women age 15-49 years who report having had more than one sexual partner in the last 12 months who also reported that a condom was used the last time they had sex	Total number of women age 15-49 years who reported having had more than one sexual partner in the last 12 months	
9.14	Sex with non-regular partners	SB	Number of sexually active women age 15-24 years who had sex with a non-marital, non-cohabitating partner in the last 12 months	Total number of women age 15-24 years who had sex in the last 12 months	
9.15	Condom use with non-regular partners	SB	Number of women age 15-24 years reporting the use of a condom during the last sexual intercourse with a non-marital, non-cohabiting sex partner in the last 12 months	Total number of women age 15-24 years who had a non-marital, non-cohabiting partner in the last 12 months	MDG 6.2
9.15a	Condom use with regular partners	SB	Number of women age 15-24 years reporting the use of a condom during the last sexual intercourse with a marital, cohabiting sex partner in the last 12 months	Total number of women age 15-24 years who had a marital, cohabiting partner in the last 12 months	

ACCESS TO MASS MEDIA AND USE OF INFORMATION/COMMUNICATION TECHNOLOGY					
10.1	Exposure to mass media	MT	Number of women age 15-49 years who, at least once a week, read a newspaper or magazine, listen to the radio, and watch television	Total number of women age 15-49 years	
10.2	Use of computers	MT	Number of young women age 15-24 years who used a computer during the last 12 months	Total number of women age 15-24 years	
10.3	Use of internet	MT	Number of young women age 15-24 who used the internet during the last 12 months	Total number of women age 15-24 years	

SUBJECTIVE WELL-BEING					
11.1	Life satisfaction		Number of young women age 15-24 years who are very or somewhat satisfied with their life, overall	Total number of young women age 15-24 years	
11.2	Happiness		Number of young women age 15-24 years who are very or somewhat happy	Total number of young women age 15-24 years	
11.3	Perception of a better life		Number of young women age 15-24 years whose life improved during the last one year, and who expect that their life will be better after one year	Total number of young women age 15-24 years	

TOBACCO AND ALCOHOL USE					
12.1	Tobacco use	TA	Number of women age 15-49 years who smoked cigarettes, or used smoked or smokeless tobacco products at any time during the last one month	Total number of women age 15-49 years	
12.2	Smoking before age 15	TA	Number of women age 15-49 years who smoked a whole cigarette before age 15	Total number of women age 15-49 years	
12.3	Use of alcohol	TA	Number of women age 15-49 years who had at least one alcoholic drink at any time during the last one month	Total number of women age 15-49 years	
12.4	Use of alcohol before age 15	TA	Number of women age 15-49 years who had at least one alcoholic drink before age 15	Total number of women age 15-49 years	

Appendix H. Turkana County MICS Questionnaires

HOUSEHOLD QUESTIONNAIRE WESTERN AND NORTH RIFT SURVEY



HOUSEHOLD INFORMATION PANEL		HH
HH1. Cluster number: _____	HH2. Household number: _____	
HH3. Interviewer's name and number: Name _____	HH4. Supervisor's name and number: Name _____	
HH5. Day / Month / Year of interview: _____ / _____ / 201 _____	HH7. Region: Bungoma..... 1 Kakamega..... 2 Turkana..... 3	
HH6. Area: Urban.....1 Rural.....2		
<p>WE ARE FROM UNIVERSITY OF NAIROBI AND KENYA NATIONAL BUREAU OF STATISTICS. WE ARE CONDUCTING A SURVEY ABOUT THE SITUATION OF CHILDREN, FAMILIES AND HOUSEHOLDS. I WOULD LIKE TO TALK TO YOU ABOUT THESE SUBJECTS. THE INTERVIEW WILL TAKE ABOUT 55 MINUTES TO ONE HOUR. ALL THE INFORMATION WE OBTAIN WILL REMAIN STRICTLY CONFIDENTIAL AND ANONYMOUS. MAY I START NOW?</p> <p><input type="checkbox"/> Yes, permission is given ⇒ Go to HH18 to record the time and then begin the interview.</p> <p><input type="checkbox"/> No, permission is not given ⇒ Circle 04 in HH9. Discuss this result with your supervisor.</p>		
<p>HH9. Result of household interview:</p> <p>Completed.....01 No household member or no competent respondent at home at time of visit02 Entire household absent for extended period of time03 Refused04 Dwelling vacant / Address not a dwelling.....05 Dwelling destroyed.....06 Dwelling not found.....07</p> <p>Other (specify) _____ 96</p>		

After the household questionnaire has been completed, fill in the following information:

HH10. Respondent to Household Questionnaire:

Name _____

HH11. Total number of household members: _____

HH12. Number of women age 15-49 years: _____

HH14. Number of children under age 5: _____

After all questionnaires for the household have been completed, fill in the following information:

HH13. Number of women's questionnaires completed: _____

HH15. Number of under-5 questionnaires completed: _____

HH16. Field editor's name and number:

Name _____

HH17. Main data entry clerk's name and number:

Name _____

HH18. Record the time.

Hour — —

Minutes..... — —

LIST OF HOUSEHOLD MEMBERS

HL

FIRST, PLEASE TELL ME THE NAME OF EACH PERSON WHO USUALLY LIVES HERE, STARTING WITH THE HEAD OF THE HOUSEHOLD.

List the head of the household in line 01. List all household members (HL2), their relationship to the household head (HL3), and their sex (HL4)

Then ask: ARE THERE ANY OTHERS WHO LIVE HERE, EVEN IF THEY ARE NOT AT HOME NOW?

If yes, complete listing for questions HL2-HL4. Then, ask questions starting with HL5 for each person at a time.

Use an additional questionnaire if all rows in the List of Household Members have been used.

							For women age 15-49	For children age 0-4	For children age 0-17 years							For children age 0-14									
HL1. Line no.	HL2. Name	HL3. WHAT IS THE RELATIONSHIP OF (name) TO THE HEAD OF HOUSEHOLD?	HL4. IS (name) MALE OR FEMALE? 1 Male 2 Female	HL5. WHAT IS (name)'S DATE OF BIRTH?		HL6. HOW OLD IS (name)? <i>Record in complete d years. If age is 95 or above, record '95'</i>	HL6A. DID (name) STAY HERE LAST NIGHT? 1 Yes 2 No	HL7. <i>Circle line no. if woman age 15-49</i>	HL7B. <i>Circle line no. if age 0-4</i>	HL11. IS (name)'S NATURAL MOTHER ALIVE? 1 Yes 2 No 8 DK HL13	HL12. DOES (name)'S NATURAL MOTHER LIVE IN THIS HOUSEHOLD? <i>If "Yes" Record line no. of mother and go to HL13</i> Record 00 for "No"	HL12A. WHERE DOES (name)'S NATURAL MOTHER LIVE? 1 In another household in this country 2 Institution in this country 3 Abroad 8 DK	HL13. IS (name)'S NATURAL FATHER ALIVE? 1 Yes 2 No 8 DK HL15	HL14. DOES (name)'S NATURAL FATHER LIVE IN THIS HOUSEHOLD? <i>If "Yes" Record line no. of father and go to HL15</i> Record 00 for "No"	HL14A. WHERE DOES (name)'S NATURAL FATHER LIVE? 1 In another household in this country 2 Institution in this country 3 Abroad 8 DK	HL15. <i>Record line no. of mother from HL12 if indicated. If HL12 is blank, or "00" ask:</i> WHO IS THE PRIMARY CARETAKER OF (name)?									
Line	Name	Relation*	M	F	Month	Year	Age	Y	N	15-49	0-4	Y	N	DK	Mother	Y	N	DK	Father	Mother					
01		01	1	2	___	___	___	1	2	01	01	1	2	8	___	1	2	8	___	1	2	3	8	___	___
02		___	1	2	___	___	___	1	2	02	02	1	2	8	___	1	2	8	___	1	2	3	8	___	___
03		___	1	2	___	___	___	1	2	03	03	1	2	8	___	1	2	8	___	1	2	3	8	___	___
04		___	1	2	___	___	___	1	2	04	04	1	2	8	___	1	2	8	___	1	2	3	8	___	___
05		___	1	2	___	___	___	1	2	05	05	1	2	8	___	1	2	8	___	1	2	3	8	___	___
06		___	1	2	___	___	___	1	2	06	06	1	2	8	___	1	2	8	___	1	2	3	8	___	___
07		___	1	2	___	___	___	1	2	07	07	1	2	8	___	1	2	8	___	1	2	3	8	___	___
08		___	1	2	___	___	___	1	2	08	08	1	2	8	___	1	2	8	___	1	2	3	8	___	___
09		___	1	2	___	___	___	1	2	09	09	1	2	8	___	1	2	8	___	1	2	3	8	___	___
10		___	1	2	___	___	___	1	2	10	10	1	2	8	___	1	2	8	___	1	2	3	8	___	___

								For women age 15-49	For children age 0-4	For children age 0-17 years						For children age 0-14							
HL1. Line no.	HL2. Name	HL3. WHAT IS THE RELATIONSHIP OF (name) TO THE HEAD OF HOUSEHOLD?	HL4. IS (name) MALE OR FEMALE? 1 Male 2 Female	HL5. WHAT IS (name)'S DATE OF BIRTH?		HL6. HOW OLD IS (name)? <i>Record in complete d years. If age is 95 or above, record '95'</i>	HL6A. DID (name) STAY HERE LAST NIGHT? 1 Yes 2 No	HL7. <i>Circle line no. if woman age 15-49</i>	HL7B. <i>Circle line no. if age 0-4</i>	HL11. IS (name)'S NATURAL MOTHER ALIVE? 1 Yes 2 No ^{HL13} 8 DK ^{HL13}	HL12. DOES (name)'S NATURAL MOTHER LIVE IN THIS HOUSEHOLD? <i>If "Yes" Record line no. of mother and go to HL13</i> Record 00 for "No"	HL12A. WHERE DOES (name)'S NATURAL MOTHER LIVE? 1 In another household in this country 2 Institution in this country 3 Abroad 8 DK	HL13. IS (name)'S NATURAL FATHER ALIVE? 1 Yes 2 No ^{HL15} 8 DK ^{HL15}	HL14. DOES (name)'S NATURAL FATHER LIVE IN THIS HOUSEHOLD? <i>If "Yes" Record line no. of father and go to HL15</i> Record 00 for "No"	HL14A. WHERE DOES (name)'S NATURAL FATHER LIVE? 1 In another household in this country 2 Institution in this country 3 Abroad 8 DK	HL15. <i>Record line no. of mother from HL12 if indicated. If HL12 is blank, or "00" ask:</i> WHO IS THE PRIMARY CARETAKER OF (name)?							
Line	Name	Relation*	M	F	Month	Year	Age	Y	N	15-49	0-4	Y	N	DK	Mother	Y	N	DK	Father	Y	N	DK	Mother
11		___ ___	1	2	___	___	___	1	2	11	11	1	2	8	___	1	2	8	___	1	2	8	___
12		___ ___	1	2	___	___	___	1	2	12	12	1	2	8	___	1	2	8	___	1	2	8	___
13		___ ___	1	2	___	___	___	1	2	13	13	1	2	8	___	1	2	8	___	1	2	8	___
14		___ ___	1	2	___	___	___	1	2	14	14	1	2	8	___	1	2	8	___	1	2	8	___
15		___ ___	1	2	___	___	___	1	2	15	15	1	2	8	___	1	2	8	___	1	2	8	___

Tick here if additional questionnaire used

Probe for additional household members.
 Probe especially for any infants or small children not listed, and others who may not be members of the family (such as servants, friends) but who usually live in the household.
 Insert names of additional members in the household list and complete form accordingly.

Now for each woman age 15-49 years, write her name and line number and other identifying information in the information panel of a separate Individual Women's Questionnaire.
 For each man age 15-49 years, write his name and line number and other identifying information in the information panel of a separate Individual Man's Questionnaire.
 For each child under age 5, write his/her name and line number AND the line number of his/her mother or caretaker in the information panel of a separate Under-5 Questionnaire.
 You should now have a separate questionnaire for each eligible woman, each eligible man, and each child under five in the household.

* Codes for HL3: Relationship to head of household:	01 Head	04 Son-In-Law / Daughter-In-Law	07 Parent-In-Law	10 Uncle / Aunt	13 Adopted / Foster/ Stepchild	96 Other (Not related)
	02 Spouse/Partner	05 Grandchild	08 Brother / Sister	11 Niece / Nephew	14 Servant (Live-in)	98 DK
	03 Son / Daughter	06 Parent	09 Brother-In-Law / Sister-In-Law	12 Other relative		

EDUCATION			ED													
			For household members age 5 and above				For household members age 5-24 years									
ED1. Line number	ED2. Name and age Copy from HL2 and HL6		ED3.		ED4A.		ED4B.		ED5.		ED6.		ED7.		ED8.	
			HAS (name) EVER ATTENDED SCHOOL OR PRE- SCHOOL?		WHAT IS THE HIGHEST LEVEL OF SCHOOL (name) HAS ATTENDED? Level: 0 Preschool 1 Primary 2 Secondary 3 Higher 8 DK If level=0, skip to ED5		WHAT IS THE HIGHEST GRADE (name) COMPLETED AT THIS LEVEL? Grade: 98 DK If grade 1 is not completed at this level, enter "00"		DURING THE CURRENT SCHOOL YEAR THAT IS 2013 - 2014, DID (name) ATTEND SCHOOL OR PRESCHOOL AT ANY TIME?		DURING THIS/THAT SCHOOL YEAR, WHICH LEVEL AND GRADE IS/WAS (name) ATTENDING? Level: 0 Preschool 1 Primary 2 Secondary 3 Higher 8 DK If level=0, skip to ED7		DURING THE PREVIOUS SCHOOL YEAR, THAT IS 2012- 2013, DID (name) ATTEND SCHOOL OR PRESCHOOL AT ANY TIME?		DURING THAT PREVIOUS SCHOOL YEAR, WHICH LEVEL AND GRADE DID (name) ATTEND? Level: 0 Preschool 1 Primary 2 Secondary 3 Higher 8 DK If level=0, go to next line'	
Line	Name	Age	Yes	No	Level	Grade	Yes	No	Level	Grade	Yes	No	DK	Level	Grade	
01		_____	1	2	0 1 2 3 8	___ __	1	2	0 1 2 3 8	___ __	1	2	8	0 1 2 3 8	___ __	
02		_____	1	2	0 1 2 3 8	___ __	1	2	0 1 2 3 8	___ __	1	2	8	0 1 2 3 8	___ __	
03		_____	1	2	0 1 2 3 8	___ __	1	2	0 1 2 3 8	___ __	1	2	8	0 1 2 3 8	___ __	
04		_____	1	2	0 1 2 3 8	___ __	1	2	0 1 2 3 8	___ __	1	2	8	0 1 2 3 8	___ __	
05		_____	1	2	0 1 2 3 8	___ __	1	2	0 1 2 3 8	___ __	1	2	8	0 1 2 3 8	___ __	
06		_____	1	2	0 1 2 3 8	___ __	1	2	0 1 2 3 8	___ __	1	2	8	0 1 2 3 8	___ __	
07		_____	1	2	0 1 2 3 8	___ __	1	2	0 1 2 3 8	___ __	1	2	8	0 1 2 3 8	___ __	
08		_____	1	2	0 1 2 3 8	___ __	1	2	0 1 2 3 8	___ __	1	2	8	0 1 2 3 8	___ __	
09		_____	1	2	0 1 2 3 8	___ __	1	2	0 1 2 3 8	___ __	1	2	8	0 1 2 3 8	___ __	
10		_____	1	2	0 1 2 3 8	___ __	1	2	0 1 2 3 8	___ __	1	2	8	0 1 2 3 8	___ __	
11		_____	1	2	0 1 2 3 8	___ __	1	2	0 1 2 3 8	___ __	1	2	8	0 1 2 3 8	___ __	
12		_____	1	2	0 1 2 3 8	___ __	1	2	0 1 2 3 8	___ __	1	2	8	0 1 2 3 8	___ __	
13		_____	1	2	0 1 2 3 8	___ __	1	2	0 1 2 3 8	___ __	1	2	8	0 1 2 3 8	___ __	
14		_____	1	2	0 1 2 3 8	___ __	1	2	0 1 2 3 8	___ __	1	2	8	0 1 2 3 8	___ __	
15		_____	1	2	0 1 2 3 8	___ __	1	2	0 1 2 3 8	___ __	1	2	8	0 1 2 3 8	___ __	

SELECTION OF ONE CHILD FOR CHILD LABOUR/CHILD DISCIPLINE					SL		
SL1. Check HL6 in the List of Household Members and write the total number of children age 1-17 years.	Total number						
SL2. Check the number of children age 1-17 years in SL1:							
<input type="checkbox"/> Zero ⇒ Go to HOUSEHOLD CHARACTERISTICS module							
<input type="checkbox"/> One ⇒ Go to SL9 and record the rank number as '1', enter the line number, child's name and age							
<input type="checkbox"/> Two or more ⇒ Continue with SL2A							
SL2A. List each of the children age 1-17 years below in the order they appear in the List of Household Members. Do not include other household members outside of the age range 1-17 years. Record the line number, name, sex, and age for each child.							
SL3. Rank number	SL4. Line number from HL1	SL5. Name from HL2	SL6. Sex from HL4		SL7. Age from HL6		
Rank	Line	Name	M	F	Age		
1	__ __		1	2	__ __		
2	__ __		1	2	__ __		
3	__ __		1	2	__ __		
4	__ __		1	2	__ __		
5	__ __		1	2	__ __		
6	__ __		1	2	__ __		
7	__ __		1	2	__ __		
8	__ __		1	2	__ __		
SL8. Check the last digit of the household number (HH2) from the cover page. This is the number of the row you should go to in the table below.							
Check the total number of children age 1-17 years in SL1 above. This is the number of the column you should go to in the table below							
Find the box where the row and the column meet and circle the number that appears in the box. This is the rank number (SL3) of the selected child.							
Last Digit of Household Number (from HH2)	Total Number of Eligible Children in the Household (from SL1)						
	2	3	4	5	6	7	8+
0	2	2	4	3	6	5	4
1	1	3	1	4	1	6	5
2	2	1	2	5	2	7	6
3	1	2	3	1	3	1	7
4	2	3	4	2	4	2	8
5	1	1	1	3	5	3	1
6	2	2	2	4	6	4	2
7	1	3	3	5	1	5	3
8	2	1	4	1	2	6	4
9	1	2	1	2	3	7	5
SL9. Record the rank number (SL3), line number (SL4), name (SL5) and age (SL7) of the selected child			Rank number			Line number	
			Name			Age	

CHILD LABOUR		CL															
CL1. Check selected child's age from SL9: <input type="checkbox"/> 1-4 years ⇒ Go to Next Module <input type="checkbox"/> 5-17 years ⇒ Continue with CL2																	
CL2. NOW I WOULD LIKE TO ASK ABOUT ANY WORK CHILDREN IN THIS HOUSEHOLD MAY DO. SINCE LAST (<i>day of the week</i>), DID (<i>name</i>) DO ANY OF THE FOLLOWING ACTIVITIES, EVEN FOR ONLY ONE HOUR? [A] DID (<i>name</i>) DO ANY WORK OR HELP ON HIS/HER OWN OR THE HOUSEHOLD'S PLOT/FARM/FOOD GARDEN OR LOOKED AFTER ANIMALS? FOR EXAMPLE, GROWING FARM PRODUCE, HARVESTING, OR FEEDING, GRAZING, MILKING ANIMALS? [B] DID (<i>name</i>) HELP IN FAMILY BUSINESS OR RELATIVE'S BUSINESS WITH OR WITHOUT PAY, OR RUN HIS/HER OWN BUSINESS? [C] DID (<i>name</i>) PRODUCE OR SELL ARTICLES, HANDICRAFTS, CLOTHES, FOOD OR AGRICULTURAL PRODUCTS? [D] SINCE LAST (<i>day of the week</i>), DID (<i>name</i>) ENGAGE IN ANY OTHER ACTIVITY IN RETURN FOR INCOME IN CASH OR IN KIND, EVEN FOR ONLY ONE HOUR? <i>If "No", Probe:</i> PLEASE INCLUDE ANY ACTIVITY (<i>name</i>) PERFORMED AS A REGULAR OR CASUAL EMPLOYEE, SELF-EMPLOYED OR EMPLOYER; OR AS AN UNPAID FAMILY WORKER HELPING OUT IN HOUSEHOLD BUSINESS OR FARM.	<table border="0"> <thead> <tr> <th></th> <th>Yes</th> <th>No</th> </tr> </thead> <tbody> <tr> <td>Worked on plot/farm/food garden/looked after animals</td> <td>1</td> <td>2</td> </tr> <tr> <td>Helped in family/relative's business/ran own business</td> <td>1</td> <td>2</td> </tr> <tr> <td>Produce/sell articles/handicrafts/clothes/food or agricultural products</td> <td>1</td> <td>2</td> </tr> <tr> <td>Any other activity</td> <td>1</td> <td>2</td> </tr> </tbody> </table>		Yes	No	Worked on plot/farm/food garden/looked after animals	1	2	Helped in family/relative's business/ran own business	1	2	Produce/sell articles/handicrafts/clothes/food or agricultural products	1	2	Any other activity	1	2	
	Yes	No															
Worked on plot/farm/food garden/looked after animals	1	2															
Helped in family/relative's business/ran own business	1	2															
Produce/sell articles/handicrafts/clothes/food or agricultural products	1	2															
Any other activity	1	2															
CL3. Check CL2, A to D <input type="checkbox"/> There is at least one 'Yes' ⇒ continue with CL4 <input type="checkbox"/> All answers are 'No' ⇒ Go to CL8																	
CL4. SINCE LAST (<i>day of the week</i>) ABOUT HOW MANY HOURS DID (<i>name</i>) ENGAGE IN THIS ACTIVITY/THESE ACTIVITIES, IN TOTAL? <i>'if less than one hour, record "00"</i>	Number of hours																
CL5. DOES THE ACTIVITY/DO THESE ACTIVITIES REQUIRE CARRYING HEAVY LOADS?	Yes1 No2	1 ⇒ CL8															
CL6. DOES THE ACTIVITY/DO THESE ACTIVITIES REQUIRE WORKING WITH DANGEROUS TOOLS (KNIVES ETC.) OR OPERATING HEAVY MACHINERY?	Yes1 No2	1 ⇒ CL8															

<p>CL7. HOW WOULD YOU DESCRIBE THE WORK ENVIRONMENT OF <i>(name)</i>?</p> <p>[A] IS <i>(name)</i> EXPOSED TO DUST, FUMES OR GAS?</p> <p>[B] IS <i>(name)</i> EXPOSED TO EXTREME COLD, HEAT OR HUMIDITY?</p> <p>[C] IS <i>(name)</i> EXPOSED TO LOUD NOISE OR VIBRATION?</p> <p>[D] IS <i>(name)</i> REQUIRED TO WORK AT HEIGHTS?</p> <p>[E] IS <i>(name)</i> REQUIRED TO WORK WITH CHEMICALS (PESTICIDES, GLUES, ETC.) OR EXPLOSIVES?</p> <p>[F] IS <i>(name)</i> EXPOSED TO OTHER THINGS, PROCESSES OR CONDITIONS BAD FOR <i>(name)</i>'S HEALTH OR SAFETY?</p>	<p>Yes 1 No 2</p> <p>Yes 1 No 2</p> <p>Yes 1 No 2</p> <p>Yes 1 No 2</p> <p>Yes 1 No 2</p> <p>Yes 1 No 2</p>	<p>1⇒ CL8</p> <p>1⇒ CL8</p> <p>1⇒ CL8</p> <p>1⇒ CL8</p> <p>1⇒ CL8</p>																								
<p>CL8. SINCE LAST <i>(day of the week)</i>, DID <i>(name)</i> FETCH WATER OR COLLECT FIREWOOD FOR HOUSEHOLD USE?</p>	<p>Yes 1 No 2</p>	<p>2⇒ CL10</p>																								
<p>CL9. IN TOTAL, HOW MANY HOURS DID <i>(name)</i> SPEND ON FETCHING WATER OR COLLECTING FIREWOOD FOR HOUSEHOLD USE, SINCE LAST <i>(day of the week)</i>?</p> <p><i>If less than one hour, record "00"</i></p>	<p>Number of hours _ _</p>																									
<p>CL10. SINCE LAST <i>(day of the week)</i>, DID <i>(name)</i> DO ANY OF THE FOLLOWING FOR THIS HOUSEHOLD?</p> <p>[A] SHOPPING FOR HOUSEHOLD?</p> <p>[B] REPAIR ANY HOUSEHOLD EQUIPMENT?</p> <p>[C] COOKING OR CLEANING UTENSILS OR THE HOUSE?</p> <p>[D] WASHING CLOTHES?</p> <p>[E] CARING FOR CHILDREN?</p> <p>[F] CARING FOR THE OLD OR SICK?</p> <p>[G] OTHER HOUSEHOLD TASKS?</p>	<table border="0"> <thead> <tr> <th></th> <th>Yes</th> <th>No</th> </tr> </thead> <tbody> <tr> <td>Shopping for household</td> <td>1</td> <td>2</td> </tr> <tr> <td>Repair household equipment</td> <td>1</td> <td>2</td> </tr> <tr> <td>Cooking/cleaning utensils/house</td> <td>1</td> <td>2</td> </tr> <tr> <td>Washing clothes</td> <td>1</td> <td>2</td> </tr> <tr> <td>Caring for children</td> <td>1</td> <td>2</td> </tr> <tr> <td>Caring for old/sick</td> <td>1</td> <td>2</td> </tr> <tr> <td>Other household tasks</td> <td>1</td> <td>2</td> </tr> </tbody> </table>		Yes	No	Shopping for household	1	2	Repair household equipment	1	2	Cooking/cleaning utensils/house	1	2	Washing clothes	1	2	Caring for children	1	2	Caring for old/sick	1	2	Other household tasks	1	2	
	Yes	No																								
Shopping for household	1	2																								
Repair household equipment	1	2																								
Cooking/cleaning utensils/house	1	2																								
Washing clothes	1	2																								
Caring for children	1	2																								
Caring for old/sick	1	2																								
Other household tasks	1	2																								
<p>CL11. Check CL10, A to G</p> <p><input type="checkbox"/> <i>There is at least one 'Yes' ⇒ Continue with CL12</i></p> <p><input type="checkbox"/> <i>All answers are 'No' ⇒ Go to Next Module</i></p>																										
<p>CL12. SINCE LAST <i>(day of the week)</i>, ABOUT HOW MANY HOURS DID <i>(name)</i> ENGAGE IN THIS ACTIVITY/THESE ACTIVITIES, IN TOTAL?</p> <p><i>If less than one hour, record "00"</i></p>	<p>Number of hours _ _</p>																									

CHILD DISCIPLINE		CD																																				
CD1. Check selected child's age from SL9: <input type="checkbox"/> 1-14 years ⇒ Continue with CD2 <input type="checkbox"/> 15-17 years ⇒ Go to Next Module																																						
CD2. Write the line number and name of the child from SL9.	Line number ____ Name																																					
CD3. ADULTS USE CERTAIN WAYS TO TEACH CHILDREN THE RIGHT BEHAVIOUR OR TO ADDRESS A BEHAVIOUR PROBLEM. I WILL READ VARIOUS METHODS THAT ARE USED. PLEASE TELL ME IF <u>YOU OR ANYONE ELSE IN YOUR HOUSEHOLD</u> HAS USED THIS METHOD WITH <u>(name)</u> IN THE PAST MONTH.	<table style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 80%;"></th> <th style="width: 10%; text-align: center;">Yes</th> <th style="width: 10%; text-align: center;">No</th> </tr> </thead> <tbody> <tr> <td>[A] TOOK AWAY PRIVILEGES, FORBADE SOMETHING (name) LIKED OR DID NOT ALLOW HIM/HER TO LEAVE THE HOUSE.</td> <td style="text-align: center;">1</td> <td style="text-align: center;">2</td> </tr> <tr> <td>[B] EXPLAINED WHY (name)'S BEHAVIOUR WAS WRONG.</td> <td style="text-align: center;">1</td> <td style="text-align: center;">2</td> </tr> <tr> <td>[C] SHOOK HIM/HER.</td> <td style="text-align: center;">1</td> <td style="text-align: center;">2</td> </tr> <tr> <td>[D] SHOUTED, YELLED AT OR SCREAMED AT HIM/HER.</td> <td style="text-align: center;">1</td> <td style="text-align: center;">2</td> </tr> <tr> <td>[E] GAVE HIM/HER SOMETHING ELSE TO DO.</td> <td style="text-align: center;">1</td> <td style="text-align: center;">2</td> </tr> <tr> <td>[F] SPANKED, HIT OR SLAPPED HIM/HER ON THE BOTTOM WITH BARE HAND.</td> <td style="text-align: center;">1</td> <td style="text-align: center;">2</td> </tr> <tr> <td>[G] HIT HIM/HER ON THE BOTTOM OR ELSEWHERE ON THE BODY WITH SOMETHING LIKE A BELT, HAIRBRUSH, STICK OR OTHER HARD OBJECT.</td> <td style="text-align: center;">1</td> <td style="text-align: center;">2</td> </tr> <tr> <td>[H] CALLED HIM/HER DUMB, LAZY, OR ANOTHER NAME LIKE THAT.</td> <td style="text-align: center;">1</td> <td style="text-align: center;">2</td> </tr> <tr> <td>[I] HIT OR SLAPPED HIM/HER ON THE FACE, HEAD OR EARS.</td> <td style="text-align: center;">1</td> <td style="text-align: center;">2</td> </tr> <tr> <td>[J] HIT OR SLAPPED HIM/HER ON THE HAND, ARM, OR LEG.</td> <td style="text-align: center;">1</td> <td style="text-align: center;">2</td> </tr> <tr> <td>[K] BEAT HIM/HER UP, THAT IS HIT HIM/HER OVER AND OVER AS HARD AS ONE COULD.</td> <td style="text-align: center;">1</td> <td style="text-align: center;">2</td> </tr> </tbody> </table>		Yes	No	[A] TOOK AWAY PRIVILEGES, FORBADE SOMETHING (name) LIKED OR DID NOT ALLOW HIM/HER TO LEAVE THE HOUSE.	1	2	[B] EXPLAINED WHY (name)'S BEHAVIOUR WAS WRONG.	1	2	[C] SHOOK HIM/HER.	1	2	[D] SHOUTED, YELLED AT OR SCREAMED AT HIM/HER.	1	2	[E] GAVE HIM/HER SOMETHING ELSE TO DO.	1	2	[F] SPANKED, HIT OR SLAPPED HIM/HER ON THE BOTTOM WITH BARE HAND.	1	2	[G] HIT HIM/HER ON THE BOTTOM OR ELSEWHERE ON THE BODY WITH SOMETHING LIKE A BELT, HAIRBRUSH, STICK OR OTHER HARD OBJECT.	1	2	[H] CALLED HIM/HER DUMB, LAZY, OR ANOTHER NAME LIKE THAT.	1	2	[I] HIT OR SLAPPED HIM/HER ON THE FACE, HEAD OR EARS.	1	2	[J] HIT OR SLAPPED HIM/HER ON THE HAND, ARM, OR LEG.	1	2	[K] BEAT HIM/HER UP, THAT IS HIT HIM/HER OVER AND OVER AS HARD AS ONE COULD.	1	2	
	Yes	No																																				
[A] TOOK AWAY PRIVILEGES, FORBADE SOMETHING (name) LIKED OR DID NOT ALLOW HIM/HER TO LEAVE THE HOUSE.	1	2																																				
[B] EXPLAINED WHY (name)'S BEHAVIOUR WAS WRONG.	1	2																																				
[C] SHOOK HIM/HER.	1	2																																				
[D] SHOUTED, YELLED AT OR SCREAMED AT HIM/HER.	1	2																																				
[E] GAVE HIM/HER SOMETHING ELSE TO DO.	1	2																																				
[F] SPANKED, HIT OR SLAPPED HIM/HER ON THE BOTTOM WITH BARE HAND.	1	2																																				
[G] HIT HIM/HER ON THE BOTTOM OR ELSEWHERE ON THE BODY WITH SOMETHING LIKE A BELT, HAIRBRUSH, STICK OR OTHER HARD OBJECT.	1	2																																				
[H] CALLED HIM/HER DUMB, LAZY, OR ANOTHER NAME LIKE THAT.	1	2																																				
[I] HIT OR SLAPPED HIM/HER ON THE FACE, HEAD OR EARS.	1	2																																				
[J] HIT OR SLAPPED HIM/HER ON THE HAND, ARM, OR LEG.	1	2																																				
[K] BEAT HIM/HER UP, THAT IS HIT HIM/HER OVER AND OVER AS HARD AS ONE COULD.	1	2																																				
CD4. DO YOU BELIEVE THAT IN ORDER TO BRING UP, RAISE, OR EDUCATE A CHILD PROPERLY, THE CHILD NEEDS TO BE PHYSICALLY PUNISHED?	Yes1 No.....2 DK / No opinion8																																					

HOUSEHOLD CHARACTERISTICS		HC
HC1A. WHAT IS THE RELIGION OF THE HEAD OF THIS HOUSEHOLD?	<i>Catholic</i> 1 <i>Other Christian</i> 2 <i>Muslim</i> 3 <i>Traditional</i> 4 Other religion (<i>specify</i>) _____ 6 No religion 7	
HC1B. WHAT IS THE MOTHER TONGUE/NATIVE LANGUAGE OF THE HEAD OF THIS HOUSEHOLD?	<i>Luhya</i> 1 <i>Turkana</i> 2 <i>Swahili</i> 3 Other language (<i>specify</i>) _____ 6	
HC1C. TO WHAT ETHNIC GROUP DOES THE HEAD OF THIS HOUSEHOLD BELONG?	<i>Luhya</i> 1 <i>Turkana</i> 2 Other ethnic group (<i>specify</i>) _____ 6	
HC2. HOW MANY ROOMS IN THIS HOUSEHOLD ARE USED FOR SLEEPING?	Number of rooms..... _ _	
HC3. <i>Main material of the dwelling floor.</i> <i>Record observation.</i>	Natural floor Earth / Sand 11 Dung 12 Rudimentary floor Wood planks 21 Palm / Bamboo 22 Finished floor Parquet or polished wood 31 Vinyl or asphalt strips 32 Ceramic tiles 33 Cement 34 Carpet 35 Other (<i>specify</i>) _____ 96	
HC4. <i>Main material of the roof.</i> <i>Record observation.</i>	Natural roofing No Roof..... 11 Thatch / Palm leaf..... 12 Sod 13 Rudimentary roofing Rustic mat..... 21 Palm / Bamboo 22 Wood planks 23 Cardboard..... 24 Finished roofing Metal/Tin 31 Wood 32 Calamine / Cement fibre..... 33 Ceramic tiles 34 Cement 35 Roofing shingles 36 Other (<i>specify</i>) _____ 96	

<p>HC5. Main material of the exterior walls.</p> <p><i>Record observation.</i></p>	<p>Natural walls</p> <p>No walls11</p> <p>Cane / Palm / Trunks12</p> <p>Dirt13</p> <p>Rudimentary walls</p> <p>Bamboo with mud21</p> <p>Stone with mud22</p> <p>Uncovered adobe23</p> <p>Plywood24</p> <p>Cardboard25</p> <p>Reused wood26</p> <p>Finished walls</p> <p>Cement31</p> <p>Stone with lime / cement32</p> <p>Bricks33</p> <p>Cement blocks34</p> <p>Covered adobe35</p> <p>Wood planks / shingles36</p> <p>Other (<i>specify</i>) 96</p>																												
<p>HC6. WHAT TYPE OF FUEL DOES YOUR HOUSEHOLD MAINLY USE FOR COOKING?</p>	<p>Electricity01</p> <p>Liquefied Petroleum Gas (LPG)02</p> <p>Natural gas03</p> <p>Biogas04</p> <p>Kerosene05</p> <p>Coal / Lignite06</p> <p>Charcoal07</p> <p>Wood08</p> <p>Straw / Shrubs / Grass09</p> <p>Animal dung10</p> <p>Agricultural crop residue11</p> <p>No food cooked in household95</p> <p>Other (<i>specify</i>) 96</p>	<p>01⇒HC8</p> <p>02⇒HC8</p> <p>03⇒HC8</p> <p>04⇒HC8</p> <p>05⇒HC8</p> <p>95⇒HC8</p>																											
<p>HC7. IS THE COOKING USUALLY DONE IN THE HOUSE, IN A SEPARATE BUILDING, OR OUTDOORS?</p> <p><i>If 'In the house', probe: IS IT DONE IN A SEPARATE ROOM USED AS A KITCHEN?</i></p>	<p>In the house</p> <p>In a separate room used as kitchen1</p> <p>Elsewhere in the house2</p> <p>In a separate building3</p> <p>Outdoors4</p> <p>Other (<i>specify</i>) 6</p>																												
<p>HC8. DOES YOUR HOUSEHOLD HAVE:</p> <p>[A] ELECTRICITY?</p> <p>[B] A RADIO?</p> <p>[C] A TELEVISION?</p> <p>[D] A NON-MOBILE TELEPHONE?</p> <p>[E] A REFRIGERATOR?</p> <p>[F] SOLAR PANEL</p> <p>[G] CHAIR</p> <p>[H] SOFA SET</p>	<table border="1"> <thead> <tr> <th></th> <th>Yes</th> <th>No</th> </tr> </thead> <tbody> <tr> <td>Electricity</td> <td>1</td> <td>2</td> </tr> <tr> <td>Radio</td> <td>1</td> <td>2</td> </tr> <tr> <td>Television</td> <td>1</td> <td>2</td> </tr> <tr> <td>Non-mobile telephone</td> <td>1</td> <td>2</td> </tr> <tr> <td>Refrigerator.....</td> <td>1</td> <td>2</td> </tr> <tr> <td>Solar Panel.....</td> <td>1</td> <td>2</td> </tr> <tr> <td>Chair</td> <td>1</td> <td>2</td> </tr> <tr> <td>Sofa set</td> <td>1</td> <td>2</td> </tr> </tbody> </table>		Yes	No	Electricity	1	2	Radio	1	2	Television	1	2	Non-mobile telephone	1	2	Refrigerator.....	1	2	Solar Panel.....	1	2	Chair	1	2	Sofa set	1	2	
	Yes	No																											
Electricity	1	2																											
Radio	1	2																											
Television	1	2																											
Non-mobile telephone	1	2																											
Refrigerator.....	1	2																											
Solar Panel.....	1	2																											
Chair	1	2																											
Sofa set	1	2																											

[I] TABLE	Table.....1	2	
[J] CUPBOARD	Cupboard.....1	2	
[K] BED	Bed1	2	
[L] CLOCK	Clock.....1	2	
[M] CAMERA	Camera.....1	2	
[N] COMPUTER	Computer1	2	
HC9. DOES ANY MEMBER OF YOUR HOUSEHOLD OWN:		Yes No	
[A] A WATCH?	Watch1	2	
[B] A MOBILE TELEPHONE?	Mobile telephone1	2	
[C] A BICYCLE?	Bicycle1	2	
[D] A MOTORCYCLE OR SCOOTER?	Motorcycle / Scooter1	2	
[E] AN ANIMAL-DRAWN CART?	Animal-drawn cart.....1	2	
[F] A CAR OR TRUCK?	Car / Truck.....1	2	
[G] A BOAT WITH A MOTOR?	Boat with motor.....1	2	
HC10. DO YOU OR SOMEONE LIVING IN THIS HOUSEHOLD OWN THIS DWELLING? <i>If "No", then ask: DO YOU RENT THIS DWELLING FROM SOMEONE NOT LIVING IN THIS HOUSEHOLD? <i>If "Rented from someone else", circle "2". For other responses, circle "6".</i></i>	Own1 Rent2 Other (<i>specify</i>) _____ 6		
HC11. DOES ANY MEMBER OF THIS HOUSEHOLD OWN ANY LAND THAT CAN BE USED FOR AGRICULTURE?	Yes1 No2		2⇒HC13
HC12. HOW MANY HECTARES OF AGRICULTURAL LAND DO MEMBERS OF THIS HOUSEHOLD OWN? <i>If less than 1, record "00". If 95 or more, record '95'. If unknown, record '98'.</i>	Hectares — —		
HC13. DOES THIS HOUSEHOLD OWN ANY LIVESTOCK, HERDS, OTHER FARM ANIMALS, OR POULTRY?	Yes1 No2		2⇒HC15
HC14. HOW MANY OF THE FOLLOWING ANIMALS DOES THIS HOUSEHOLD HAVE?			
[A] CATTLE, MILK COWS, OR BULLS?	Cattle, milk cows, or bulls..... — —		
[B] HORSES, DONKEYS, OR MULES?	Horses, donkeys, or mules..... — —		
[C] GOATS?	Goats — —		
[D] SHEEP?	Sheep — —		

<p>[E] CHICKENS?</p> <p>[F] PIGS?</p> <p>[G] CAMELS</p> <p><i>If none, record '00'. If 95 or more, record '95'. If unknown, record '98'.</i></p>	<p>Chickens __ __</p> <p>Pigs __ __</p> <p>Camels __ __</p>	
<p>HC15. DOES ANY MEMBER OF THIS HOUSEHOLD HAVE A BANK ACCOUNT?</p>	<p>Yes 1</p> <p>No 2</p> <p>Dk 8</p>	

INSECTICIDE TREATED NETS		TN
TN1. DOES YOUR HOUSEHOLD HAVE ANY MOSQUITO NETS THAT CAN BE USED WHILE SLEEPING?	Yes 1 No 2	2⇒Next Module
TN2. HOW MANY MOSQUITO NETS DOES YOUR HOUSEHOLD HAVE?	Number of nets..... ____ ____	
TN3. Ask the respondent to show you the nets in the household. If more than 3 nets, use additional questionnaire(s).		

	1 st Net	2 nd Net	3 rd Net
TN4. Mosquito net observed?	Observed 1 Not observed..... 2	Observed..... 1 Not observed2	Observed 1 Not observed 2
TN5. Observe or ask the brand/type of mosquito net. <i>If brand is unknown and you cannot observe the net, show pictures of typical net types/brands to respondent.</i>	Long-lasting treated nets Perma Net 11 Olyset 12 Supernet 13 Other (specify) 16 DK brand 18 Pre-treated nets Supanet 21 Other (specify) 26 DK brand 28 Other net (specify) 36 DK brand / type 98	Long-lasting treated nets Perma Net 11 Olyset 12 Supernet 13 Other (specify) 16 DK brand 18 Pre-treated nets Supanet 21 Other (specify) 26 DK brand 28 Other net (specify) 36 DK brand / type 98	Long-lasting treated nets Perma Net 11 Olyset 12 Supernet 13 Other (specify) 16 DK brand 18 Pre-treated nets Supanet 21 Other (specify) 26 DK brand 28 Other net (specify) 36 DK brand / type 98
TN6. HOW MANY MONTHS AGO DID YOUR HOUSEHOLD GET THE MOSQUITO NET? <i>If less than one month, record "00"</i>	Months ago ____ ____ More than 36 mo. ago... 95 DK / Not sure 98	Months ago ____ ____ More than 36 mo. ago ... 95 DK / Not sure 98	Months ago ____ ____ More than 36 mo. ago... 95 DK / Not sure 98
TN7. Check TN5 for type of net	<input type="checkbox"/> Long-lasting (11-18) ⇒ TN11 <input type="checkbox"/> Pre-treated (21-28) ⇒ TN9 <input type="checkbox"/> Else ⇒ Continue	<input type="checkbox"/> Long-lasting (11-18) ⇒ TN11 <input type="checkbox"/> Pre-treated (21-28) ⇒ TN9 <input type="checkbox"/> Else ⇒ Continue	<input type="checkbox"/> Long-lasting (11-18) ⇒ TN11 <input type="checkbox"/> Pre-treated (21-28) ⇒ TN9 <input type="checkbox"/> Else ⇒ Continue
TN8. WHEN YOU GOT THE NET, WAS IT ALREADY TREATED WITH AN INSECTICIDE TO KILL OR REPEL MOSQUITOES?	Yes 1 No 2 DK / Not sure 8	Yes 1 No 2 DK / Not sure 8	Yes 1 No 2 DK / Not sure 8
TN9. SINCE YOU GOT THE NET, WAS IT EVER SOAKED OR DIPPED IN A LIQUID TO KILL OR REPEL MOSQUITOES?	Yes 1 No 2 ⇒ TN11 DK / Not sure 8 ⇒ TN11	Yes 1 No 2 ⇒ TN11 DK / Not sure 8 ⇒ TN11	Yes 1 No 2 ⇒ TN11 DK / Not sure 8 ⇒ TN11

<p>TN10. HOW MANY MONTHS AGO WAS THE NET LAST SOAKED OR DIPPED?</p> <p><i>If less than one month, record "00"</i></p>	<p>Months ago ____ ____</p> <p>More than 24 mo. ago... 95</p> <p>DK / Not sure 98</p>	<p>Months ago ____ ____</p> <p>More than 24 mo. ago ...95</p> <p>DK / Not sure 98</p>	<p>Months ago ____ ____</p> <p>More than 24 mo. ago... 95</p> <p>DK / Not sure 98</p>
<p>TN11. DID ANYONE SLEEP UNDER THIS MOSQUITO NET LAST NIGHT?</p>	<p>Yes 1</p> <p>No 2</p> <p style="text-align: right;">⇒ TN13</p> <p>DK / Not sure 8</p> <p style="text-align: right;">⇒ TN13</p>	<p>Yes 1</p> <p>No 2</p> <p style="text-align: right;">⇒ TN13</p> <p>DK / Not sure 8</p> <p style="text-align: right;">⇒ TN13</p>	<p>Yes 1</p> <p>No 2</p> <p style="text-align: right;">⇒ TN13</p> <p>DK / Not sure 8</p> <p style="text-align: right;">⇒ TN13</p>
<p>TN12. WHO SLEPT UNDER THIS MOSQUITO NET LAST NIGHT?</p> <p><i>Record the person's line number from the List of Household Members</i></p> <p><i>If someone not in the List of Household Members slept under the mosquito net, record "00"</i></p>	<p>Name _____</p> <p>Line number ____ ____</p> <p>Name _____</p> <p>Line number ____ ____</p> <p>Name _____</p> <p>Line number ____ ____</p> <p>Name _____</p> <p>Line number ____ ____</p> <p>Name _____</p> <p>Line number ____ ____</p> <p>Name _____</p> <p>Line number ____ ____</p> <p>Name _____</p> <p>Line number ____ ____</p>	<p>Name _____</p> <p>Line number ____ ____</p> <p>Name _____</p> <p>Line number ____ ____</p> <p>Name _____</p> <p>Line number ____ ____</p> <p>Name _____</p> <p>Line number ____ ____</p> <p>Name _____</p> <p>Line number ____ ____</p> <p>Name _____</p> <p>Line number ____ ____</p>	<p>Name _____</p> <p>Line number ____ ____</p> <p>Name _____</p> <p>Line number ____ ____</p> <p>Name _____</p> <p>Line number ____ ____</p> <p>Name _____</p> <p>Line number ____ ____</p> <p>Name _____</p> <p>Line number ____ ____</p> <p>Name _____</p> <p>Line number ____ ____</p>
<p>TN13.</p>	<p><i>Go back to TN4 for next net. If no more nets, go to next module</i></p>	<p><i>Go back to TN4 for next net. If no more nets, go to next module</i></p>	<p><i>Go back to TN4 in first column of a new questionnaire for next net. If no more nets, go to next module</i></p>
			<p><i>Tick here if additional questionnaire used</i> <input type="checkbox"/></p>

INDOOR RESIDUAL SPRAYING		IR
IR1. AT ANY TIME IN THE PAST 12 MONTHS, HAS ANYONE COME INTO YOUR DWELLING TO SPRAY THE INTERIOR WALLS AGAINST MOSQUITOES?	Yes 1 No 2 DK 8	2⇒Next Module 8⇒Next Module
IR2. WHO SPRAYED THE DWELLING? <i>Circle all that apply.</i>	Government worker / programA Private companyB Non-governmental organizationC Other (<i>specify</i>)X DKZ	

WATER AND SANITATION		WS
WS1. WHAT IS THE <u>MAIN</u> SOURCE OF DRINKING WATER FOR MEMBERS OF YOUR HOUSEHOLD?	Piped water Piped into dwelling 11 Piped into compound, yard or plot 12 Piped to neighbour 13 Public tap / standpipe 14 Tube Well, Borehole.....21 Dug well Protected well 31 Unprotected well..... 32 Water from spring Protected spring 41 Unprotected spring 42 Rainwater collection 51 Tanker-truck 61 Cart with small tank / drum..... 71 Surface water (river, stream, dam, lake, pond, canal, irrigation channel) 81 Bottled water 91 Other (<i>specify</i>) 96	11⇒WS6 12⇒WS6 13⇒WS6 14⇒WS3 21⇒WS3 31⇒WS3 32⇒WS3 41⇒WS3 42⇒WS3 51⇒WS3 61⇒WS3 71⇒WS3 81⇒WS3 96⇒WS3
WS2. WHAT IS THE <u>MAIN</u> SOURCE OF WATER USED BY YOUR HOUSEHOLD FOR OTHER PURPOSES SUCH AS COOKING AND HANDWASHING?	Piped water Piped into dwelling 11 Piped into compound, yard or plot 12 Piped to neighbour 13 Public tap / standpipe 14 Tube Well, Borehole.....21 Dug well Protected well 31 Unprotected well..... 32 Water from spring Protected spring 41 Unprotected spring 42 Rainwater collection 51 Tanker-truck 61 Cart with small tank / drum..... 71 Surface water (river, stream, dam, lake, pond, canal, irrigation channel) 81 Other (<i>specify</i>) 96	11⇒WS6 12⇒WS6 13⇒WS6 14⇒WS3 21⇒WS3 31⇒WS3 32⇒WS3 41⇒WS3 42⇒WS3 51⇒WS3 61⇒WS3 71⇒WS3 81⇒WS3 96⇒WS3
WS3. WHERE IS THAT WATER SOURCE LOCATED?	In own dwelling 1 In own yard / plot 2 Elsewhere 3	1⇒WS6 2⇒WS6
WS4. HOW LONG DOES IT TAKE TO GO THERE, GET WATER, AND COME BACK?	Number of minutes ___ ___ ___ DK 998	

<p>WS5. WHO USUALLY GOES TO THIS SOURCE TO COLLECT THE WATER FOR YOUR HOUSEHOLD?</p> <p><i>Probe:</i> IS THIS PERSON UNDER AGE 15? WHAT SEX?</p>	<p>Adult woman (age 15+ years) 1 Adult man (age 15+ years)..... 2 Female child (under 15) 3 Male child (under 15) 4</p> <p>DK 8</p>	
<p>WS6. DO YOU DO ANYTHING TO THE WATER TO MAKE IT SAFER TO DRINK?</p>	<p>Yes 1 No 2</p> <p>DK 8</p>	<p>2⇒WS8 8⇒WS8</p>
<p>WS7. WHAT DO YOU USUALLY DO TO MAKE THE WATER SAFER TO DRINK?</p> <p><i>Probe:</i> ANYTHING ELSE?</p> <p><i>Record all items mentioned.</i></p>	<p>Boil A Add bleach / chlorine B Strain it through a cloth C Use water filter (ceramic, sand, composite, etc.) D Solar disinfection E Let it stand and settle F</p> <p>Other (<i>specify</i>) X DK Z</p>	
<p>WS8. WHAT KIND OF TOILET FACILITY DO MEMBERS OF YOUR HOUSEHOLD USUALLY USE?</p> <p><i>If “flush” or “pour flush”, probe:</i> WHERE DOES IT FLUSH TO?</p> <p><i>If not possible to determine, ask permission to observe the facility.</i></p>	<p>Flush / Pour flush Flush to piped sewer system 11 Flush to septic tank 12 Flush to pit (latrine)..... 13 Flush to somewhere else 14 Flush to unknown place / Not sure / DK where 15</p> <p>Pit latrine Ventilated Improved Pit latrine (VIP) 21 Pit latrine with slab 22 Pit latrine without slab / Open pit..... 23</p> <p>Composting toilet..... 31 Bucket 41 Hanging toilet, Hanging latrine 51</p> <p>No facility, Bush, Field..... 95</p> <p>Other (<i>specify</i>) 96</p>	<p>95⇒Next Module</p>
<p>WS9. DO YOU SHARE THIS FACILITY WITH OTHERS WHO ARE NOT MEMBERS OF YOUR HOUSEHOLD?</p>	<p>Yes 1 No 2</p>	<p>2⇒Next Module</p>
<p>WS10. DO YOU SHARE THIS FACILITY ONLY WITH MEMBERS OF OTHER HOUSEHOLDS THAT YOU KNOW, OR IS THE FACILITY OPEN TO THE USE OF THE GENERAL PUBLIC?</p>	<p>Other households only (not public) 1 Public facility..... 2</p>	<p>2⇒Next Module</p>
<p>WS11. HOW MANY HOUSEHOLDS IN TOTAL USE THIS TOILET FACILITY, INCLUDING YOUR OWN HOUSEHOLD?</p>	<p>Number of households (if less than 10) 0 __</p> <p>Ten or more households 10</p> <p>DK 98</p>	

HANDWASHING	HW	
<p>HW1. WE WOULD LIKE TO LEARN ABOUT THE PLACES THAT HOUSEHOLDS USE TO WASH THEIR HANDS.</p> <p>CAN YOU PLEASE SHOW ME WHERE MEMBERS OF YOUR HOUSEHOLD <u>MOST OFTEN</u> WASH THEIR HANDS?</p>	<p>Observed 1</p> <p>Not observed</p> <p>Moving basin/kettle/bucket..... 2</p> <p>Not in dwelling / plot / yard..... 3</p> <p>No permission to see 4</p> <p>Other reason (specify)..... 6</p>	<p>2 ⇨HW4</p> <p>3 ⇨HW4</p> <p>4 ⇨HW4</p> <p>6 ⇨HW4</p>
<p>HW2. <i>Observe presence of water at the place for handwashing.</i></p> <p><i>Verify by checking the tap/pump, or basin, bucket, water container or similar objects for presence of water.</i></p>	<p>Water is available..... 1</p> <p>Water is not available 2</p>	
<p>HW3A. <i>Is soap, detergent or ash/mud/sand present at the place for handwashing?</i></p>	<p>Yes, present..... 1</p> <p>No, not present 2</p>	<p>2⇨HW4</p>
<p>HW3B. <i>Record your observation.</i></p> <p><i>Circle all that apply.</i></p>	<p>Bar soap.....A</p> <p>Detergent (Powder / Liquid / Paste).....B</p> <p>Liquid soap.....C</p> <p>Ash / Mud / SandD</p>	<p>A⇨HH19</p> <p>B⇨HH19</p> <p>C⇨HH19</p> <p>D⇨HH19</p>
<p>HW4. DO YOU HAVE ANY SOAP OR DETERGENT OR ASH/MUD/SAND IN YOUR HOUSE FOR WASHING HANDS?</p>	<p>Yes..... 1</p> <p>No 2</p>	<p>2⇨HH19</p>
<p>HW5A. CAN YOU PLEASE SHOW IT TO ME?</p>	<p>Yes, shown 1</p> <p>No, not shown 2</p>	<p>2⇨HH19</p>
<p>HW5B. <i>Record your observation.</i></p> <p><i>Circle all that apply.</i></p>	<p>Bar soap.....A</p> <p>Detergent (Powder / Liquid / Paste).....B</p> <p>Liquid soap.....C</p> <p>Ash / Mud / SandD</p>	

HH19. Record the time.	Hour and minutes : ..	
-------------------------------	-----------------------------	--

SALT IODIZATION		SI
<p>SI1. WE WOULD LIKE TO CHECK WHETHER THE SALT USED IN YOUR HOUSEHOLD IS IODIZED. MAY I HAVE A SAMPLE OF THE SALT USED TO <u>COOK MEALS</u> IN YOUR HOUSEHOLD?</p> <p><i>Once you have tested the salt, circle number that corresponds to test outcome.</i></p>	<p>Not iodized - 0 PPM 1</p> <p>More than 0 PPM & less than 15 PPM..... 2</p> <p>15 PPM or more 3</p> <p>No salt in the house..... 4</p> <p>Salt not tested (specify reason) _____ 5</p>	

<p>HH20. Thank the respondent for his/her cooperation and check the List of Household Members:</p> <p><input type="checkbox"/> A separate <i>QUESTIONNAIRE FOR INDIVIDUAL WOMEN</i> has been issued for each woman age 15-49 years in the List of Household Members (HL7)</p> <p><input type="checkbox"/> A separate <i>QUESTIONNAIRE FOR CHILDREN UNDER FIVE</i> has been issued for each child under age 5 years in the List of Household Members (HL7B)</p> <p><i>Return to the cover page and make sure that all information is entered, including the number of eligible women (HH12) and under-5s (HH14)</i></p> <p><i>Make arrangements for the administration of the remaining questionnaire(s) in this household.</i></p>

Interviewer's Observations

Field Editor's Observations

Supervisor's Observations

QUESTIONNAIRE FOR INDIVIDUAL WOMEN WESTERN AND NORTH RIFT SURVEY



WOMAN'S INFORMATION PANEL	WM
<i>This questionnaire is to be administered to all women age 15 through 49 (see List of Household Members, column HL7). A separate questionnaire should be used for each eligible woman.</i>	
WM1. Cluster number: <div style="text-align: right;">_____</div>	WM2. Household number: <div style="text-align: right;">_____</div>
WM3. Woman's name: Name _____	WM4. Woman's line number: <div style="text-align: right;">_____</div>
WM5. Interviewer's name and number: Name _____	WM6. Day/Month/Year of interview: <div style="text-align: right;">_____ / _____ / 201__</div>

<p><i>Repeat greeting if not already read to this woman:</i></p> <p>WE ARE FROM THE UNIVERSITY OF NAIROBI AND KENYA NATIONAL BUREAU OF STATISTICS. WE ARE CONDUCTING A SURVEY ABOUT THE SITUATION OF CHILDREN, FAMILIES AND HOUSEHOLDS. I WOULD LIKE TO TALK TO YOU ABOUT THESE SUBJECTS. THE INTERVIEW WILL TAKE ABOUT 45 MINUTES. ALL THE INFORMATION WE OBTAIN WILL REMAIN STRICTLY CONFIDENTIAL AND ANONYMOUS.</p>	<p><i>If greeting at the beginning of the household questionnaire has already been read to this woman, then read the following:</i></p> <p>NOW I WOULD LIKE TO TALK TO YOU MORE ABOUT YOUR HEALTH AND OTHER TOPICS. THIS INTERVIEW WILL TAKE ABOUT 45 MINUTES. AGAIN, ALL THE INFORMATION WE OBTAIN WILL REMAIN STRICTLY CONFIDENTIAL AND ANONYMOUS.</p>
<p>MAY I START NOW?</p> <p><input type="checkbox"/> <i>Yes, permission is given ⇒ Go to WM10 to record the time and then begin the interview.</i></p> <p><input type="checkbox"/> <i>No, permission is not given ⇒ Circle '03' in WM7. Discuss this result with your supervisor.</i></p>	

WM7. Result of woman's interview	Completed..... 01 Not at home..... 02 Refused..... 03 Partly completed 04 Incapacitated..... 05 Other (specify) _____ 96
---	---

WM8. Field editor's name and number: Name _____	WM9. Main data entry clerk's name and number: Name _____
---	--

WM10. Record the time.	Hour and minutes : ..	
-------------------------------	-----------------------------	--

WOMAN'S BACKGROUND		WB
WB1. IN WHAT MONTH AND YEAR WERE YOU BORN?	Date of birth Month..... DK month.....98 Year DK year.....9998	
WB2. HOW OLD ARE YOU? <i>Probe: HOW OLD WERE YOU AT YOUR LAST BIRTHDAY?</i> <i>Compare and correct WB1 and/or WB2 if inconsistent</i>	Age (in completed years)	
WB3. HAVE YOU EVER ATTENDED SCHOOL OR PRESCHOOL?	Yes1 No2	2⇒WB7
WB4. WHAT IS THE HIGHEST LEVEL OF SCHOOL YOU ATTENDED?	Preschool0 Primary1 Secondary2 Higher3	0⇒WB7
WB5. WHAT IS THE HIGHEST GRADE YOU COMPLETED AT THAT LEVEL? <i>If the first grade at this level is not completed, enter "00"</i>	Grade	
WB6. Check WB4:		
<input type="checkbox"/> Secondary or higher (WB4=2 or 3) ⇒ Go to Next Module <input type="checkbox"/> Primary (WB4=1) ⇒ Continue with WB7		
WB7. NOW I WOULD LIKE YOU TO READ THIS SENTENCE TO ME. <i>Show sentence on the card to the respondent. If respondent cannot read whole sentence, probe:</i> CAN YOU READ PART OF THE SENTENCE TO ME?	Cannot read at all1 Able to read only parts of sentence.....2 Able to read whole sentence3 No sentence in required language4 <i>(specify language)</i> Blind/visually impaired.....5	

ACCESS TO MASS MEDIA AND USE OF INFORMATION/COMMUNICATION TECHNOLOGY		MT
MT1. Check WB7: <input type="checkbox"/> Question left blank (Respondent has secondary or higher education) ⇒ Continue with MT2 <input type="checkbox"/> Able to read or no sentence in required language (WB7 = 2, 3 or 4) ⇒ Continue with MT2 <input type="checkbox"/> Cannot read at all or blind/visually impaired (WB7 = 1 or 5) ⇒ Go to MT3		
MT2. HOW OFTEN DO YOU READ A NEWSPAPER OR MAGAZINE: ALMOST EVERY DAY, AT LEAST ONCE A WEEK, LESS THAN ONCE A WEEK OR NOT AT ALL?	Almost every day 1 At least once a week 2 Less than once a week 3 Not at all 4	
MT3. DO YOU LISTEN TO THE RADIO ALMOST EVERY DAY, AT LEAST ONCE A WEEK, LESS THAN ONCE A WEEK OR NOT AT ALL?	Almost every day 1 At least once a week 2 Less than once a week 3 Not at all 4	
MT4. HOW OFTEN DO YOU WATCH TELEVISION: WOULD YOU SAY THAT YOU WATCH ALMOST EVERY DAY, AT LEAST ONCE A WEEK, LESS THAN ONCE A WEEK OR NOT AT ALL?	Almost every day 1 At least once a week 2 Less than once a week 3 Not at all 4	
MT5. Check WB2: Age of respondent? <input type="checkbox"/> Age 15-24 ⇒ Continue with MT6 <input type="checkbox"/> Age 25-49 ⇒ Go to Next Module		
MT6. HAVE YOU EVER USED A COMPUTER?	Yes 1 No 2	2 ⇒ MT9
MT7. HAVE YOU USED A COMPUTER FROM ANY LOCATION IN THE LAST 12 MONTHS?	Yes 1 No 2	2 ⇒ MT9
MT8. DURING THE LAST ONE MONTH, HOW OFTEN DID YOU USE A COMPUTER: ALMOST EVERY DAY, AT LEAST ONCE A WEEK, LESS THAN ONCE A WEEK OR NOT AT ALL?	Almost every day 1 At least once a week 2 Less than once a week 3 Not at all 4	
MT9. HAVE YOU EVER USED THE INTERNET?	Yes 1 No 2	2 ⇒ Next Module
MT10. IN THE LAST 12 MONTHS, HAVE YOU USED THE INTERNET? <i>If necessary, probe for use from any location, with any device.</i>	Yes 1 No 2	2 ⇒ Next Module
MT11. DURING THE LAST ONE MONTH, HOW OFTEN DID YOU USE THE INTERNET: ALMOST EVERY DAY, AT LEAST ONCE A WEEK, LESS THAN ONCE A WEEK OR NOT AT ALL?	Almost every day 1 At least once a week 2 Less than once a week 3 Not at all 4	

FERTILITY/BIRTH HISTORY		CM
CM1. NOW I WOULD LIKE TO ASK ABOUT ALL THE BIRTHS YOU HAVE HAD DURING YOUR LIFE. HAVE YOU EVER GIVEN BIRTH?	Yes 1 No 2	2⇒CM8
CM4. DO YOU HAVE ANY SONS OR DAUGHTERS TO WHOM YOU HAVE GIVEN BIRTH WHO ARE NOW LIVING WITH YOU?	Yes 1 No 2	2⇒CM6
CM5. HOW MANY SONS LIVE WITH YOU? HOW MANY DAUGHTERS LIVE WITH YOU? <i>If none, record '00'.</i>	Sons at home __ __ Daughters at home __ __	
CM6. DO YOU HAVE ANY SONS OR DAUGHTERS TO WHOM YOU HAVE GIVEN BIRTH WHO ARE ALIVE BUT DO NOT LIVE WITH YOU?	Yes 1 No 2	2⇒CM8
CM7. HOW MANY SONS ARE ALIVE BUT DO NOT LIVE WITH YOU? HOW MANY DAUGHTERS ARE ALIVE BUT DO NOT LIVE WITH YOU? <i>If none, record '00'.</i>	Sons elsewhere __ __ Daughters elsewhere __ __	
CM8. HAVE YOU EVER GIVEN BIRTH TO A BOY OR GIRL WHO WAS BORN ALIVE BUT LATER DIED? <i>If "No" probe by asking: I MEAN, TO A CHILD WHO EVER BREATHED OR CRIED OR SHOWED OTHER SIGNS OF LIFE – EVEN IF HE OR SHE LIVED ONLY A FEW MINUTES OR HOURS?</i>	Yes 1 No 2	2⇒CM10
CM9. HOW MANY BOYS HAVE DIED? HOW MANY GIRLS HAVE DIED? <i>If none, record '00'.</i>	Boys dead __ __ Girls dead __ __	
CM10. <i>Sum answers to CM5, CM7, and CM9.</i>	Sum __ __	
CM11. JUST TO MAKE SURE THAT I HAVE THIS RIGHT, YOU HAVE HAD IN TOTAL (<i>total number in CM10</i>) LIVE BIRTHS DURING YOUR LIFE. IS THIS CORRECT? <input type="checkbox"/> <i>Yes. Check below:</i> <ul style="list-style-type: none"> <input type="checkbox"/> <i>No live births ⇒ Go to ILLNESS SYMPTOMS Module</i> <input type="checkbox"/> <i>One or more live births ⇒ Continue with the BIRTH HISTORY module</i> <input type="checkbox"/> <i>No. ⇒ Check responses to CM1-CM10 and make corrections as necessary before proceeding to the BIRTH HISTORY Module or ILLNESS SYMPTOMS Module</i>		

BIRTH HISTORY													BH		
<p>NOW I WOULD LIKE TO RECORD THE NAMES OF ALL OF YOUR BIRTHS, WHETHER STILL ALIVE OR NOT, STARTING WITH THE FIRST ONE YOU HAD. <i>Record names of all of the births in BH1. Record twins and triplets on separate lines. If there are more than 14 births, use an additional questionnaire.</i></p>															
BH Line No.	BH1. WHAT NAME WAS GIVEN TO YOUR (first/next) BABY?	BH2. WERE ANY OF THESE BIRTHS TWINS?	BH3. IS (name) A BOY OR A GIRL?	BH4. IN WHAT MONTH AND YEAR WAS (name) BORN? <i>Probe: WHAT IS HIS/HER BIRTHDAY?</i>		BH5. IS (name) STILL ALIVE?		BH6. HOW OLD WAS (name) AT HIS/HER LAST BIRTHDAY?	BH7. IS (name) LIVING WITH YOU?		BH8. <i>Record household line number of child (from HLI)</i>	BH9. <i>If dead:</i> HOW OLD WAS (name) WHEN HE/SHE DIED? <i>If "1 year", probe:</i> HOW MANY MONTHS OLD WAS (name)? <i>Record days if less than 1 month; record months if less than 2 years; or years</i>		BH10. WERE THERE ANY OTHER LIVE BIRTHS BETWEEN (name of previous birth) AND (name), INCLUDING ANY CHILDREN WHO DIED AFTER BIRTH?	
		1 Single 2 Multiple	1 Boy 2 Girl			1 Yes 2 No		<i>Record age in completed years.</i>	1 Yes 2 No		<i>Record "00" if child is not listed.</i>		1 Yes 2 No		
Line	Name	SM	BG	Month	Year	Y	N	Age	Y	N	Line No	Unit	Number	Y	N
01		1 2	1 2	___	___	1	2	___	1	2	___	Days 1 Months 2 Years 3			
							↓ BH9				⇒Next Line				
02		1 2	1 2	___	___	1	2	___	1	2	___	Days 1 Months 2 Years 3		1	2
							↓ BH9				⇒BH10			Add Birth	Next Birth
03		1 2	1 2	___	___	1	2	___	1	2	___	Days 1 Months 2 Years 3		1	2
							↓ BH9				⇒BH10			Add Birth	Next Birth
04		1 2	1 2	___	___	1	2	___	1	2	___	Days 1 Months 2 Years 3		1	2
							↓ BH9				⇒BH10			Add Birth	Next Birth
05		1 2	1 2	___	___	1	2	___	1	2	___	Days 1 Months 2 Years 3		1	2
							↓ BH9				⇒BH10			Add Birth	Next Birth
06		1 2	1 2	___	___	1	2	___	1	2	___	Days 1 Months 2 Years 3		1	2
							↓ BH9				⇒BH10			Add Birth	Next Birth
07		1 2	1 2	___	___	1	2	___	1	2	___	Days 1 Months 2 Years 3		1	2
							↓ BH9				⇒BH10			Add Birth	Next Birth

<i>BH Line No.</i>	BH1. WHAT NAME WAS GIVEN TO YOUR (<i>first/next</i>) BABY?	BH2. WERE ANY OF THESE BIRTHS TWINS? 1 Single 2 Multiple	BH3. IS (<i>name</i>) A BOY OR A GIRL? 1 Boy 2 Girl	BH4. IN WHAT MONTH AND YEAR WAS (<i>name</i>) BORN? <i>Probe: WHAT IS HIS/HER BIRTHDAY?</i>		BH5. IS (<i>name</i>) STILL ALIVE? 1 Yes 2 No	BH6. HOW OLD WAS (<i>name</i>) AT HIS/HER LAST BIRTHDAY? <i>Record age in completed years.</i>	BH7. IS (<i>name</i>) LIVING WITH YOU? 1 Yes 2 No	BH8. <i>Record household line number of child (from HLI)</i> <i>Record "00" if child is not listed.</i>	BH9. <i>If dead:</i> HOW OLD WAS (<i>name</i>) WHEN HE/SHE DIED? <i>If "1 year", probe:</i> HOW MANY MONTHS OLD WAS (<i>name</i>)? <i>Record days if less than 1 month; record months if less than 2 years; or years</i>		BH10. WERE THERE ANY OTHER LIVE BIRTHS BETWEEN (<i>name of previous birth</i>) AND (<i>name</i>), INCLUDING ANY CHILDREN WHO DIED AFTER BIRTH? 1 Yes 2 No
08		1 2	1 2	___	___	1 2 ↓ BH9	___	1 2	___ ⇒BH10	Days 1 Months 2 Years 3	___	1 2 Add Next Birth Birth
09		1 2	1 2	___	___	1 2 ↓ BH9	___	1 2	___ ⇒BH10	Days 1 Months 2 Years 3	___	1 2 Add Next Birth Birth
10		1 2	1 2	___	___	1 2 ↓ BH9	___	1 2	___ ⇒BH10	Days 1 Months 2 Years 3	___	1 2 Add Next Birth Birth
11		1 2	1 2	___	___	1 2 ↓ BH9	___	1 2	___ ⇒BH10	Days 1 Months 2 Years 3	___	1 2 Add Next Birth Birth
12		1 2	1 2	___	___	1 2 ↓ BH9	___	1 2	___ ⇒BH10	Days 1 Months 2 Years 3	___	1 2 Add Next Birth Birth
13		1 2	1 2	___	___	1 2 ↓ BH9	___	1 2	___ ⇒BH10	Days 1 Months 2 Years 3	___	1 2 Add Next Birth Birth
14		1 2	1 2	___	___	1 2 ↓ BH9	___	1 2	___ ⇒BH10	Days 1 Months 2 Years 3	___	1 2 Add Next Birth Birth
BH11. HAVE YOU HAD ANY LIVE BIRTHS SINCE THE BIRTH OF (<i>name of last birth in BIRTH HISTORY Module</i>)?							Yes..... 1 No 2				1⇒Record birth(s) in Birth History	

CM12A. Compare number in CM10 with number of births in the *BIRTH HISTORY* Module above and check:

- Numbers are same ⇒ Continue with CM13
- Numbers are different ⇒ Probe and reconcile

CM13. Check BH4 in *BIRTH HISTORY* Module: Last birth occurred within the last 2 years, that is, since (month of interview) in **2011** (if the month of interview and the month of birth are the same, and the year of birth is **2011**, consider this as a birth within the last 2 years)

- No live birth in last 2 years. ⇒ Go to *ILLNESS SYMPTOMS* Module.
- One or more live births in last 2 years. ⇒ Record name of last born child and continue with Next Module

Name of last-born child _____

If child has died, take special care when referring to this child by name in the following modules.

DESIRE FOR LAST BIRTH		DB
<p><i>This module is to be administered to all women with a live birth in the 2 years preceding the date of interview. Record name of last-born child from CM13 here _____. Use this child's name in the following questions, where indicated.</i></p>		
<p>DB1. WHEN YOU GOT PREGNANT WITH (<i>name</i>), DID YOU WANT TO GET PREGNANT AT THAT TIME?</p>	<p>Yes 1</p> <p>No 2</p>	<p>1⇒Next Module</p>
<p>DB2. DID YOU WANT TO HAVE A BABY LATER ON, OR DID YOU NOT WANT ANY (MORE) CHILDREN?</p>	<p>Later 1</p> <p>No more..... 2</p>	<p>2⇒Next Module</p>
<p>DB3. HOW MUCH LONGER DID YOU WANT TO WAIT?</p> <p><i>Record the answer as stated by respondent.</i></p>	<p>Months..... 1 __ __</p> <p>Years 2 __ __</p> <p>DK..... 998</p>	

MATERNAL AND NEWBORN HEALTH		MN												
<p><i>This module is to be administered to all women with a live birth in the 2 years preceding the date of interview. Record name of last-born child from CM13 here _____. Use this child's name in the following questions, where indicated.</i></p>														
<p>MN1. DID YOU SEE ANYONE FOR ANTENATAL CARE DURING YOUR PREGNANCY WITH (name)?</p>	<p>Yes 1 No 2</p>	2⇒MN5												
<p>MN2. WHOM DID YOU SEE?</p> <p><i>Probe:</i> ANYONE ELSE?</p> <p><i>Probe for the type of person seen and circle all answers given.</i></p>	<p>Health professional: Doctor A Nurse/Midwife B Clinical Officer C Community Nurse D Other person Relative / friend E Traditional birth attendant F Community health worker G Other (specify) X</p>													
<p>MN2A. HOW MANY WEEKS OR MONTHS PREGNANT WERE YOU WHEN YOU FIRST RECEIVED ANTENATAL CARE FOR THIS PREGNANCY?</p> <p><i>Record the answer as stated by respondent.</i></p>	<p>Weeks 1 ___ Months 2 0 ___ DK 998</p>													
<p>MN3. HOW MANY TIMES DID YOU RECEIVE ANTENATAL CARE DURING THIS PREGNANCY?</p> <p><i>Probe to identify the number of times antenatal care was received. If a range is given, record the minimum number of times antenatal care received.</i></p>	<p>Number of times ___ DK 98</p>													
<p>MN4. AS PART OF YOUR ANTENATAL CARE DURING THIS PREGNANCY, WERE ANY OF THE FOLLOWING DONE AT LEAST ONCE:</p> <p>[A] WAS YOUR BLOOD PRESSURE MEASURED?</p> <p>[B] DID YOU GIVE A URINE SAMPLE?</p> <p>[C] DID YOU GIVE A BLOOD SAMPLE?</p>	<table border="0"> <thead> <tr> <th></th> <th>Yes</th> <th>No</th> </tr> </thead> <tbody> <tr> <td>Blood pressure</td> <td>1</td> <td>2</td> </tr> <tr> <td>Urine sample</td> <td>1</td> <td>2</td> </tr> <tr> <td>Blood sample</td> <td>1</td> <td>2</td> </tr> </tbody> </table>		Yes	No	Blood pressure	1	2	Urine sample	1	2	Blood sample	1	2	
	Yes	No												
Blood pressure	1	2												
Urine sample	1	2												
Blood sample	1	2												
<p>MN5. DO YOU HAVE A CARD OR OTHER DOCUMENT WITH YOUR OWN IMMUNIZATIONS LISTED?</p> <p>MAY I SEE IT PLEASE?</p> <p><i>If a card is presented, use it to assist with answers to the following questions.</i></p>	<p>Yes (card seen) 1 Yes (card not seen) 2 No 3 DK 8</p>													
<p>MN6. WHEN YOU WERE PREGNANT WITH (name), DID YOU RECEIVE ANY INJECTION IN THE ARM OR SHOULDER TO PREVENT THE BABY FROM GETTING TETANUS, THAT IS CONVULSIONS AFTER BIRTH?</p>	<p>Yes 1 No 2 DK 8</p>	2⇒MN9 8⇒MN9												
<p>MN7. HOW MANY TIMES DID YOU RECEIVE THIS TETANUS INJECTION DURING YOUR PREGNANCY WITH (name)?</p>	<p>Number of times ___ DK 8</p>	8⇒MN9												

MN8. How many tetanus injections during last pregnancy were reported in MN7? <input type="checkbox"/> At least two tetanus injections during last pregnancy. ⇒ Go to MN12 <input type="checkbox"/> Only one tetanus injection during last pregnancy. ⇒ Continue with MN9		
MN9. DID YOU RECEIVE ANY TETANUS INJECTION AT ANY TIME BEFORE YOUR PREGNANCY WITH (<i>name</i>), EITHER TO PROTECT YOURSELF OR ANOTHER BABY?	Yes 1 No 2 DK 8	 2⇒MN12 8⇒MN12
MN10. HOW MANY TIMES DID YOU RECEIVE A TETANUS INJECTION BEFORE YOUR PREGNANCY WITH (<i>name</i>)? <i>If 7 or more times, record '7'.</i>	Number of times..... DK 8	 8⇒MN12
MN11. HOW MANY YEARS AGO DID YOU RECEIVE THE LAST TETANUS INJECTION BEFORE YOUR PREGNANCY WITH (<i>name</i>)? <i>If less than 1 year, record '00'.</i>	Years ago.....	
MN12. Check MN1 for presence of antenatal care during this pregnancy: <input type="checkbox"/> Yes, antenatal care received. ⇒ Continue with MN13 <input type="checkbox"/> No antenatal care received ⇒ Go to MN17		
MN13. DURING (ANY OF) YOUR ANTENATAL VISIT(S) FOR THE PREGNANCY WITH (<i>name</i>), DID YOU TAKE ANY MEDICINE IN ORDER TO PREVENT YOU FROM GETTING MALARIA?	Yes 1 No 2 DK 8	 2⇒MN17 8⇒MN17
MN14. WHICH MEDICINES DID YOU TAKE TO PREVENT MALARIA? <i>Circle all medicines taken. If type of medicine is not determined, show typical anti-malarial to respondent.</i>	SP/Fansidar A Chloroquine B Other (<i>specify</i>) X DK Z	
MN15. Check MN14 for medicine taken: <input type="checkbox"/> SP/Fansidar taken. ⇒ Continue with MN16 <input type="checkbox"/> SP/Fansidar not taken. ⇒ Go to MN17		
MN16. DURING YOUR PREGNANCY WITH (<i>name</i>), HOW MANY TIMES DID YOU TAKE SP/FANSIDAR IN TOTAL? PLEASE INCLUDE ALL THAT YOU OBTAINED EITHER DURING AN ANTENATAL CARE VISIT, DURING A VISIT TO A HEALTH FACILITY OR FROM ANOTHER SOURCE?	Number of times..... DK 98	

<p>MN17. WHO ASSISTED WITH THE DELIVERY OF (name)?</p> <p><i>Probe:</i> ANYONE ELSE?</p> <p><i>Probe for the type of person assisting and circle all answers given.</i></p> <p><i>If respondent says no one assisted, probe to determine whether any adults were present at the delivery.</i></p>	<p>Health professional:</p> <p>Doctor A</p> <p>Nurse / Midwife B</p> <p>Clinical Officer C</p> <p>Community Nurse D</p> <p>Other person</p> <p>Traditional birth attendant F</p> <p>Community health worker G</p> <p>Relative / Friend H</p> <p>Other (<i>specify</i>) X</p> <p>No one Y</p>	
<p>MN18. WHERE DID YOU GIVE BIRTH TO (name)?</p> <p><i>Probe to identify the type of source.</i></p> <p><i>If unable to determine whether public or private, write the name of the place.</i></p> <p>_____</p> <p>(<i>Name of place</i>)</p>	<p>Home</p> <p>Respondent's home 11</p> <p>Other home 12</p> <p>Public sector</p> <p>Government hospital 21</p> <p>Government clinic/health centre 22</p> <p>Government dispensary 23</p> <p>Other public (<i>specify</i>) 26</p> <p>Private Medical Sector</p> <p>Private hospital 31</p> <p>Private clinic 32</p> <p>Private maternity home 33</p> <p>Mission hospital /clinic 34</p> <p>Other private medical (<i>specify</i>) 36</p> <p>Other (<i>specify</i>) 96</p>	<p>11⇒MN20</p> <p>12⇒MN20</p> <p>96⇒MN20</p>
<p>MN19. WAS (name) DELIVERED BY CAESAREAN SECTION? THAT IS, DID THEY CUT YOUR BELLY OPEN TO TAKE THE BABY OUT?</p>	<p>Yes 1</p> <p>No 2</p>	<p>2⇒MN20</p>
<p>MN19A. WHEN WAS THE DECISION MADE TO HAVE THE CAESAREAN SECTION?</p> <p>WAS IT BEFORE OR AFTER YOUR LABOUR PAINS STARTED?</p>	<p>Before 1</p> <p>After 2</p>	
<p>MN20. WHEN (name) WAS BORN, WAS HE/SHE VERY LARGE, LARGER THAN AVERAGE, AVERAGE, SMALLER THAN AVERAGE, OR VERY SMALL?</p>	<p>Very large 1</p> <p>Larger than average 2</p> <p>Average 3</p> <p>Smaller than average 4</p> <p>Very small 5</p> <p>DK 8</p>	
<p>MN21. WAS (name) WEIGHED AT BIRTH?</p>	<p>Yes 1</p> <p>No 2</p> <p>DK 8</p>	<p>2⇒MN23</p> <p>8⇒MN23</p>
<p>MN22. HOW MUCH DID (name) WEIGH?</p> <p><i>If a card is available, record weight from card.</i></p>	<p>From card 1 (kg)</p> <p>From recall 2 (kg)</p>	

	DK 99998	
MN23. HAS YOUR MENSTRUAL PERIOD RETURNED SINCE THE BIRTH OF (<i>name</i>)?	Yes 1 No 2	
MN24. DID YOU EVER BREASTFEED (<i>name</i>)?	Yes 1 No 2	2⇒Next Module
MN25. HOW LONG AFTER BIRTH DID YOU FIRST PUT (<i>name</i>) TO THE BREAST? <i>If less than 1 hour, record '00' hours. If less than 24 hours, record hours. Otherwise, record days.</i>	Immediately 000 Hours 1 __ __ Days 2 __ __ DK/Don't remember 998	
MN26. IN THE FIRST THREE DAYS AFTER DELIVERY, WAS (<i>name</i>) GIVEN ANYTHING TO DRINK OTHER THAN BREAST MILK?	Yes 1 No 2	2⇒Next Module
MN27. WHAT WAS (<i>name</i>) GIVEN TO DRINK? <i>Probe:</i> ANYTHING ELSE?	Milk (other than breast milk) A Plain water B Sugar or glucose water C Gripe water D Sugar-salt-water solution E Fruit juice F Infant formula G Tea / Infusions H Honey I Other (<i>specify</i>) X	

POST-NATAL HEALTH CHECKS		PN
<p><i>This module is to be administered to all women with a live birth in the 2 years preceding the date of interview. Record name of last-born child from CM13 here _____.</i></p> <p><i>Use this child's name in the following questions, where indicated.</i></p>		
<p>PN1. Check MN18: Was the child delivered in a health facility?</p> <p><input type="checkbox"/> Yes, the child was delivered in a health facility (MN18=21-26 or 31-36) ⇒ Continue with PN2</p> <p><input type="checkbox"/> No, the child was not delivered in a health facility (MN18=11-12 or 96) ⇒ Go to PN6</p>		
<p>PN2. NOW I WOULD LIKE TO ASK YOU SOME QUESTIONS ABOUT WHAT HAPPENED IN THE HOURS AND DAYS AFTER THE BIRTH OF (<i>name</i>).</p> <p>YOU HAVE SAID THAT YOU GAVE BIRTH IN (<i>name or type of facility in MN18</i>). HOW LONG DID YOU STAY THERE AFTER THE DELIVERY?</p> <p><i>If less than one day, record hours.</i> <i>If less than one week, record days.</i> <i>Otherwise, record weeks.</i></p>	<p>Hours..... 1 ___</p> <p>Days 2 ___</p> <p>Weeks 3 ___</p> <p>DK / Don't remember 998</p>	
<p>PN3. I WOULD LIKE TO TALK TO YOU ABOUT CHECKS ON (<i>name</i>)'S HEALTH AFTER DELIVERY – FOR EXAMPLE, SOMEONE EXAMINING (<i>name</i>), CHECKING THE CORD, OR SEEING IF (<i>name</i>) IS OK.</p> <p>BEFORE YOU LEFT THE (<i>name or type of facility in MN18</i>), DID ANYONE CHECK ON (<i>name</i>)'S HEALTH?</p>	<p>Yes 1</p> <p>No..... 2</p>	
<p>PN4. AND WHAT ABOUT CHECKS ON <u>YOUR</u> HEALTH – I MEAN, SOMEONE ASSESSING YOUR HEALTH, FOR EXAMPLE ASKING QUESTIONS ABOUT YOUR HEALTH OR EXAMINING YOU?</p> <p>DID ANYONE CHECK ON <u>YOUR</u> HEALTH BEFORE YOU LEFT (<i>name or type of facility in MN18</i>)?</p>	<p>Yes 1</p> <p>No..... 2</p>	
<p>PN5. NOW I WOULD LIKE TO TALK TO YOU ABOUT WHAT HAPPENED AFTER YOU LEFT (<i>name or type of facility in MN18</i>).</p> <p>DID ANYONE CHECK ON (<i>name</i>)'S HEALTH AFTER YOU LEFT (<i>name or type of facility in MN18</i>)?</p>	<p>Yes 1</p> <p>No..... 2</p>	<p>1⇒PN11</p> <p>2⇒PN16</p>
<p>PN6. Check MN17: Did a health professional, traditional birth attendant, or community health worker assist with the delivery?</p> <p><input type="checkbox"/> Yes, delivery assisted by a health professional, traditional birth attendant, or community health worker (MN17=A-G) ⇒Continue with PN7</p> <p><input type="checkbox"/> No, delivery not assisted by a health professional, traditional birth attendant, or community health worker (A-G not circled in MN17) ⇒ Go to PN10</p>		

<p>PN7. YOU HAVE ALREADY SAID THAT (<i>person or persons in MN17</i>) ASSISTED WITH THE BIRTH. NOW I WOULD LIKE TO TALK TO YOU ABOUT CHECKS ON (<i>name</i>)’S HEALTH AFTER DELIVERY, FOR EXAMPLE EXAMINING (<i>name</i>), CHECKING THE CORD, OR SEEING IF (<i>name</i>) IS OK.</p> <p>AFTER THE DELIVERY WAS OVER AND BEFORE (<i>person or persons in MN17</i>) LEFT YOU, DID (<i>person or persons in MN17</i>) CHECK ON (<i>name</i>)’S HEALTH?</p>	<p>Yes 1 No..... 2</p>	
<p>PN8. AND DID (<i>person or persons in MN17</i>) CHECK ON <u>YOUR</u> HEALTH BEFORE LEAVING?</p> <p>BY CHECK ON YOUR HEALTH, I MEAN ASSESSING YOUR HEALTH, FOR EXAMPLE ASKING QUESTIONS ABOUT YOUR HEALTH OR EXAMINING YOU.</p>	<p>Yes 1 No..... 2</p>	
<p>PN9. AFTER THE (<i>person or persons in MN17</i>) LEFT YOU, DID ANYONE CHECK ON THE HEALTH OF (<i>name</i>)?</p>	<p>Yes 1 No..... 2</p>	<p>1⇒PN11 2⇒PN18</p>
<p>PN10. I WOULD LIKE TO TALK TO YOU ABOUT CHECKS ON (<i>name</i>)’S HEALTH AFTER DELIVERY – FOR EXAMPLE, SOMEONE EXAMINING (<i>name</i>), CHECKING THE CORD, OR SEEING IF THE BABY IS OK.</p> <p>AFTER (<i>name</i>) WAS DELIVERED, DID ANYONE CHECK ON HIS/HER HEALTH?</p>	<p>Yes 1 No..... 2</p>	<p>2⇒PN19</p>
<p>PN11. DID SUCH A CHECK HAPPEN ONLY ONCE, OR MORE THAN ONCE?</p>	<p>Once..... 1 More than once 2</p>	<p>1⇒PN12A 2⇒PN12B</p>
<p>PN12A. HOW LONG AFTER DELIVERY DID THAT CHECK HAPPEN?</p> <p>PN12B. HOW LONG AFTER DELIVERY DID THE FIRST OF THESE CHECKS HAPPEN?</p> <p><i>If less than one day, record hours. If less than one week, record days. Otherwise, record weeks.</i></p>	<p>Hours..... 1 ___</p> <p>Days 2 ___</p> <p>Weeks 3 ___</p> <p>DK / Don’t remember 998</p>	

<p>PN13. WHO CHECKED ON (<i>name</i>)'S HEALTH AT THAT TIME?</p>	<p>Health professional:</p> <p>Doctor A</p> <p>Nurse / Midwife B</p> <p>Clinical Officer C</p> <p>Community Nurse D</p> <p>Other person</p> <p>Traditional birth attendant F</p> <p>Community health worker G</p> <p>Relative / Friend H</p> <p>Other (<i>specify</i>) X</p>	
<p>PN14. WHERE DID THIS CHECK TAKE PLACE?</p> <p><i>Probe to identify the type of source.</i></p> <p><i>If unable to determine whether public or private, write the name of the place.</i></p> <p>_____</p> <p>(<i>Name of place</i>)</p>	<p>Home</p> <p>Respondent's home 11</p> <p>Other home 12</p> <p>Public sector</p> <p>Government hospital 21</p> <p>Government clinic / health centre 22</p> <p>Government dispensary 23</p> <p>Other public (<i>specify</i>) 26</p> <p>Private Medical Sector</p> <p>Private hospital 31</p> <p>Private clinic 32</p> <p>Private maternity home 33</p> <p>Mission hospital /clinic 34</p> <p>Other Private Medical 35</p> <p>Other (<i>specify</i>) 96</p>	
<p>PN15. Check MN18: Was the child delivered in a health facility?</p> <p><input type="checkbox"/> Yes, the child was delivered in a health facility (MN18=21-26 or 31-36) ⇒ Continue with PN16</p> <p><input type="checkbox"/> No, the child was not delivered in a health facility (MN18=11-12 or 96) ⇒ Go to PN17</p>		
<p>PN16. AFTER YOU LEFT (<i>name or type of facility in MN18</i>), DID ANYONE CHECK ON <u>YOUR</u> HEALTH?</p>	<p>Yes 1</p> <p>No 2</p>	<p>1 ⇒ PN20</p> <p>2 ⇒ Next Module</p>
<p>PN17. Check MN17: Did a health professional, traditional birth attendant, or community health worker assist with the delivery?</p> <p><input type="checkbox"/> Yes, delivery assisted by a health professional, traditional birth attendant, or community health worker (MN17=A-G) ⇒ Continue with PN18</p> <p><input type="checkbox"/> No, delivery not assisted by a health professional, traditional birth attendant, or community health worker (A-G not circled in MN17) ⇒ Go to PN19</p>		
<p>PN18. AFTER THE DELIVERY WAS OVER AND (<i>person or persons in MN17</i>) LEFT, DID ANYONE CHECK ON <u>YOUR</u> HEALTH?</p>	<p>Yes 1</p> <p>No 2</p>	<p>1 ⇒ PN20</p> <p>2 ⇒ Next Module</p>

<p>PN19. AFTER THE BIRTH OF (<i>name</i>), DID ANYONE CHECK ON <u>YOUR</u> HEALTH?</p> <p>I MEAN SOMEONE ASSESSING YOUR HEALTH, FOR EXAMPLE ASKING QUESTIONS ABOUT YOUR HEALTH OR EXAMINING YOU.</p>	<p>Yes 1 No..... 2</p>	<p>2⇒Next Module</p>
<p>PN20. DID SUCH A CHECK HAPPEN ONLY ONCE, OR MORE THAN ONCE?</p>	<p>Once..... 1 More than once 2</p>	<p>1⇒PN21A 2⇒PN21B</p>
<p>PN21A. HOW LONG AFTER DELIVERY DID THAT CHECK HAPPEN?</p> <p>PN21B. HOW LONG AFTER DELIVERY DID THE FIRST OF THESE CHECKS HAPPEN?</p> <p><i>If less than one day, record hours. If less than one week, record days. Otherwise, record weeks.</i></p>	<p>Hours..... 1 ___ ___ Days 2 ___ ___ Weeks 3 ___ ___ DK / Don't remember 998</p>	
<p>PN22. WHO CHECKED ON <u>YOUR</u> HEALTH AT THAT TIME?</p>	<p>Health professional: Doctor.....A Nurse / MidwifeB Clinical OfficerC Community NurseD</p> <p>Other person Traditional birth attendant F Community health worker G Relative / FriendH</p> <p>Other (<i>specify</i>)X</p>	
<p>PN23. WHERE DID THIS CHECK TAKE PLACE?</p> <p><i>Probe to identify the type of source.</i></p> <p><i>If unable to determine whether public or private, write the name of the place.</i></p> <p>_____</p> <p>(<i>Name of place</i>)</p>	<p>Home Respondent's home 11 Other home 12</p> <p>Public sector Government hospital 21 Government clinic / health centre 22 Government dispensary 23 Other public (<i>specify</i>) 26</p> <p>Private Medical Sector Private hospital..... 31 Private clinic 32 Private maternity home 33 Mission hospital /clinic..... 34 Other Private Medical 35</p> <p>Other (<i>specify</i>) 96</p>	

ILLNESS SYMPTOMS		IS
<p>IS1. Check List of Household Members, columns HL7B and HL15</p> <p>Is the respondent the mother or caretaker of any child under age 5?</p> <p><input type="checkbox"/> Yes ⇒ Continue with IS2.</p> <p><input type="checkbox"/> No ⇒ Go to Next Module.</p>		
<p>IS2. SOMETIMES CHILDREN HAVE SEVERE ILLNESSES AND SHOULD BE TAKEN IMMEDIATELY TO A HEALTH FACILITY. WHAT TYPES OF SYMPTOMS WOULD CAUSE YOU TO TAKE A CHILD UNDER THE AGE OF 5 TO A HEALTH FACILITY RIGHT AWAY?</p> <p><i>Probe:</i> ANY OTHER SYMPTOMS?</p> <p><i>Keep asking for more signs or symptoms until the mother/caretaker cannot recall any additional symptoms.</i></p> <p><i>Circle all symptoms mentioned, but do <u>not</u> prompt with any suggestions</i></p>	<p>Child not able to drink or breastfeed A</p> <p>Child becomes sicker B</p> <p>Child develops a fever C</p> <p>Child has fast breathing D</p> <p>Child has difficulty breathing E</p> <p>Child has blood in stool F</p> <p>Child is drinking poorly G</p> <p>Other (<i>specify</i>) _____ X</p> <p>Other (<i>specify</i>) _____ Y</p> <p>Other (<i>specify</i>) _____ Z</p>	

CONTRACEPTION		CP
<p>CP1. I WOULD LIKE TO TALK WITH YOU ABOUT ANOTHER SUBJECT – FAMILY PLANNING.</p> <p>ARE YOU PREGNANT NOW?</p>	<p>Yes, currently pregnant..... 1</p> <p>No..... 2</p> <p>Unsure or DK 8</p>	1⇒CP2A
<p>CP2. COUPLES USE VARIOUS WAYS OR METHODS TO DELAY OR AVOID A PREGNANCY.</p> <p>ARE YOU CURRENTLY DOING SOMETHING OR USING ANY METHOD TO DELAY OR AVOID GETTING PREGNANT?</p>	<p>Yes 1</p> <p>No..... 2</p>	1⇒CP3
<p>CP2A. HAVE YOU EVER DONE SOMETHING OR USED ANY METHOD TO DELAY OR AVOID GETTING PREGNANT?</p>	<p>Yes 1</p> <p>No..... 2</p>	1⇒Next Module 2⇒Next Module
<p>CP3. WHAT ARE YOU DOING TO DELAY OR AVOID A PREGNANCY?</p> <p><i>Do not prompt.</i> <i>If more than one method is mentioned, circle each one.</i></p>	<p>Female sterilizationA</p> <p>Male sterilizationB</p> <p>IUDC</p> <p>InjectablesD</p> <p>Implants.....E</p> <p>PillF</p> <p>Male condom.....G</p> <p>Female condomH</p> <p>Diaphragm.....I</p> <p>Foam/ Jelly.....J</p> <p>Lactational amenorrhoea method (LAM)K</p> <p>Periodic abstinence/RhythmL</p> <p>WithdrawalM</p> <p>Other (<i>specify</i>).....X</p>	

UNMET NEED		UN
UN1. Check CP1. Currently pregnant? <input type="checkbox"/> Yes, currently pregnant ⇒ Continue with UN2 <input type="checkbox"/> No, unsure or DK ⇒ Go to UN5		
UN2. NOW I WOULD LIKE TO TALK TO YOU ABOUT YOUR CURRENT PREGNANCY. WHEN YOU GOT PREGNANT, DID YOU WANT TO GET PREGNANT AT THAT TIME?	Yes 1 No 2	1⇒UN4
UN3. DID YOU WANT TO HAVE A BABY LATER ON OR DID YOU NOT WANT ANY (MORE) CHILDREN?	Later 1 No more 2	
UN4. NOW I WOULD LIKE TO ASK SOME QUESTIONS ABOUT THE FUTURE. AFTER THE CHILD YOU ARE NOW EXPECTING, WOULD YOU LIKE TO HAVE ANOTHER CHILD, OR WOULD YOU PREFER NOT TO HAVE ANY MORE CHILDREN?	Have another child 1 No more / None 2 Undecided / DK 8	1⇒UN7 2⇒UN13 8⇒UN13
UN5. Check CP3. Currently using "Female sterilization"? <input type="checkbox"/> Yes ⇒ Go to UN13 <input type="checkbox"/> No ⇒ Continue with UN6		
UN6. NOW I WOULD LIKE TO ASK YOU SOME QUESTIONS ABOUT THE FUTURE. WOULD YOU LIKE TO HAVE (A/ANOTHER) CHILD, OR WOULD YOU PREFER NOT TO HAVE ANY (MORE) CHILDREN?	Have (a/another) child 1 No more / None 2 Says she cannot get pregnant 3 Undecided / DK 8	2⇒UN9 3⇒UN11 8⇒UN9
UN7. HOW LONG WOULD YOU LIKE TO WAIT BEFORE THE BIRTH OF (A/ANOTHER) CHILD? <i>Record the answer as stated by respondent.</i>	Months 1 ___ ___ Years 2 ___ ___ Does not want to wait (soon/how) 993 Says she cannot get pregnant 994 After marriage 995 Other 996 DK 998	994⇒UN11
UN8. Check CP1. Currently pregnant? <input type="checkbox"/> Yes, currently pregnant ⇒ Go to UN13 <input type="checkbox"/> No, unsure or DK ⇒ Continue with UN9		

UN9. Check CP2. Currently using a method? <input type="checkbox"/> Yes ⇒ Go to UN13 <input type="checkbox"/> No ⇒ Continue with UN10		
UN10. DO YOU THINK YOU ARE PHYSICALLY ABLE TO GET PREGNANT AT THIS TIME?	Yes 1 No 2 DK 8	1 ⇒ UN13 8 ⇒ UN13
UN11. WHY DO YOU THINK YOU ARE NOT PHYSICALLY ABLE TO GET PREGNANT?	Infrequent sex / No sex A Menopausal B Never menstruated C Hysterectomy (surgical removal of uterus) D Has been trying to get pregnant for 2 years or more without result E Postpartum amenorrhic F Breastfeeding G Too old H Fatalistic I Other (<i>specify</i>) X DK Z	
UN12. Check UN11. "Never menstruated" mentioned? <input type="checkbox"/> Mentioned ⇒ Go to Next Module <input type="checkbox"/> Not mentioned ⇒ Continue with UN13		
UN13. WHEN DID YOUR LAST MENSTRUAL PERIOD START? <i>Record the answer using the same unit stated by the respondent</i>	Days ago 1 ___ Weeks ago 2 ___ Months ago 3 ___ Years ago 4 ___ In menopause / Has had hysterectomy 994 Before last birth 995 Never menstruated 996	

FEMALE GENITAL MUTILATION/CUTTING		FG
FG1. HAVE YOU EVER HEARD OF FEMALE CIRCUMCISION?	Yes 1 No 2	1⇒FG3
FG2. IN SOME COUNTRIES, THERE IS A PRACTICE IN WHICH A GIRL MAY HAVE PART OF HER GENITALS CUT. HAVE YOU EVER HEARD ABOUT THIS PRACTICE?	Yes 1 No 2	2⇒Next Module
FG3. HAVE YOU YOURSELF EVER BEEN CIRCUMCISED?	Yes 1 No 2	2⇒FG9
FG4. NOW I WOULD LIKE TO ASK YOU WHAT WAS DONE TO YOU AT THAT TIME. WAS ANY FLESH REMOVED FROM THE GENITAL AREA?	Yes 1 No 2 DK 8	1⇒FG6
FG5. WAS THE GENITAL AREA JUST NICKED WITHOUT REMOVING ANY FLESH?	Yes 1 No 2 DK 8	
FG6. WAS THE GENITAL AREA SOWN CLOSED? <i>If necessary, probe: WAS IT SEALED?</i>	Yes 1 No 2 DK 8	
FG7. HOW OLD WERE YOU WHEN YOU WERE CIRCUMCISED? <i>If the respondent does not know the exact age, probe to get an estimate</i>	Age at circumcision __ __ DK/Don't remember/Not sure 98	
FG8. WHO PERFORMED THE CIRCUMCISION?	Health professional Doctor 11 Nurse/Midwife 12 Other health professional (<i>specify</i>) 16 Traditional persons Traditional 'circumciser' 21 Traditional birth attendant 22 Other traditional (<i>specify</i>) 26 DK 98	
FG9. <i>Check CM5 for Number of daughters at home and CM7 for Number of daughters elsewhere, and sum the answers here</i>	Total number of living daughters __ __	
FG10. JUST TO MAKE SURE THAT I HAVE THIS RIGHT, YOU HAVE (<i>total number in FG9</i>) LIVING DAUGHTERS. IS THIS CORRECT?		
<input type="checkbox"/> Yes		
<input type="checkbox"/> One or more living daughters ⇒ Continue with FG11		
<input type="checkbox"/> Does not have any living daughters ⇒ Go to FG22		
<input type="checkbox"/> No ⇒ Check responses to CM1 – CM10 and make corrections as necessary, until FG10 = Yes		

FG11. Ask the respondent to tell you the name(s) of her daughter(s), beginning with the youngest daughter (if more than one daughter). Write down the name of each daughter in FG12. Then, ask questions FG13 to FG20 for each daughter at a time.

The total number of daughters in FG12 should be equal to the number in FG9

If more than 4 daughters, use additional questionnaires

	Daughter #1	Daughter #2	Daughter #3	Daughter #4
FG12. Name of daughter	_____	_____	_____	_____
FG13. HOW OLD IS (name)?	Age ____	Age ____	Age ____	Age ____
FG14. Is (name) younger than 15 years of age?	Yes1 No2 <i>If "No", go to FG13 for next daughter. If no more daughters, go to FG22</i>	Yes1 No2 <i>If "No", go to FG13 for next daughter. If no more daughters, go to FG22</i>	Yes1 No2 <i>If "No", go to FG13 for next daughter. If no more daughters, go to FG22</i>	Yes1 No2 <i>If "No", go to FG13 for next daughter. If no more daughters, go to FG22</i>
FG15. IS (name) CIRCUMCISED?	Yes1 No2 <i>If "No", go to FG13 for next daughter. If no more daughters, go to FG22</i>	Yes1 No2 <i>If "No", go to FG13 for next daughter. If no more daughters, go to FG22</i>	Yes1 No2 <i>If "No", go to FG13 for next daughter. If no more daughters, go to FG22</i>	Yes1 No2 <i>If "No", go to FG13 for next daughter. If no more daughters, go to FG22</i>
FG16. HOW OLD WAS (name) WHEN THIS OCCURRED? <i>If the respondent does not know the age, probe to get an estimate.</i>	Age ____ DK.....98	Age ____ DK.....98	Age ____ DK98	Age ____ DK98
FG17. NOW I WOULD LIKE TO ASK YOU WHAT WAS DONE TO (name) AT THAT TIME. WAS ANY FLESH REMOVED FROM THE GENITAL AREA?	Yes1 ⇒FG19 No2 DK.....8	Yes1 ⇒FG19 No2 DK.....8	Yes1 ⇒FG19 No2 DK8	Yes1 ⇒FG19 No2 DK8
FG18. WAS HER GENITAL AREA JUST NICKED WITHOUT REMOVING ANY FLESH?	Yes1 No2 DK.....8	Yes1 No2 DK.....8	Yes1 No2 DK8	Yes1 No2 DK8

FG19. WAS HER GENITAL AREA SEWN CLOSED? <i>If necessary, probe:</i> WAS IT SEALED?	Yes1 No2 DK.....8	Yes1 No2 DK.....8	Yes1 No2 DK8	Yes1 No2 DK8
FG20. WHO PERFORMED THE CIRCUMCISION?	Health professional Doctor11 Nurse/midwife...12 Other health professional (specify) _____16 Traditional persons Traditional 'circumciser'21 Traditional birth attendant.....22 Other traditional (specify) _____26 DK.....98	Health professional Doctor11 Nurse/midwife...12 Other health professional (specify) _____16 Traditional persons Traditional 'circumciser'21 Traditional birth attendant.....22 Other traditional (specify) _____26 DK.....98	Health professional Doctor11 Nurse/midwife...12 Other health professional (specify) _____16 Traditional persons Traditional 'circumciser'21 Traditional birth attendant.....22 Other traditional (specify) _____26 DK98	Health professional Doctor11 Nurse/midwife...12 Other health professional (specify) _____16 Traditional persons Traditional 'circumciser'21 Traditional birth attendant.....22 Other traditional (specify) _____26 DK98
FG21.	<i>Go back to FG13 for next daughter. If no more daughters, continue with FG22</i>	<i>Go back to FG13 for next daughter. If no more daughters, continue with FG22</i>	<i>Go back to FG13 for next daughter. If no more daughters, continue with FG22</i>	<i>Go back to FG13 in first column of additional questionnaire for next daughter. If no more daughters, continue with FG22</i>
				<i>Tick here if additional questionnaire used</i> <input type="checkbox"/>

FG22. DO YOU THINK THIS PRACTICE SHOULD BE CONTINUED OR SHOULD IT BE DISCONTINUED?	Continued.....1 Discontinued2 Depends.....3 DK8	
---	--	--

ATTITUDES TOWARD DOMESTIC VIOLENCE		DV		
DV1. SOMETIMES A HUSBAND IS ANNOYED OR ANGERED BY THINGS THAT HIS WIFE DOES. IN YOUR OPINION, IS A HUSBAND JUSTIFIED IN HITTING OR BEATING HIS WIFE IN THE FOLLOWING SITUATIONS:				
		Yes	No	DK
[A] IF SHE GOES OUT WITHOUT TELLING HIM?	Goes out without telling	1	2	8
[B] IF SHE NEGLECTS THE CHILDREN?	Neglects children	1	2	8
[C] IF SHE ARGUES WITH HIM?	Argues with him	1	2	8
[D] IF SHE REFUSES TO HAVE SEX WITH HIM?	Refuses sex.....	1	2	8
[E] IF SHE BURNS THE FOOD?	Burns food	1	2	8

MARRIAGE/UNION		MA
MA1. ARE YOU CURRENTLY MARRIED OR LIVING TOGETHER WITH A MAN AS IF MARRIED?	Yes, currently married1 Yes, living with a man.....2 No, not in union3	3⇒MA5
MA2. HOW OLD IS YOUR HUSBAND/PARTNER? <i>Probe:</i> HOW OLD WAS YOUR HUSBAND/PARTNER ON HIS LAST BIRTHDAY?	Age in years..... _ _ DK..... 98	
MA3. BESIDES YOURSELF, DOES YOUR HUSBAND/PARTNER HAVE ANY OTHER WIVES OR PARTNERS OR DOES HE LIVE WITH OTHER WOMEN AS IF MARRIED?	Yes1 No2	2⇒MA7
MA4. HOW MANY OTHER WIVES OR PARTNERS DOES HE HAVE?	Number..... _ _ DK..... 98	⇒MA7 98⇒MA7
MA5. HAVE YOU EVER BEEN MARRIED OR LIVED TOGETHER WITH A MAN AS IF MARRIED?	Yes, formerly married1 Yes, formerly lived with a man2 No3	3 ⇒Next Module
MA6. WHAT IS YOUR MARITAL STATUS NOW: ARE YOU WIDOWED, DIVORCED OR SEPARATED?	Widowed.....1 Divorced2 Separated3	
MA7. HAVE YOU BEEN MARRIED OR LIVED WITH A MAN ONLY ONCE OR MORE THAN ONCE?	Only once1 More than once.....2	1 ⇒MA8A 2 ⇒MA8B
MA8A. IN WHAT MONTH AND YEAR DID YOU MARRY OR START LIVING WITH A MAN AS IF MARRIED?	Date of (first) marriage Month..... _ _ DK month..... 98	
MA8B. IN WHAT MONTH AND YEAR DID YOU <u>FIRST</u> MARRY OR START LIVING WITH A MAN AS IF MARRIED?	Year _ _ _ _ DK year..... 9998	⇒Next Module
MA9. HOW OLD WERE YOU WHEN YOU FIRST STARTED LIVING WITH YOUR (<u>FIRST</u>) HUSBAND/PARTNER?	Age in years..... _ _	

SEXUAL BEHAVIOUR		SB
Check for the presence of others. Before continuing, ensure privacy.		
SB1. NOW I WOULD LIKE TO ASK YOU SOME QUESTIONS ABOUT SEXUAL ACTIVITY IN ORDER TO GAIN A BETTER UNDERSTANDING OF SOME IMPORTANT LIFE ISSUES. THE INFORMATION YOU SUPPLY WILL REMAIN STRICTLY CONFIDENTIAL. HOW OLD WERE YOU WHEN YOU HAD SEXUAL INTERCOURSE FOR THE VERY FIRST TIME?	Never had intercourse 00 Age in years ___ ___ First time when started living with (first) husband/partner 95	00⇒Next Module
SB2. THE FIRST TIME YOU HAD SEXUAL INTERCOURSE, WAS A CONDOM USED?	Yes 1 No 2 DK / Don't remember 8	
SB3. WHEN WAS THE LAST TIME YOU HAD SEXUAL INTERCOURSE? <i>Record answers in days, weeks or months if less than 12 months (one year). If 12 months (one year) or more, answer must be recorded in years.</i>	Days ago 1 ___ ___ Weeks ago 2 ___ ___ Months ago 3 ___ ___ Years ago 4 ___ ___	4⇒SB15
SB4. THE LAST TIME YOU HAD SEXUAL INTERCOURSE, WAS A CONDOM USED?	Yes 1 No 2	
SB5. WHAT WAS YOUR RELATIONSHIP TO THIS PERSON WITH WHOM YOU LAST HAD SEXUAL INTERCOURSE? <i>Probe to ensure that the response refers to the relationship at the time of sexual intercourse</i> <i>If 'boyfriend', then ask: WERE YOU LIVING TOGETHER AS IF MARRIED? If 'yes', circle '2'. If 'no', circle '3'.</i>	Husband 1 Cohabiting partner 2 Boyfriend 3 Casual acquaintance 4 Other (specify) 6	3⇒SB7 4⇒SB7 6⇒SB7
SB6. Check MAI: <input type="checkbox"/> Currently married or living with a man (MAI = 1 or 2) ⇒ Go to SB8 <input type="checkbox"/> Not married / Not in union (MAI = 3) ⇒ Continue with SB7		
SB7. HOW OLD IS THIS PERSON? <i>If response is DK, probe: ABOUT HOW OLD IS THIS PERSON?</i>	Age of sexual partner ___ ___ DK 98	
SB8. HAVE YOU HAD SEXUAL INTERCOURSE WITH ANY OTHER PERSON IN THE LAST 12 MONTHS?	Yes 1 No 2	2⇒SB15
SB9. THE LAST TIME YOU HAD SEXUAL INTERCOURSE WITH THIS OTHER PERSON, WAS A CONDOM USED?	Yes 1 No 2	

<p>SB10. WHAT WAS YOUR RELATIONSHIP TO THIS PERSON?</p> <p><i>Probe to ensure that the response refers to the relationship at the time of sexual intercourse</i></p> <p><i>If 'boyfriend' then ask:</i> WERE YOU LIVING TOGETHER AS IF MARRIED? <i>If 'yes', circle '2'. If 'no', circle '3'.</i></p>	<p>Husband1 Cohabiting partner2 Boyfriend.....3 Casual acquaintance4 Other (<i>specify</i>)6</p>	<p>3⇒SB12 4⇒SB12 6⇒SB12</p>
<p>SB11. Check MA1 and MA7:</p> <p><input type="checkbox"/> <i>Currently married or living with a man (MA1 = 1 or 2) AND Married only once or lived with a man only once (MA7 = 1) ⇒ Go to SB13</i></p> <p><input type="checkbox"/> <i>Else ⇒ Continue with SB12</i></p>		
<p>SB12. HOW OLD IS THIS PERSON?</p> <p><i>If response is DK, probe:</i> ABOUT HOW OLD IS THIS PERSON?</p>	<p>Age of sexual partner..... __ __ DK..... 98</p>	
<p>SB13. OTHER THAN THESE TWO PERSONS, HAVE YOU HAD SEXUAL INTERCOURSE WITH ANY OTHER PERSON IN THE LAST 12 MONTHS?</p>	<p>Yes.....1 No2</p>	<p>2⇒SB15</p>
<p>SB14. IN TOTAL, WITH HOW MANY DIFFERENT PEOPLE HAVE YOU HAD SEXUAL INTERCOURSE IN THE LAST 12 MONTHS?</p>	<p>Number of partners..... __ __</p>	
<p>SB15. IN TOTAL, WITH HOW MANY DIFFERENT PEOPLE HAVE YOU HAD SEXUAL INTERCOURSE IN YOUR LIFETIME?</p> <p><i>If a non-numeric answer is given, probe to get an estimate.</i></p> <p><i>If number of partners is 95 or more, write '95'.</i></p>	<p>Number of lifetime partners __ __ DK..... 98</p>	

HIV/AIDS		HA																
HA1. NOW I WOULD LIKE TO TALK WITH YOU ABOUT SOMETHING ELSE. HAVE YOU EVER HEARD OF AN ILLNESS CALLED AIDS?	Yes 1 No 2 DK 8	2 ⇒ Next Module																
HA2. CAN PEOPLE REDUCE THEIR CHANCE OF GETTING THE AIDS VIRUS BY HAVING JUST ONE UNINFECTED SEX PARTNER WHO HAS NO OTHER SEX PARTNERS?	Yes 1 No 2 DK 8																	
HA3. CAN PEOPLE GET THE AIDS VIRUS BECAUSE OF WITCHCRAFT OR OTHER SUPERNATURAL MEANS?	Yes 1 No 2 DK 8																	
HA4. CAN PEOPLE REDUCE THEIR CHANCE OF GETTING THE AIDS VIRUS BY USING A CONDOM EVERY TIME THEY HAVE SEX?	Yes 1 No 2 DK 8																	
HA5. CAN PEOPLE GET THE AIDS VIRUS FROM MOSQUITO BITES?	Yes 1 No 2 DK 8																	
HA6. CAN PEOPLE GET THE AIDS VIRUS BY SHARING FOOD WITH A PERSON WHO HAS THE AIDS VIRUS?	Yes 1 No 2 DK 8																	
HA7. IS IT POSSIBLE FOR A HEALTHY-LOOKING PERSON TO HAVE THE AIDS VIRUS?	Yes 1 No 2 DK 8																	
HA8. CAN THE VIRUS THAT CAUSES AIDS BE TRANSMITTED FROM A MOTHER TO HER BABY: [A] DURING PREGNANCY? [B] DURING DELIVERY? [C] BY BREASTFEEDING?	<table style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th></th> <th style="text-align: center;">Yes</th> <th style="text-align: center;">No</th> <th style="text-align: center;">DK</th> </tr> </thead> <tbody> <tr> <td>During pregnancy</td> <td style="text-align: center;">1</td> <td style="text-align: center;">2</td> <td style="text-align: center;">8</td> </tr> <tr> <td>During delivery</td> <td style="text-align: center;">1</td> <td style="text-align: center;">2</td> <td style="text-align: center;">8</td> </tr> <tr> <td>By breastfeeding</td> <td style="text-align: center;">1</td> <td style="text-align: center;">2</td> <td style="text-align: center;">8</td> </tr> </tbody> </table>		Yes	No	DK	During pregnancy	1	2	8	During delivery	1	2	8	By breastfeeding	1	2	8	
	Yes	No	DK															
During pregnancy	1	2	8															
During delivery	1	2	8															
By breastfeeding	1	2	8															
HA9. IN YOUR OPINION, IF A FEMALE TEACHER HAS THE AIDS VIRUS BUT IS NOT SICK, SHOULD SHE BE ALLOWED TO CONTINUE TEACHING IN SCHOOL?	Yes 1 No 2 DK/Not sure/Depends 8																	
HA10. WOULD YOU BUY FRESH VEGETABLES FROM A SHOPKEEPER OR VENDOR IF YOU KNEW THAT THIS PERSON HAD THE AIDS VIRUS?	Yes 1 No 2 DK/Not sure/Depends 8																	
HA11. IF A MEMBER OF YOUR FAMILY GOT INFECTED WITH THE AIDS VIRUS, WOULD YOU WANT IT TO REMAIN A SECRET?	Yes 1 No 2 DK/Not sure/Depends 8																	
HA12. IF A MEMBER OF YOUR FAMILY BECAME SICK WITH AIDS, WOULD YOU BE WILLING TO CARE FOR HER OR HIM IN YOUR OWN HOUSEHOLD?	Yes 1 No 2 DK/Not sure/Depends 8																	

HA13. Check CM13: Any live birth in last 2 years? <input type="checkbox"/> No live birth in last 2 years (CM13="No" or blank) ⇒ Go to HA24 <input type="checkbox"/> One or more live births in last 2 years ⇒ Continue with HA14																						
HA14. Check MN1: Received antenatal care? <input type="checkbox"/> Received antenatal care ⇒ Continue with HA15 <input type="checkbox"/> Did not receive antenatal care ⇒ Go to HA24																						
HA15. DURING ANY OF THE ANTENATAL VISITS FOR YOUR PREGNANCY WITH (name), WERE YOU GIVEN ANY INFORMATION ABOUT: [A] BABIES GETTING THE AIDS VIRUS FROM THEIR MOTHER? [B] THINGS THAT YOU CAN DO TO PREVENT GETTING THE AIDS VIRUS? [C] GETTING TESTED FOR THE AIDS VIRUS? WERE YOU: [D] OFFERED A TEST FOR THE AIDS VIRUS?	<table> <thead> <tr> <th></th> <th>Y</th> <th>N</th> <th>DK</th> </tr> </thead> <tbody> <tr> <td>AIDS from mother.....</td> <td>1</td> <td>2</td> <td>8</td> </tr> <tr> <td>Things to do.....</td> <td>1</td> <td>2</td> <td>8</td> </tr> <tr> <td>Tested for AIDS.....</td> <td>1</td> <td>2</td> <td>8</td> </tr> <tr> <td>Offered a test.....</td> <td>1</td> <td>2</td> <td>8</td> </tr> </tbody> </table>		Y	N	DK	AIDS from mother.....	1	2	8	Things to do.....	1	2	8	Tested for AIDS.....	1	2	8	Offered a test.....	1	2	8	
	Y	N	DK																			
AIDS from mother.....	1	2	8																			
Things to do.....	1	2	8																			
Tested for AIDS.....	1	2	8																			
Offered a test.....	1	2	8																			
HA16. I DON'T WANT TO KNOW THE RESULTS, BUT WERE YOU TESTED FOR THE AIDS VIRUS AS PART OF YOUR ANTENATAL CARE?	Yes1 No2 DK.....8	2⇒HA19 8⇒HA19																				
HA17. I DON'T WANT TO KNOW THE RESULTS, BUT DID YOU GET THE RESULTS OF THE TEST?	Yes1 No2 DK.....8	2⇒HA22 8⇒HA22																				
HA18. REGARDLESS OF THE RESULT, ALL WOMEN WHO ARE TESTED ARE SUPPOSED TO RECEIVE COUNSELLING AFTER GETTING THE RESULT. AFTER YOU WERE TESTED, DID YOU RECEIVE COUNSELLING?	Yes1 No2 DK.....8	1⇒HA22 2⇒HA22 8⇒HA22																				
HA19. Check MN17: Birth delivered by health professional (A, B or C)? <input type="checkbox"/> Yes, birth delivered by health professional (MN17 = A, B or C) ⇒ Continue with HA20 <input type="checkbox"/> No, birth not delivered by health professional (MN17 = else) ⇒ Go to HA24																						
HA20. I DON'T WANT TO KNOW THE RESULTS, BUT WERE YOU TESTED FOR THE AIDS VIRUS BETWEEN THE TIME YOU WENT FOR DELIVERY BUT BEFORE THE BABY WAS BORN?	Yes1 No2	2⇒HA24																				
HA21. I DON'T WANT TO KNOW THE RESULTS, BUT DID YOU GET THE RESULTS OF THE TEST?	Yes1 No2																					
HA22. HAVE YOU BEEN TESTED FOR THE AIDS VIRUS SINCE THAT TIME YOU WERE TESTED DURING YOUR PREGNANCY?	Yes1 No2	1⇒HA25																				

HA23. WHEN WAS THE MOST RECENT TIME YOU WERE TESTED FOR THE AIDS VIRUS?	Less than 12 months ago..... 1 12-23 months ago 2 2 or more years ago 3	1 ⇨Next Module 2 ⇨Next Module 3 ⇨Next Module
HA24. I DON'T WANT TO KNOW THE RESULTS, BUT HAVE YOU EVER BEEN TESTED TO SEE IF YOU HAVE THE AIDS VIRUS?	Yes 1 No..... 2	2⇨HA27
HA25. WHEN WAS THE MOST RECENT TIME YOU WERE TESTED?	Less than 12 months ago..... 1 12-23 months ago 2 2 or more years ago 3	
HA26. I DON'T WANT TO KNOW THE RESULTS, BUT DID YOU GET THE RESULTS OF THE TEST?	Yes 1 No..... 2 DK 8	1 ⇨Next Module 2 ⇨Next Module 8 ⇨Next Module
HA27. DO YOU KNOW OF A PLACE WHERE PEOPLE CAN GO TO GET TESTED FOR THE AIDS VIRUS?	Yes 1 No..... 2	

TOBACCO AND ALCOHOL USE		TA
TA1. HAVE YOU EVER TRIED CIGARETTE SMOKING, EVEN ONE OR TWO PUFFS?	Yes 1 No 2	2⇒TA6
TA2. HOW OLD WERE YOU WHEN YOU SMOKED A WHOLE CIGARETTE FOR THE FIRST TIME?	Never smoked a whole cigarette 00 Age ____ ____	00⇒TA6
TA3. DO YOU CURRENTLY SMOKE CIGARETTES?	Yes 1 No 2	2⇒TA6
TA4. IN THE LAST 24 HOURS, HOW MANY CIGARETTES DID YOU SMOKE?	Number of cigarettes ____ ____	
TA5. DURING THE LAST ONE MONTH, ON HOW MANY DAYS DID YOU SMOKE CIGARETTES? <i>If less than 10 days, record the number of days. If 10 days or more but less than a month, circle "10". If "everyday" or "almost every day", circle "30"</i>	Number of days 0 ____ 10 days or more but less than a month 10 Everyday / Almost every day 30	
TA6. HAVE YOU EVER TRIED ANY SMOKED TOBACCO PRODUCTS OTHER THAN CIGARETTES, SUCH AS CIGARS, WATER PIPE, CIGARILLOS OR PIPE?	Yes 1 No 2	2⇒TA10
TA7. DURING THE LAST ONE MONTH, DID YOU USE ANY SMOKED TOBACCO PRODUCTS?	Yes 1 No 2	2⇒TA10
TA8. WHAT TYPE OF SMOKED TOBACCO PRODUCT DID YOU USE OR SMOKE DURING THE LAST ONE MONTH? <i>Circle all mentioned.</i>	Cigars A Water pipe B Cigarillos C Pipe D Other (<i>specify</i>) X	
TA9. DURING THE LAST ONE MONTH, ON HOW MANY DAYS DID YOU USE SMOKED TOBACCO PRODUCTS? <i>If less than 10 days, record the number of days. If 10 days or more but less than a month, circle "10". If "everyday" or "almost every day", circle "30"</i>	Number of days 0 ____ 10 days or more but less than a month 10 Everyday / Almost every day 30	
TA10. HAVE YOU EVER TRIED ANY FORM OF SMOKELESS TOBACCO PRODUCTS, SUCH AS CHEWING TOBACCO, SNUFF, OR DIP?	Yes 1 No 2	2 ⇒TA14
TA11. DURING THE LAST ONE MONTH, DID YOU USE ANY SMOKELESS TOBACCO PRODUCTS?	Yes 1 No 2	2 ⇒TA14

<p>TA12. WHAT TYPE OF SMOKELESS TOBACCO PRODUCT DID YOU USE DURING THE LAST ONE MONTH?</p> <p><i>Circle all mentioned.</i></p>	<p>Chewing tobacco A Snuff B Dip C Other (<i>specify</i>) _____ X</p>	
<p>TA13. DURING THE LAST ONE MONTH, ON HOW MANY DAYS DID YOU USE SMOKELESS TOBACCO PRODUCTS?</p> <p><i>If less than 10 days, record the number of days. If 10 days or more but less than a month, circle "10". If "everyday" or "almost every day", circle "30"</i></p>	<p>Number of days 0 ____ 10 days or more but less than a month 10 Everyday / Almost every day 30</p>	
<p>TA14. NOW I WOULD LIKE TO ASK YOU SOME QUESTIONS ABOUT DRINKING ALCOHOL.</p> <p>HAVE YOU EVER DRUNK ALCOHOL?</p>	<p>Yes 1 No 2</p>	2⇒Next Module
<p>TA15. WE COUNT ONE DRINK OF ALCOHOL AS ONE CAN OR BOTTLE OF BEER, ONE GLASS OF WINE, OR ONE SHOT OF COGNAC, VODKA, WHISKEY, RUM OR CHANG'A</p> <p>HOW OLD WERE YOU WHEN YOU HAD YOUR FIRST DRINK OF ALCOHOL, OTHER THAN A FEW SIPS?</p>	<p>Never had one drink of alcohol 00 Age ____ ____</p>	00⇒Next Module
<p>TA16. DURING THE LAST ONE MONTH, ON HOW MANY DAYS DID YOU HAVE AT LEAST ONE DRINK OF ALCOHOL?</p> <p><i>If respondent did not drink, circle "00". If less than 10 days, record the number of days. If 10 days or more but less than a month, circle "10". If "everyday" or "almost every day", circle "30"</i></p>	<p>Did not have one drink in last one month.. 00 Number of days 0 ____ 10 days or more but less than a month 10 Everyday / Almost every day 30</p>	00⇒Next Module
<p>TA17. IN THE LAST ONE MONTH, ON THE DAYS THAT YOU DRANK ALCOHOL, HOW MANY DRINKS DID YOU USUALLY HAVE PER DAY?</p>	<p>Number of drinks ____ ____</p>	

LIFE SATISFACTION		LS
<p>LS1. Check WB2: Age of respondent is between 15 and 24?</p> <p><input type="checkbox"/> Age 25-49 ⇒ Go to WM11</p> <p><input type="checkbox"/> Age 15-24 ⇒ Continue with LS2</p>		
<p>LS2. I WOULD LIKE TO ASK YOU SOME SIMPLE QUESTIONS ON HAPPINESS AND SATISFACTION.</p> <p>FIRST, TAKING ALL THINGS TOGETHER, WOULD YOU SAY YOU ARE VERY HAPPY, SOMEWHAT HAPPY, NEITHER HAPPY NOR UNHAPPY, SOMEWHAT UNHAPPY OR VERY UNHAPPY?</p> <p>YOU CAN ALSO LOOK AT THESE PICTURES TO HELP YOU WITH YOUR RESPONSE.</p> <p><i>Show side 1 of response card and explain what each symbol represents. Circle the response code selected by the respondent.</i></p>	<p>Very happy 1</p> <p>Somewhat happy 2</p> <p>Neither happy nor unhappy 3</p> <p>Somewhat unhappy 4</p> <p>Very unhappy 5</p>	
<p>LS3. NOW I WILL ASK YOU QUESTIONS ABOUT YOUR LEVEL OF SATISFACTION IN DIFFERENT AREAS.</p> <p>IN EACH CASE, WE HAVE FIVE POSSIBLE RESPONSES: PLEASE TELL ME, FOR EACH QUESTION, WHETHER YOU ARE VERY SATISFIED, SOMEWHAT SATISFIED, NEITHER SATISFIED NOR UNSATISFIED, SOMEWHAT UNSATISFIED OR VERY UNSATISFIED.</p> <p>AGAIN, YOU CAN LOOK AT THESE PICTURES TO HELP YOU WITH YOUR RESPONSE.</p> <p><i>Show side 2 of response card and explain what each symbol represents. Circle the response code selected by the respondent, for questions LS3 to LS13.</i></p> <p>HOW SATISFIED ARE YOU WITH YOUR FAMILY LIFE?</p>	<p>Very satisfied 1</p> <p>Somewhat satisfied 2</p> <p>Neither satisfied nor unsatisfied 3</p> <p>Somewhat unsatisfied 4</p> <p>Very unsatisfied 5</p>	
<p>LS4. HOW SATISFIED ARE YOU WITH YOUR FRIENDSHIPS?</p>	<p>Very satisfied 1</p> <p>Somewhat satisfied 2</p> <p>Neither satisfied nor unsatisfied 3</p> <p>Somewhat unsatisfied 4</p> <p>Very unsatisfied 5</p>	
<p>LS5. DURING THE current 2013/14 SCHOOL YEAR, DID YOU ATTEND SCHOOL AT ANY TIME?</p>	<p>Yes 1</p> <p>No 2</p>	2 ⇒ LS7
<p>LS6. HOW SATISFIED (are/were) YOU WITH YOUR SCHOOL?</p>	<p>Very satisfied 1</p> <p>Somewhat satisfied 2</p> <p>Neither satisfied nor unsatisfied 3</p> <p>Somewhat unsatisfied 4</p> <p>Very unsatisfied 5</p>	

<p>LS7. HOW SATISFIED ARE YOU WITH YOUR CURRENT JOB?</p> <p><i>If the respondent says that she does not have a job, circle "0" and continue with the next question. Do not probe to find out how she feels about not having a job, unless she tells you herself.</i></p>	<p>Does not have a job 0</p> <p>Very satisfied..... 1</p> <p>Somewhat satisfied 2</p> <p>Neither satisfied nor unsatisfied 3</p> <p>Somewhat unsatisfied 4</p> <p>Very unsatisfied..... 5</p>	
<p>LS8. HOW SATISFIED ARE YOU WITH YOUR HEALTH?</p>	<p>Very satisfied..... 1</p> <p>Somewhat satisfied 2</p> <p>Neither satisfied nor unsatisfied 3</p> <p>Somewhat unsatisfied 4</p> <p>Very unsatisfied..... 5</p>	
<p>LS9. HOW SATISFIED ARE YOU WITH WHERE YOU LIVE?</p> <p><i>If necessary, explain that the question refers to the living environment, including the neighbourhood and the dwelling.</i></p>	<p>Very satisfied..... 1</p> <p>Somewhat satisfied 2</p> <p>Neither satisfied nor unsatisfied 3</p> <p>Somewhat unsatisfied 4</p> <p>Very unsatisfied..... 5</p>	
<p>LS10. HOW SATISFIED ARE YOU WITH HOW PEOPLE AROUND YOU GENERALLY TREAT YOU?</p>	<p>Very satisfied..... 1</p> <p>Somewhat satisfied 2</p> <p>Neither satisfied nor unsatisfied 3</p> <p>Somewhat unsatisfied 4</p> <p>Very unsatisfied..... 5</p>	
<p>LS11. HOW SATISFIED ARE YOU WITH THE WAY YOU LOOK?</p>	<p>Very satisfied..... 1</p> <p>Somewhat satisfied 2</p> <p>Neither satisfied nor unsatisfied 3</p> <p>Somewhat unsatisfied 4</p> <p>Very unsatisfied..... 5</p>	
<p>LS12. HOW SATISFIED ARE YOU WITH YOUR LIFE, OVERALL?</p>	<p>Very satisfied..... 1</p> <p>Somewhat satisfied 2</p> <p>Neither satisfied nor unsatisfied 3</p> <p>Somewhat unsatisfied 4</p> <p>Very unsatisfied..... 5</p>	
<p>LS13. HOW SATISFIED ARE YOU WITH YOUR CURRENT INCOME?</p> <p><i>If the respondent says that she does not have any income, circle "0" and continue with the next question. Do not probe to find out how she feels about not having any income, unless she tells you herself.</i></p>	<p>Does not have any income 0</p> <p>Very satisfied..... 1</p> <p>Somewhat satisfied 2</p> <p>Neither satisfied nor unsatisfied 3</p> <p>Somewhat unsatisfied 4</p> <p>Very unsatisfied..... 5</p>	
<p>LS14. COMPARED TO THIS TIME LAST YEAR, WOULD YOU SAY THAT YOUR LIFE HAS IMPROVED, STAYED MORE OR LESS THE SAME, OR WORSENEDED, OVERALL?</p>	<p>Improved 1</p> <p>More or less the same 2</p> <p>Worsened..... 3</p>	
<p>LS15. AND IN ONE YEAR FROM NOW, DO YOU EXPECT THAT YOUR LIFE WILL BE BETTER, WILL BE MORE OR LESS THE SAME, OR WILL BE WORSE, OVERALL?</p>	<p>Better..... 1</p> <p>More or less the same 2</p> <p>Worse..... 3</p>	

WM11. <i>Record the time.</i>	Hour and minutes :	
--------------------------------------	--------------------------------	--

<p>WM12. <i>Check List of Household Members, columns HL7B and HL15.</i> <i>Is the respondent the mother or caretaker of any child age 0-4 living in this household?</i></p> <p><input type="checkbox"/> <i>Yes</i> ⇒ <i>Proceed to complete the result of woman's interview (WM7) on the cover page and then go to QUESTIONNAIRE FOR CHILDREN UNDER FIVE for that child and start the interview with this respondent.</i></p> <p><input type="checkbox"/> <i>No</i> ⇒ <i>End the interview with this respondent by thanking her for her cooperation and proceed to complete the result of the woman's interview (WM7) on the cover page</i></p>






Interviewer's Observations

Field Editor's Observations


Supervisor's Observations

RESPONSE CARD:

SIDE 1

Very happy	Somewhat happy	Neither happy, nor unhappy	Somewhat unhappy	Very unhappy
				

SIDE 2



QUESTIONNAIRE FOR CHILDREN UNDER FIVE WESTERN AND NORTH RIFT SURVEY



UNDER-FIVE CHILD INFORMATION PANEL		UF
<p><i>This questionnaire is to be administered to all mothers or caretakers (see List of Household Members, column HL15) who care for a child that lives with them and is under the age of 5 years (see List of Household Members, column HL7B).</i></p> <p><i>A separate questionnaire should be used for each eligible child.</i></p>		
UF1. Cluster number: <div style="text-align: right;">_____</div>	UF2. Household number: <div style="text-align: right;">_____</div>	
UF3. Child's name: Name _____	UF4. Child's line number: <div style="text-align: right;">_____</div>	
UF5. Mother's/Caretaker's name: Name _____	UF6. Mother's/Caretaker's line number: <div style="text-align: right;">_____</div>	
UF7. Interviewer's name and number: Name _____	UF8. Day/Month/Year of interview: <div style="text-align: right;">_____ / _____ / 201__</div>	

<p><i>Repeat greeting if not already read to this respondent:</i></p> <p>WE ARE FROM THE UNIVERSITY OF NAIROBI AND KENYA NATIONAL BUREAU OF STATISTICS. WE ARE CONDUCTING A SURVEY ABOUT THE SITUATION OF CHILDREN, FAMILIES AND HOUSEHOLDS. I WOULD LIKE TO TALK TO YOU ABOUT <i>(child's name from UF3)</i>'S HEALTH AND WELL-BEING. THE INTERVIEW WILL TAKE ABOUT 20 TO 35 MINUTES. ALL THE INFORMATION WE OBTAIN WILL REMAIN STRICTLY CONFIDENTIAL AND ANONYMOUS.</p>	<p><i>If greeting at the beginning of the household questionnaire has already been read to this person, then read the following:</i></p> <p>NOW I WOULD LIKE TO TALK TO YOU MORE ABOUT <i>(child's name from UF3)</i>'S HEALTH AND OTHER TOPICS. THIS INTERVIEW WILL TAKE ABOUT 20 TO 35 MINUTES. AGAIN, ALL THE INFORMATION WE OBTAIN WILL REMAIN STRICTLY CONFIDENTIAL AND ANONYMOUS.</p>
<p>MAY I START NOW?</p> <p><input type="checkbox"/> <i>Yes, permission is given ⇒ Go to UF12 to record the time and then begin the interview.</i></p> <p><input type="checkbox"/> <i>No, permission is not given ⇒ Circle '03' in UF9. Discuss this result with your supervisor</i></p>	

UF9. Result of interview for children under 5 <i>Codes refer to mother/caretaker.</i>	Completed01 Not at home02 Refused03 Partly completed04 Incapacitated05 Other (<i>specify</i>) _____ 96
---	---

<p>UF10. Field editor's name and number: Name _____</p>	<p>UF11. Main data entry clerk's name and number: Name _____</p>
--	---

<p>UF12. Record the time.</p>	<p>Hour and minutes ____ : ____</p>	
--------------------------------------	--	--

AGE	AG
<p>AG1 NOW I WOULD LIKE TO ASK YOU SOME QUESTIONS ABOUT THE DEVELOPMENT AND HEALTH OF (name). ON WHAT DAY, MONTH AND YEAR WAS (name) BORN? <i>Probe:</i> WHAT IS HIS/HER BIRTHDAY? <i>If the mother/caretaker knows the exact birth date, also enter the day; otherwise, circle 98 for day</i> <i>Month and year must be recorded.</i></p>	<p>Date of birth Day ____ DK day.....98 Month..... ____ Year 20 ____</p>
<p>AG2. HOW OLD IS (name)? <i>Probe:</i> HOW OLD WAS (name) AT HIS/HER LAST BIRTHDAY? <i>Record age in completed years.</i> <i>Record '0' if less than 1 year.</i> <i>Compare and correct AG1 and/or AG2 if inconsistent.</i></p>	<p>Age (in completed years) ____</p>

BIRTH REGISTRATION		BR
BR1. DOES (<i>name</i>) HAVE A BIRTH CERTIFICATE? <i>If yes, ask:</i> MAY I SEE IT?	Yes, seen..... 1 Yes, not seen.....2 No.....3 DK.....8	1⇒Next Module 2⇒Next Module
BR2. HAS (<i>name</i>)'S BIRTH BEEN REGISTERED WITH THE CIVIL AUTHORITIES?	Yes 1 No2 DK.....8	1⇒Next Module
BR3. DO YOU KNOW HOW TO REGISTER (<i>name</i>)'S BIRTH?	Yes 1 No2	

EARLY CHILDHOOD DEVELOPMENT		EC																
<p>EC1. HOW MANY CHILDREN'S BOOKS OR PICTURE BOOKS DO YOU HAVE FOR <i>(name)</i>?</p>	<p>None00</p> <p>Number of children's books.....0 ___</p> <p>Ten or more books 10</p>																	
<p>EC2. I AM INTERESTED IN LEARNING ABOUT THE THINGS THAT <i>(name)</i> PLAYS WITH WHEN HE/SHE IS AT HOME.</p> <p>DOES HE/SHE PLAY WITH:</p> <p>[A] HOMEMADE TOYS (SUCH AS DOLLS, CARS, OR OTHER TOYS MADE AT HOME)?</p> <p>[B] TOYS FROM A SHOP OR MANUFACTURED TOYS?</p> <p>[C] HOUSEHOLD OBJECTS (SUCH AS BOWLS OR POTS) OR OBJECTS FOUND OUTSIDE (SUCH AS STICKS, ROCKS, ANIMAL SHELLS OR LEAVES)?</p> <p><i>If the respondent says "YES" to the categories above, then probe to learn specifically what the child plays with to ascertain the response</i></p>	<table> <thead> <tr> <th></th> <th>Y</th> <th>N</th> <th>DK</th> </tr> </thead> <tbody> <tr> <td>Homemade toys</td> <td>1</td> <td>2</td> <td>8</td> </tr> <tr> <td>Toys from a shop.....</td> <td>1</td> <td>2</td> <td>8</td> </tr> <tr> <td>Household objects or outside objects</td> <td>1</td> <td>2</td> <td>8</td> </tr> </tbody> </table>		Y	N	DK	Homemade toys	1	2	8	Toys from a shop.....	1	2	8	Household objects or outside objects	1	2	8	
	Y	N	DK															
Homemade toys	1	2	8															
Toys from a shop.....	1	2	8															
Household objects or outside objects	1	2	8															
<p>EC3. SOMETIMES ADULTS TAKING CARE OF CHILDREN HAVE TO LEAVE THE HOUSE TO GO SHOPPING, WASH CLOTHES, OR FOR OTHER REASONS AND HAVE TO LEAVE YOUNG CHILDREN.</p> <p>ON HOW MANY DAYS IN THE PAST WEEK WAS <i>(name)</i>:</p> <p>[A] LEFT ALONE FOR MORE THAN AN HOUR?</p> <p>[B] LEFT IN THE CARE OF ANOTHER CHILD, THAT IS, SOMEONE LESS THAN 10 YEARS OLD, FOR MORE THAN AN HOUR?</p> <p><i>If 'none' enter '0'. If 'don't know' enter '8'</i></p>	<p>Number of days left alone for more than an hour</p> <p>Number of days left with other child for more than an hour</p>																	
<p>EC4. Check AG2: Age of child</p> <p><input type="checkbox"/> Child age 0, 1 or 2 ⇒ Go to Next Module</p> <p><input type="checkbox"/> Child age 3 or 4 ⇒ Continue with EC5</p>																		
<p>EC5. DOES <i>(name)</i> ATTEND ANY ORGANIZED LEARNING OR EARLY CHILDHOOD EDUCATION PROGRAMME, SUCH AS A PRIVATE OR GOVERNMENT FACILITY, INCLUDING KINDERGARTEN OR COMMUNITY CHILD CARE?</p>	<p>Yes 1</p> <p>No 2</p> <p>DK..... 8</p>																	

<p>EC7. IN THE PAST 3 DAYS, DID YOU OR ANY HOUSEHOLD MEMBER AGE 15 OR OVER ENGAGE IN ANY OF THE FOLLOWING ACTIVITIES WITH <i>(name)</i>:</p> <p><i>If yes, ask:</i> WHO ENGAGED IN THIS ACTIVITY WITH <i>(name)</i>?</p> <p><i>Circle all that apply.</i></p> <p>[A] READ BOOKS TO OR LOOKED AT PICTURE BOOKS WITH <i>(name)</i>?</p> <p>[B] TOLD STORIES TO <i>(name)</i>?</p> <p>[C] SANG SONGS TO <i>(name)</i> OR WITH <i>(name)</i>, INCLUDING LULLABIES?</p> <p>[D] TOOK <i>(name)</i> OUTSIDE THE HOME, COMPOUND, YARD OR ENCLOSURE?</p> <p>[E] PLAYED WITH <i>(name)</i>?</p> <p>[F] NAMED, COUNTED, OR DREW THINGS TO OR WITH <i>(name)</i>?</p>	<table> <thead> <tr> <th></th> <th>Mother</th> <th>Father</th> <th>Other</th> <th>No one</th> </tr> </thead> <tbody> <tr> <td>Read books</td> <td>A</td> <td>B</td> <td>X</td> <td>Y</td> </tr> <tr> <td>Told stories</td> <td>A</td> <td>B</td> <td>X</td> <td>Y</td> </tr> <tr> <td>Sang songs</td> <td>A</td> <td>B</td> <td>X</td> <td>Y</td> </tr> <tr> <td>Took outside</td> <td>A</td> <td>B</td> <td>X</td> <td>Y</td> </tr> <tr> <td>Played with</td> <td>A</td> <td>B</td> <td>X</td> <td>Y</td> </tr> <tr> <td>Named/counted</td> <td>A</td> <td>B</td> <td>X</td> <td>Y</td> </tr> </tbody> </table>		Mother	Father	Other	No one	Read books	A	B	X	Y	Told stories	A	B	X	Y	Sang songs	A	B	X	Y	Took outside	A	B	X	Y	Played with	A	B	X	Y	Named/counted	A	B	X	Y	
	Mother	Father	Other	No one																																	
Read books	A	B	X	Y																																	
Told stories	A	B	X	Y																																	
Sang songs	A	B	X	Y																																	
Took outside	A	B	X	Y																																	
Played with	A	B	X	Y																																	
Named/counted	A	B	X	Y																																	
<p>EC8. I WOULD LIKE TO ASK YOU SOME QUESTIONS ABOUT THE HEALTH AND DEVELOPMENT OF <i>(name)</i>. CHILDREN DO NOT ALL DEVELOP AND LEARN AT THE SAME RATE. FOR EXAMPLE, SOME WALK EARLIER THAN OTHERS. THESE QUESTIONS ARE RELATED TO SEVERAL ASPECTS OF <i>(name)</i>'S DEVELOPMENT.</p> <p>CAN <i>(name)</i> IDENTIFY OR NAME AT LEAST TEN LETTERS OF THE ALPHABET?</p>	<p>Yes1</p> <p>No2</p> <p>DK.....8</p>																																				
<p>EC9. CAN <i>(name)</i> READ AT LEAST FOUR SIMPLE, POPULAR WORDS?</p>	<p>Yes1</p> <p>No2</p> <p>DK.....8</p>																																				
<p>EC10. DOES <i>(name)</i> KNOW THE NAME AND RECOGNIZE THE SYMBOL OF ALL NUMBERS FROM 1 TO 10?</p>	<p>Yes1</p> <p>No2</p> <p>DK.....8</p>																																				
<p>EC11. CAN <i>(name)</i> PICK UP A SMALL OBJECT WITH TWO FINGERS, LIKE A STICK OR A ROCK FROM THE GROUND?</p>	<p>Yes1</p> <p>No2</p> <p>DK.....8</p>																																				
<p>EC12. IS <i>(name)</i> SOMETIMES TOO SICK TO PLAY?</p>	<p>Yes1</p> <p>No2</p> <p>DK.....8</p>																																				
<p>EC13. DOES <i>(name)</i> FOLLOW SIMPLE DIRECTIONS ON HOW TO DO SOMETHING CORRECTLY?</p>	<p>Yes1</p> <p>No2</p> <p>DK.....8</p>																																				

EC14. WHEN GIVEN SOMETHING TO DO, IS <i>(name)</i> ABLE TO DO IT INDEPENDENTLY?	Yes1 No2 DK.....8	
EC15. DOES <i>(name)</i> GET ALONG WELL WITH OTHER CHILDREN?	Yes1 No2 DK.....8	
EC16. DOES <i>(name)</i> KICK, BITE, OR HIT OTHER CHILDREN OR ADULTS?	Yes1 No2 DK.....8	
EC17. DOES <i>(name)</i> GET DISTRACTED EASILY?	Yes1 No2 DK.....8	

IMMUNIZATION										IM
<i>If an immunization (child health) card is available, copy the dates in IM3 for each type of immunization and Vitamin A recorded on the card. IM6-IM17 will only be asked when a card is not available.</i>										
IM1. DO YOU HAVE A CARD WHERE (name)'S VACCINATIONS ARE WRITTEN DOWN? <i>If yes: MAY I SEE IT PLEASE?</i>				Yes, seen..... 1 Yes, not seen..... 2 No card 3				1⇒IM3 2⇒IM6		
IM2. DID YOU EVER HAVE A VACCINATION CARD FOR (name)?				Yes..... 1 No 2				1⇒IM6 2⇒IM6		
IM3. (a) Copy dates for each vaccination from the card. (b) Write '44' in day column if card shows that vaccination was given but no date recorded.				Date of Immunization						
				Day		Month		Year		
BCG	BCG									
POLIO AT BIRTH	OPV0									
POLIO 1	OPV1									
POLIO 2	OPV2									
POLIO 3	OPV3									
DPT 1	DPT1									
DPT 2	DPT2									
DPT 3	DPT3									
HEPB AT BIRTH	HEP0									
HEPB 1	HEP1									
HEPB 2	HEP2									
HEPB 3	HEP3									
HIB 1	HIB1									
HIB 2	HIB2									
HIB 3	HIB3									
MEASLES (OR MMR OR MR)	MEASLES									
YELLOW FEVER	YF									
VITAMIN A (FIRST DOSE)	VITA1									
VITAMIN A (SECOND DOSE)	VITA2									
IM4. Check IM3. Are all vaccines (BCG to Yellow Fever) recorded? <input type="checkbox"/> Yes ⇒Go to IM19 <input type="checkbox"/> No ⇒Continue with IM5										

<p>IM5. IN ADDITION TO WHAT IS RECORDED ON THIS CARD, DID (<i>name</i>) RECEIVE ANY OTHER VACCINATIONS – INCLUDING VACCINATIONS RECEIVED IN CAMPAIGNS OR IMMUNIZATION DAYS OR CHILD HEALTH DAYS?</p> <p><input type="checkbox"/> <i>Yes</i> ⇒ Go back to IM3 and probe for these vaccinations and write ‘66’ in the corresponding day column for each vaccine mentioned. When finished, skip to IM19</p> <p><input type="checkbox"/> <i>No/DK</i> ⇒ Go to IM19</p>		
<p>IM6. HAS (<i>name</i>) EVER RECEIVED ANY VACCINATIONS TO PREVENT HIM/HER FROM GETTING DISEASES, INCLUDING VACCINATIONS RECEIVED IN A CAMPAIGN OR IMMUNIZATION DAY OR CHILD HEALTH DAY?</p>	<p>Yes..... 1</p> <p>No 2</p> <p>DK..... 8</p>	<p>2⇒IM19</p> <p>8⇒IM19</p>
<p>IM7. HAS (<i>name</i>) EVER RECEIVED A BCG VACCINATION AGAINST TUBERCULOSIS – THAT IS, AN INJECTION IN THE ARM OR SHOULDER THAT USUALLY CAUSES A SCAR?</p>	<p>Yes..... 1</p> <p>No 2</p> <p>DK..... 8</p>	
<p>IM8. HAS (<i>name</i>) EVER RECEIVED ANY VACCINATION DROPS IN THE MOUTH TO PROTECT HIM/HER FROM POLIO?</p>	<p>Yes..... 1</p> <p>No 2</p> <p>DK..... 8</p>	<p>2⇒IM11</p> <p>8⇒IM11</p>
<p>IM9. WAS THE FIRST POLIO VACCINE RECEIVED IN THE FIRST TWO WEEKS AFTER BIRTH?</p>	<p>Yes..... 1</p> <p>No 2</p>	
<p>IM10. HOW MANY TIMES WAS THE POLIO VACCINE RECEIVED?</p>	<p>Number of times _</p>	
<p>IM11. HAS (<i>name</i>) EVER RECEIVED A DPT VACCINATION – THAT IS, AN INJECTION IN THE THIGH TO PREVENT HIM/HER FROM GETTING TETANUS, WHOOPING COUGH, OR DIPHTHERIA?</p> <p><i>Probe by indicating that DPT vaccination is sometimes given at the same time as Polio</i></p>	<p>Yes..... 1</p> <p>No 2</p> <p>DK..... 8</p>	<p>2⇒IM13</p> <p>8⇒IM13</p>
<p>IM12. HOW MANY TIMES WAS THE DPT VACCINE RECEIVED?</p>	<p>Number of times _</p>	
<p>IM13. HAS (<i>name</i>) EVER RECEIVED A HEPATITIS B VACCINATION – THAT IS, AN INJECTION IN THE THIGH TO PREVENT HIM/HER FROM GETTING HEPATITIS B?</p> <p><i>Probe by indicating that the Hepatitis B vaccine is sometimes given at the same time as Polio and DPT vaccines</i></p>	<p>Yes..... 1</p> <p>No 2</p> <p>DK..... 8</p>	<p>2⇒IM15A</p> <p>8⇒IM15A</p>
<p>IM14. WAS THE FIRST HEPATITIS B VACCINE RECEIVED WITHIN 24 HOURS AFTER BIRTH?</p>	<p>Yes..... 1</p> <p>No 2</p> <p>DK..... 8</p>	
<p>IM15. HOW MANY TIMES WAS THE HEPATITIS B RECEIVED?</p>	<p>Number of times _</p>	
<p>IM15A. HAS (<i>name</i>) EVER RECEIVED A HIB VACCINATION – THAT IS, AN INJECTION IN THE THIGH TO PREVENT HIM/HER FROM GETTING HAEMOPHILUS INFLUENZAE TYPE B?</p> <p><i>Probe by indicating that the Hib vaccine is sometimes given at the same time as Polio and DPT vaccines</i></p>	<p>Yes..... 1</p> <p>No 2</p> <p>DK..... 8</p>	<p>2⇒IM16</p> <p>8⇒IM16</p>

IM15B. HOW MANY TIMES WAS THE HIB VACCINE RECEIVED?	Number of times _																					
IM16. HAS (<i>name</i>) EVER RECEIVED A MEASLES INJECTION (OR AN MMR OR MR) – THAT IS, A SHOT IN THE ARM AT THE AGE OF 9 MONTHS OR OLDER - TO PREVENT HIM/HER FROM GETTING MEASLES?	Yes..... 1 No 2 DK..... 8																					
IM17. HAS (<i>name</i>) EVER RECEIVED THE YELLOW FEVER VACCINATION – THAT IS, A SHOT IN THE ARM AT THE AGE OF 9MONTHS OR OLDER - TO PREVENT HIM/HER FROM GETTING YELLOW FEVER? <i>Probe by indicating that the Yellow Fever vaccine is sometimes given at the same time as the measles vaccine</i>	Yes..... 1 No 2 DK..... 8																					
IM19. PLEASE TELL ME IF (NAME) HAS PARTICIPATED IN ANY OF THE FOLLOWING CAMPAIGNS, NATIONAL IMMUNIZATION DAYS AND/OR VITAMIN A OR CHILD HEALTH DAYS: [A] MALEZI BORA AND MEASLES IMMUNIZATION CAMPAIGNS FROM NOVEMBER 2012 [B] MALEZI BORA AND MEASLES IMMUNIZATION CAMPAIGNS FROM MAY 2013 [C] POLIO CAMPAIGN JULY 2013 [D] POLIO CAMPAIGN AUGUST 2013	<table style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 80%;"></th> <th style="width: 10%; text-align: center;">Y</th> <th style="width: 10%; text-align: center;">N</th> <th style="width: 10%; text-align: center;">DK</th> </tr> </thead> <tbody> <tr> <td>Malezi bora, November 2012</td> <td style="text-align: center;">1</td> <td style="text-align: center;">2</td> <td style="text-align: center;">8</td> </tr> <tr> <td>Malezi bora, May 2013</td> <td style="text-align: center;">1</td> <td style="text-align: center;">2</td> <td style="text-align: center;">8</td> </tr> <tr> <td>Polio campaign, July 2013.....</td> <td style="text-align: center;">1</td> <td style="text-align: center;">2</td> <td style="text-align: center;">8</td> </tr> <tr> <td>Polio campaign, August 2013.....</td> <td style="text-align: center;">1</td> <td style="text-align: center;">2</td> <td style="text-align: center;">8</td> </tr> </tbody> </table>		Y	N	DK	Malezi bora, November 2012	1	2	8	Malezi bora, May 2013	1	2	8	Polio campaign, July 2013.....	1	2	8	Polio campaign, August 2013.....	1	2	8	
	Y	N	DK																			
Malezi bora, November 2012	1	2	8																			
Malezi bora, May 2013	1	2	8																			
Polio campaign, July 2013.....	1	2	8																			
Polio campaign, August 2013.....	1	2	8																			
IM20. <i>Is the vaccination card of the child kept at the health facility?</i> <input type="checkbox"/> <i>Yes</i> ⇒ Issue a QUESTIONNAIRE FORM FOR VACCINATION RECORDS AT HEALTH FACILITY for this child. Complete the Information Panel on that Questionnaire and go to Next Module. <input type="checkbox"/> <i>No</i> ⇒ Continue with Next Module																						

BREASTFEEDING AND DIETARY INTAKE		BD
BD1. Check AG2: Age of child <input type="checkbox"/> Child age 0, 1 or 2 ⇒ Continue with BD2 <input type="checkbox"/> Child age 3 or 4 ⇒ Go to CARE OF ILLNESS Module		
BD2. HAS (name) EVER BEEN BREASTFED?	Yes 1 No 2 DK 8	2⇒BD4 8⇒BD4
BD3. IS (name) STILL BEING BREASTFED?	Yes 1 No 2 DK 8	
BD4. YESTERDAY, DURING THE DAY OR NIGHT, DID (name) DRINK ANYTHING FROM A BOTTLE WITH A NIPPLE?	Yes 1 No 2 DK 8	
BD5. DID (name) DRINK ORS (ORAL REHYDRATION SOLUTION) YESTERDAY, DURING THE DAY OR NIGHT?	Yes 1 No 2 DK 8	
BD6. DID (name) DRINK OR EAT VITAMIN OR MINERAL SUPPLEMENTS OR ANY MEDICINES YESTERDAY, DURING THE DAY OR NIGHT?	Yes 1 No 2 DK 8	
BD7. NOW I WOULD LIKE TO ASK YOU ABOUT (OTHER) LIQUIDS THAT (name) MAY HAVE HAD YESTERDAY DURING THE DAY OR THE NIGHT. I AM INTERESTED TO KNOW WHETHER (name) HAD THE ITEM EVEN IF COMBINED WITH OTHER FOODS. PLEASE INCLUDE LIQUIDS CONSUMED OUTSIDE OF YOUR HOME. DID (name) DRINK (Name of item) YESTERDAY DURING THE DAY OR THE NIGHT:	Yes No DK	
[A] PLAIN WATER?	Plain water	1 2 8
[B] JUICE OR JUICE DRINKS?	Juice or juice drinks	1 2 8
[C] SOUP?	Soup	1 2 8
[D] MILK SUCH AS TINNED, POWDERED, OR FRESH ANIMAL MILK?	Milk	1 2 8
<i>If yes: HOW MANY TIMES DID (name) DRINK MILK? If 7 or more times, record '7'. If unknown, record '8'.</i>	Number of times drank milk.....	___
[E] INFANT FORMULA?	Infant formula	1 2 8
<i>If yes: HOW MANY TIMES DID (name) DRINK INFANT FORMULA? If 7 or more times, record '7'. If unknown, record '8'.</i>	Number of times drank infant formula.....	___
[F] ANY OTHER LIQUIDS?	(Specify) _____	1 2 8

<p>BD8. NOW I WOULD LIKE TO ASK YOU ABOUT (OTHER) FOODS THAT (<i>name</i>) MAY HAVE HAD YESTERDAY DURING THE DAY OR THE NIGHT. AGAIN, I AM INTERESTED TO KNOW WHETHER (<i>name</i>) HAD THE ITEM EVEN IF COMBINED WITH OTHER FOODS.</p> <p>PLEASE INCLUDE FOODS CONSUMED OUTSIDE OF YOUR HOME.</p> <p>DID (<i>name</i>) EAT (<i>Name of food</i>) YESTERDAY DURING THE DAY OR THE NIGHT:</p>				
		Yes	No	DK
[A] YOGURT?	Yogurt	1	2	8
<p><i>If yes: HOW MANY TIMES DID (<i>name</i>) DRINK OR EAT YOGURT? If 7 or more times, record '7'. If unknown, record '8'.</i></p>		Number of times drank/ate yogurt __		
[B] ANY FORTIFIED BABY FOOD E.G. CERELAC?	Cerelac	1	2	8
[C] BREAD, RICE, NOODLES, PORRIDGE, OR OTHER FOODS MADE FROM GRAINS?	Foods made from grains	1	2	8
[D] PUMPKIN, CARROTS, SQUASH OR SWEET POTATOES THAT ARE YELLOW OR ORANGE INSIDE?	Pumpkin, carrots, squash, etc.	1	2	8
[E] WHITE POTATOES, WHITE YAMS, MANIOC, CASSAVA, OR ANY OTHER FOODS MADE FROM ROOTS?	White potatoes, white yams, manioc, cassava, etc.	1	2	8
[F] ANY DARK GREEN, LEAFY VEGETABLES?	Dark green, leafy vegetables	1	2	8
[G] RIPE MANGOES, PAPAYAS?	Ripe mangoes or papayas	1	2	8
[H] ANY OTHER FRUITS OR VEGETABLES?	Other fruits or vegetables	1	2	8
[I] LIVER, KIDNEY, HEART OR OTHER ORGAN MEATS?	Liver, kidney, heart or other organ meats	1	2	8
[J] ANY MEAT, SUCH AS BEEF, PORK, LAMB, GOAT, CHICKEN, OR DUCK?	Meat, such as beef, pork, lamb, goat, etc.	1	2	8
[K] EGGS?	Eggs	1	2	8
[L] FRESH OR DRIED FISH OR SHELLFISH?	Fresh or dried fish	1	2	8
[M] ANY FOODS MADE FROM BEANS, PEAS, LENTILS, OR NUTS?	Foods made from beans, peas, etc.	1	2	8
[N] CHEESE OR OTHER FOOD MADE FROM MILK?	Cheese or other food made from milk	1	2	8
[O] ANY OTHER SOLID, SEMI-SOLID, OR SOFT FOOD THAT I HAVE NOT MENTIONED (<i>specify</i>)?	(<i>Specify</i>) _____	1	2	8
<p>BD9. Check BD8 (Categories "A" through "O")</p> <p><input type="checkbox"/> At least one "Yes" or all "DK" ⇒ Go to BD11</p> <p><input type="checkbox"/> Else ⇒ Continue with BD10</p>				
<p>BD10. Probe to determine whether the child ate any solid, semi-solid or soft foods yesterday during the day or night</p> <p><input type="checkbox"/> The child did not eat or the respondent does not know ⇒ Go to Next Module</p> <p><input type="checkbox"/> The child ate at least one solid, semi-solid or soft food item mentioned by the respondent ⇒ Go back to BD8 and record food eaten yesterday [A to O]. When finished, continue with BD11</p>				
<p>BD11. HOW MANY TIMES DID (<i>name</i>) EAT ANY SOLID, SEMI-SOLID OR SOFT FOODS YESTERDAY DURING THE DAY OR NIGHT?</p> <p><i>If 7 or more times, record '7'.</i></p>		<p>Number of times.....__</p> <p>DK.....8</p>		

CARE OF ILLNESS		CA
<p>CA1. IN THE LAST TWO WEEKS, HAS (<i>name</i>) HAD DIARRHOEA?</p>	Yes 1 No 2 DK..... 8	2⇒CA6A 8⇒CA6A
<p>CA2. I WOULD LIKE TO KNOW HOW MUCH (<i>name</i>) WAS GIVEN TO DRINK DURING THE DIARRHOEA (INCLUDING BREAST MILK).</p> <p>DURING THE TIME (<i>name</i>) HAD DIARRHOEA, WAS HE/SHE GIVEN LESS THAN USUAL TO DRINK, ABOUT THE SAME AMOUNT, OR MORE THAN USUAL?</p> <p><i>If 'less', probe:</i> WAS HE/SHE GIVEN MUCH LESS THAN USUAL TO DRINK, OR SOMEWHAT LESS?</p>	Much less 1 Somewhat less 2 About the same 3 More 4 Nothing to drink 5 DK..... 8	
<p>CA3. DURING THE TIME (<i>name</i>) HAD DIARRHOEA, WAS HE/SHE GIVEN LESS THAN USUAL TO EAT, ABOUT THE SAME AMOUNT, MORE THAN USUAL, OR NOTHING TO EAT?</p> <p><i>If 'less', probe:</i> WAS HE/SHE GIVEN MUCH LESS THAN USUAL TO EAT OR SOMEWHAT LESS?</p>	Much less 1 Somewhat less 2 About the same 3 More 4 Stopped food 5 Never gave food 6 DK..... 8	
<p>CA3A. DID YOU SEEK ANY ADVICE OR TREATMENT FOR THE DIARRHOEA FROM ANY SOURCE?</p>	Yes 1 No 2 DK..... 8	2⇒CA4 8⇒CA4
<p>CA3B. FROM WHERE DID YOU SEEK ADVICE OR TREATMENT?</p> <p><i>Probe:</i> ANYWHERE ELSE?</p> <p><i>Circle all providers mentioned, but do NOT prompt with any suggestions.</i></p> <p><i>Probe to identify each type of source.</i></p> <p><i>If unable to determine if public or private sector, write the name of the place.</i></p> <p>_____</p> <p>(<i>Name of place</i>)</p>	Public sector Government hospital A Government health centre B Government dispensary C Community health worker D Mobile / Outreach clinic E Other public (<i>specify</i>) _____ H Private medical sector Private hospital / clinic I Private physician J Private pharmacy K Mobile clinic L Mission hospital / clinic M Other private medical (<i>specify</i>) _____ O Other source Relative / Friend P Shop Q Traditional practitioner R Other (<i>specify</i>) _____ X	

<p>CA4. DURING THE TIME (<i>name</i>) HAD DIARRHOEA, WAS (<i>name</i>) GIVEN TO DRINK :</p> <p>[A] A FLUID MADE FROM A SPECIAL PACKET CALLED ORS?</p> <p>[B] A PRE-PACKAGED ORS FLUID FOR DIARRHOEA?</p>	<p style="text-align: right;">Y N DK</p> <p>Fluid from ORS packet 1 2 8</p> <p>Pre-packaged ORS fluid 1 2 8</p>	
<p>CA4A. Check CA4: ORS</p> <p><input type="checkbox"/> Child was given ORS ('Yes' circled in 'A' or 'B' in CA4) ⇒ Continue with CA4B</p> <p><input type="checkbox"/> Child was not given ORS ⇒ Go to CA4C</p>		

<p>CA4B. WHERE DID YOU GET THE ORS?</p> <p><i>Probe to identify the type of source.</i></p> <p><i>If unable to determine whether public or private, write the name of the place.</i></p> <p>_____</p> <p>(Name of place)</p>	<p>Public sector</p> <p>Government hospital 11</p> <p>Government health centre 12</p> <p>Government dispensary 13</p> <p>Community health worker 14</p> <p>Mobile / Outreach clinic 15</p> <p>Other public (<i>specify</i>) 16</p> <p>Private medical sector</p> <p>Private hospital / clinic 21</p> <p>Private physician 22</p> <p>Private pharmacy 23</p> <p>Mobile clinic 24</p> <p>Mission hospital / clinic 25</p> <p>Other private medical (<i>specify</i>) 26</p> <p>Other source</p> <p>Relative / Friend 31</p> <p>Shop 32</p> <p>Traditional practitioner 33</p> <p>Already had at home 40</p> <p>Other (<i>specify</i>) 96</p>	
<p>CA4C. DURING THE TIME (name) HAD DIARRHOEA, WAS (name) GIVEN:</p> <p>[A] ZINC TABLETS?</p> <p>[B] ZINC SYRUP?</p>	<p style="text-align: right;">Y N DK</p> <p>Zinc tablets 1 2 8</p> <p>Zinc syrup 1 2 8</p>	
<p>CA4D. Check CA4C: Any zinc?</p> <p><input type="checkbox"/> Child given any zinc ('Yes' circled in 'A' or 'B' in CA4C) ⇒ Continue with CA4E</p> <p><input type="checkbox"/> Child was not given any zinc' ⇒ Go to CA4F</p>		
<p>CA4E. WHERE DID YOU GET THE ZINC?</p> <p><i>Probe to identify the type of source.</i></p> <p><i>If unable to determine whether public or private, write the name of the place.</i></p> <p>_____</p> <p>(Name of place)</p>	<p>Public sector</p> <p>Government hospital 11</p> <p>Government health centre 12</p> <p>Government dispensary 13</p> <p>Community health worker 14</p> <p>Mobile / Outreach clinic 15</p> <p>Other public (<i>specify</i>) 16</p> <p>Private medical sector</p> <p>Private hospital / clinic 21</p> <p>Private physician 22</p> <p>Private pharmacy 23</p> <p>Mobile clinic 24</p> <p>Mission hospital / clinic 25</p> <p>Other private medical (<i>specify</i>) 26</p> <p>Other source</p> <p>Relative / Friend 31</p> <p>Shop 32</p> <p>Traditional practitioner 33</p> <p>Already had at home 40</p> <p>Other (<i>specify</i>) 96</p>	

<p>CA4F. DURING THE TIME (<i>name</i>) HAD DIARRHOEA, WAS (<i>name</i>) GIVEN TO DRINK ANY OF THE FOLLOWING:</p> <p><i>Read each item aloud and record response before proceeding to the next item.</i></p> <p>[A] CEREAL GRUEL (UJI)?</p> <p>[B] FRESH OR FERMENTED MILK?</p> <p>[C] FRESH FRUIT JUICES?</p> <p>[D] SOUPS PREPARED FROM MEAT, FISH AND CHICKEN?</p> <p>[E] CLEAN, SAFE WATER?</p> <p>[F] BREAST FEEDING?</p>	<p style="text-align: right;">Y N DK</p> <p>Cereal gruel (uji) 1 2 8</p> <p>Fresh or fermented milk 1 2 8</p> <p>Fresh fruit juices 1 2 8</p> <p>Soups 1 2 8</p> <p>Clean, Safe water 1 2 8</p> <p>Breast feeding 1 2 8</p>	
<p>CA5. WAS ANYTHING (ELSE) GIVEN TO TREAT THE DIARRHOEA?</p>	<p>Yes 1</p> <p>No 2</p> <p>DK 8</p>	<p>2⇒CA6A</p> <p>8⇒CA6A</p>
<p>CA6. WHAT (ELSE) WAS GIVEN TO TREAT THE DIARRHOEA?</p> <p><i>Probe:</i> ANYTHING ELSE?</p> <p><i>Record all treatments given. Write brand name(s) of all medicines mentioned.</i></p> <p>_____</p> <p style="text-align: center;">(<i>Name</i>)</p>	<p>Pill or Syrup</p> <p>Antibiotic A</p> <p>Antimotility B</p> <p>Other pill or syrup (Not antibiotic, antimotility or zinc) G</p> <p>Unknown pill or syrup H</p> <p>Injection</p> <p>Antibiotic L</p> <p>Non-antibiotic M</p> <p>Unknown injection N</p> <p>Intravenous O</p> <p>Home remedy/Herbal medicine Q</p> <p>Other (<i>specify</i>) _____ X</p>	
<p>CA6A. IN THE LAST TWO WEEKS, HAS (<i>name</i>) BEEN ILL WITH A FEVER AT ANY TIME?</p>	<p>Yes 1</p> <p>No 2</p> <p>DK 8</p>	<p>2⇒CA7</p> <p>8⇒CA7</p>
<p>CA6B. AT ANY TIME DURING THE ILLNESS, DID (<i>name</i>) HAVE BLOOD TAKEN FROM HIS/HER FINGER OR HEEL FOR TESTING?</p>	<p>Yes 1</p> <p>No 2</p> <p>DK 8</p>	
<p>CA7. AT ANY TIME IN THE LAST TWO WEEKS, HAS (<i>name</i>) HAD AN ILLNESS WITH A COUGH?</p>	<p>Yes 1</p> <p>No 2</p> <p>DK 8</p>	<p>2⇒CA9A</p> <p>8⇒CA9A</p>
<p>CA8. WHEN (<i>name</i>) HAD AN ILLNESS WITH A COUGH, DID HE/SHE BREATHE FASTER THAN</p>	<p>Yes 1</p> <p>No 2</p>	<p>2⇒CA10</p>

USUAL WITH SHORT, RAPID BREATHS OR HAVE DIFFICULTY BREATHING?	DK.....8	8⇒CA10
CA9. WAS THE FAST OR DIFFICULT BREATHING DUE TO A PROBLEM IN THE CHEST OR A BLOCKED OR RUNNY NOSE?	Problem in chest only 1 Blocked or runny nose only 2 Both 3 Other (<i>specify</i>) 6 DK..... 8	1⇒CA10 2⇒CA10 3⇒CA10 6⇒CA10 8⇒CA10
CA9A. Check CA6A: Had fever?		
<input type="checkbox"/> Child had fever ⇒ Continue with CA10 <input type="checkbox"/> Child did not have fever ⇒ Go to CA14		
CA10. DID YOU SEEK ANY ADVICE OR TREATMENT FOR THE ILLNESS FROM ANY SOURCE?	Yes 1 No 2 DK..... 8	2⇒CA12 8⇒CA12
CA11. FROM WHERE DID YOU SEEK ADVICE OR TREATMENT? <i>Probe:</i> ANYWHERE ELSE? <i>Circle all providers mentioned, but do NOT prompt with any suggestions.</i> <i>Probe to identify each type of source.</i> <i>If unable to determine if public or private sector, write the name of the place.</i> _____ (<i>Name of place</i>)	Public sector Government hospital A Government health centre B Government dispensary C Community health worker D Mobile / Outreach clinic E Other public (<i>specify</i>) F Private medical sector Private hospital / clinic G Private physician H Private pharmacy I Mobile clinic J Mission hospital / clinic K Other source Relative / Friend L Shop M Traditional practitioner N Other (<i>specify</i>) X	
CA12. AT ANY TIME DURING THE ILLNESS, WAS (<i>name</i>) GIVEN ANY MEDICINE FOR THE ILLNESS?	Yes 1 No 2 DK..... 8	2⇒CA14 8⇒CA14
CA13. WHAT MEDICINE WAS (<i>name</i>) GIVEN? <i>Probe:</i> ANY OTHER MEDICINE? <i>Circle all medicines given. Write brand name(s) of all medicines mentioned.</i> _____ (<i>Names of medicines</i>)	Anti-malarials: SP / Fansidar A Chloroquine B Amodiaquine C Quinine D Combination with Artemisinin E Other anti-malarial (<i>specify</i>) H Antibiotics: Pill / Syrup I Injection J	

	Other medications: Paracetamol/ Panadol /Acetaminophen . P Aspirin..... Q Ibuprofen R Other (<i>specify</i>) X DK.....Z	
CA13A. Check CA13: Antibiotic mentioned (codes I or J)? <input type="checkbox"/> Yes ⇒ Continue with CA13B <input type="checkbox"/> No ⇒ Go to CA13C		
CA13B. WHERE DID YOU GET THE ANTIBIOTICS? <i>Probe to identify the type of source.</i> <i>If unable to determine whether public or private, write the name of the place.</i> _____ (<i>Name of place</i>)	Public sector Government hospital 11 Government health centre 12 Government dispensary 13 Community health worker..... 14 Mobile / Outreach clinic 15 Other public (<i>specify</i>) 16 Private medical sector Private hospital / clinic 21 Private physician 22 Private pharmacy 23 Mobile clinic 24 Mission hospital /clinic..... 25 Other private medical (<i>specify</i>) 26 Other source Relative / Friend 31 Shop 32 Traditional practitioner 33 Already had at home 40 Other (<i>specify</i>) 96	
CA13C. Check CA13: Anti-malarial mentioned (codes A - H)? <input type="checkbox"/> Yes ⇒ Continue with CA13D <input type="checkbox"/> No ⇒ Go to CA14		
CA13D. WHERE DID YOU GET THIS ANTI-MALARIAL? <i>Probe to identify the type of source.</i> <i>If unable to determine whether public or private, write the name of the place.</i> _____ (<i>Name of place</i>)	Public sector Government hospital 11 Government health centre 12 Government dispensary 13 Community health worker..... 14 Mobile / Outreach clinic 15 Other public (<i>specify</i>) 16 Private medical sector Private hospital / clinic 21 Private physician 22 Private pharmacy 23 Mobile clinic 24 Mission hospital /clinic..... 25 Other private medical (<i>specify</i>) 26 Other source	

	Relative / Friend31 Shop32 Traditional practitioner33 Already had at home40 Other (<i>specify</i>) _____ 96	
CA13E. HOW LONG AFTER THE FEVER STARTED DID (name) FIRST TAKE (name of anti-malarial from CA13)? <i>If multiple anti-malarials mentioned in CA13, name all anti-malarial medicines mentioned.</i>	Same day0 Next day1 2 days after the fever.....2 3 days after the fever.....3 4 or more days after the fever4 DK.....8	
CA14. Check AG2: Age of child <input type="checkbox"/> Child age 0, 1 or 2 ⇒ Continue with CA15 <input type="checkbox"/> Child age 3 or 4 ⇒ Go to UF13		
CA15. THE LAST TIME (name) PASSED STOOLS, WHAT WAS DONE TO DISPOSE OF THE STOOLS?	Child used toilet/latrine01 Put / Rinsed into toilet or latrine02 Put / Rinsed into drain or ditch03 Thrown into garbage (solid waste)04 Buried05 Left in the open.....06 Other (<i>specify</i>) _____ 96 DK.....98	

UF13. Record the time.	Hour and minutes__ : __	
-------------------------------	-------------------------------	--

UF14. Check List of Household Members, columns HL7B and HL15. <i>Is the respondent the mother or caretaker of another child age 0-4 living in this household?</i> <input type="checkbox"/> Yes ⇒ Indicate to the respondent that you will need to measure the weight and height of the child later. Go to the next QUESTIONNAIRE FOR CHILDREN UNDER FIVE to be administered to the same respondent <input type="checkbox"/> No ⇒ End the interview with this respondent by thanking her/him for her/his cooperation and tell her/him that you will need to measure the weight and height of the child before you leave the household <i>Check to see if there are other woman's, man's or under-5 questionnaires to be administered in this household.</i>
--

ANTHROPOMETRY		AN
<p><i>After questionnaires for all children are complete, the measurer weighs and measures each child. Record weight and length/height below, taking care to record the measurements on the correct questionnaire for each child. Check the child's name and line number in the List of Household Members before recording measurements.</i></p>		
AN1. <i>Measurer's name and number:</i>	Name _____	
AN2. <i>Result of height/length and weight measurement</i>	Either or both measured..... 1	
	Child not present 2	2⇒AN6
	Child or mother/caretaker refused..... 3	3⇒AN6
	Other (specify) _____ 6	6⇒AN6
AN3. <i>Child's weight</i>	Kilograms (kg) _ . _	
	Weight not measured 99.9	
AN3A. <i>Was the child undressed to the minimum?</i>		
<input type="checkbox"/> Yes <input type="checkbox"/> No, the child could not be undressed to the minimum		
AN3B. <i>Check age of child in AG2:</i>		
<input type="checkbox"/> Child under 2 years old. ⇒ Measure length (lying down). <input type="checkbox"/> Child age 2 or more years. ⇒ Measure height (standing up).		
AN4. <i>Child's length or height</i>	Length / Height (cm)..... _ . _	
	Length/ Height not measured..... 999.9	⇒AN6
AN4A. <i>How was the child actually measured? Lying down or standing up?</i>	Lying down 1	
	Standing up 2	

AN6. <i>Is there another child in the household who is eligible for measurement?</i>		
<input type="checkbox"/> Yes ⇒ Record measurements for next child. <input type="checkbox"/> No ⇒ Check if there are any other individual questionnaires to be completed in the household.		

Interviewer's Observations

Field Editor's Observations

Supervisor's Observations

Measurer's Observations